

THE SCHRÖDINGER-POISSON SYSTEM ON THE SPHERE

PATRICK GÉRARD AND FLORIAN MÉHATS

In memory of Naoufel Ben Abdallah, with our friendship

ABSTRACT. We study the Schrödinger-Poisson system on the unit sphere \mathbb{S}^2 of \mathbb{R}^3 , modeling the quantum transport of charged particles confined on a sphere by an external potential. Our first results concern the Cauchy problem for this system. We prove that this problem is regularly well-posed on every $H^s(\mathbb{S}^2)$ with $s > 0$, and not uniformly well-posed on $L^2(\mathbb{S}^2)$. The proof of well-posedness relies on multilinear Strichartz estimates, the proof of ill-posedness relies on the construction of a counterexample which concentrates exponentially on a closed geodesic. In a second part of the paper, we prove that this model can be obtained as the limit of the three dimensional Schrödinger-Poisson system, singularly perturbed by an external potential that confines the particles in the vicinity of the sphere.

1. Introduction and main results

Let $\mathbb{S}^2 \subset \mathbb{R}^3$ be the unit sphere. For functions defined on \mathbb{S}^2 , one considers the operator G defined by

$$G(f)(x) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \frac{1}{|x-y|} f(y) d\sigma(y), \quad (1.1)$$

where σ denote the surface measure on \mathbb{S}^2 and $|\cdot|$ is the Euclidean norm on \mathbb{R}^3 .

In this paper, we are interested in the following Schrödinger-Poisson system on \mathbb{S}^2 :

$$i\partial_t u + \Delta_\sigma u = G(|u|^2)u, \quad u(t=0) = u_0. \quad (1.2)$$

Here Δ_σ denotes the Laplace-Beltrami operator on \mathbb{S}^2 . This system models the transport of a gas of quantum charged particles confined on a surface, here the sphere, and interacting through the Poisson potential, which is the Coulombian interaction in the Hartree approximation. Our purpose in studying this ideal system is twofold.

First, we are interested in understanding the wellposedness of the Cauchy problem for (1.2): choice of the function space, local in time or global in time solutions, stability of the flow map, ... From this point of view, this paper comes in the continuity of several recent works concerning the nonlinear Schrödinger equation on Riemannian manifolds or in inhomogeneous media [9, 10, 11, 12, 13, 19, 20].

Second, we wish to justify this system for modeling a quantum gas via some asymptotic analysis, starting from a more conventional 3D Schrödinger-Poisson system with a singular perturbation which stands for a strong confinement potential. Strongly confined Schrödinger-Poisson systems have previously been studied in Euclidean spaces: in [5, 16] for the confinement on a plane and in [3] for the confinement on an axis. The idea here is to investigate the influence of the geometry on

the confinement procedure. The case of the sphere can be seen as a step before more general manifolds, which can be interesting in some applications in the field of nanoelectronics. Quantum dynamical systems confined on a surface have been studied previously in [15, 17, 28] in linear situations. Starting from a similar scaling on the transverse Hamiltonian, these authors consider the linear Schrödinger equation with a confinement on a general surface and derive an effective Hamiltonian which locally depends on the curvature properties of the surface. Here our approach is mainly concentrated on understanding the nonlinear effects.

Our first result states that this problem is locally well-posed in H^s , $s > 0$ and globally well-posed in the energy space:

Theorem 1.1. *Let $s > 0$ a real number. For every bounded subset $\mathcal{B} \subset H^s(\mathbb{S}^2)$, there exists $T \in (0, +\infty]$ and a subspace X_T of $\mathcal{C}((-T, T), H^s(\mathbb{S}^2))$ such that, for $u_0 \in \mathcal{B}$, the Cauchy problem (1.2) admits a unique solution $u \in X_T$. For all $0 < T' < T$, the application $u_0 \mapsto u \in C([-T', T'], H^s(\mathbb{S}^2))$ is Lipschitz continuous on \mathcal{B} . Moreover, if $s \geq 1$ one can choose $T = +\infty$ and this global solution satisfies the following two conservation laws:*

$$\|u(t)\|_{L^2} = Q_0, \quad \|\nabla_\sigma u(t)\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{S}^2} G(|u|^2)|u|^2 d\sigma = E_0. \quad (1.3)$$

Let us now deal with the limit case $s = 0$: after Theorem 1.1, the question whether this system is well-posed on $L^2(\mathbb{S}^2)$ is natural. In the case of a plane, the answer is positive thanks to Strichartz estimates in dimension 2. Our second result is that, in the case of the sphere, this system is not locally (uniformly) well-posed on \mathbb{S}^2 . The precise statement is as follows.

Theorem 1.2. *For all ball \mathcal{B} of $L^2(\mathbb{S}^2)$ centered on 0 and for all $T > 0$, the application $u_0 \mapsto u$ is not uniformly continuous from $\mathcal{B} \cap H^1(\mathbb{S}^2)$, endowed with the L^2 norm, into $C([-T, T], L^2(\mathbb{S}^2))$.*

Our third result concerns the asymptotic analysis. The starting model is the Schrödinger-Poisson system (or Hartree equation) in \mathbb{R}^3 with a strong potential that confines the particles near the sphere. We consider the following system, on the unknown $u^\varepsilon(t, x)$:

$$i\partial_t u^\varepsilon = -\Delta u^\varepsilon + V_c^\varepsilon u^\varepsilon + V(|u^\varepsilon|^2)u^\varepsilon, \quad u^\varepsilon(t=0) = u_0^\varepsilon, \quad (1.4)$$

where V_c^ε and $V(|u|^2)$ are respectively the applied confinement potential and the selfconsistent Poisson potential, given by

$$V_c^\varepsilon(x) = \frac{1}{\varepsilon^2} V_c \left(\frac{|x| - 1}{\varepsilon} \right) \quad (1.5)$$

and

$$V(|u|^2)(x) = \frac{1}{4\pi} \int \frac{1}{|x - x'|} |u(x')|^2 dx'. \quad (1.6)$$

The parameter $\varepsilon > 0$ is the order of magnitude of the width of the confined gas, compared to a typical length, for instance the radius of the sphere. The square of ε also denotes the ratio between the kinetic energy of the particles and the confinement energy. One refers to [5] and [16], where similar singularly perturbed systems were derived and put in dimensionless form, starting from systems in physical variables. Our aim is to derive a simpler asymptotic system satisfied by u^ε as $\varepsilon \rightarrow 0$.

The function V_c is fixed, independent of ε , and is supposed to go to infinity at the infinity, faster than the harmonic potential, as stated in the following assumption. Note that we do not solve the asymptotic problem in the precise case of a harmonic confinement potential, due to a lack of a priori estimates (Assumption $\alpha > 2$ is used several times in the proofs of Lemma 3.1 and Proposition 3.5).

Assumption 1.3. *The confinement potential is a C^∞ function such that*

$$\forall z \in \mathbb{R} \quad V_c(z) \geq C|z|^\alpha, \quad (1.7)$$

where $\alpha > 2$. Moreover, it satisfies the following condition:

$$\forall k \in \mathbb{N}, \quad \exists C_k > 0 : \forall z \in \mathbb{R} \quad \left| \frac{\partial^k V_c}{\partial z^k}(z) \right| \leq C_k V_c(z). \quad (1.8)$$

Let us define the energy adapted to our system. We set

$$\mathcal{B}^1 = \left\{ u \in \dot{H}^1(\mathbb{R}^3) : (V_c^\varepsilon)^{1/2} u \in L^2(\mathbb{R}^3) \right\},$$

equipped with the norm

$$\|u\|_{\mathcal{B}^1}^2 = \|\nabla u\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} u\|_{L^2}^2. \quad (1.9)$$

The two conservations laws associated to (1.4) are the conservation of mass and the conservation of energy:

$$\|u^\varepsilon(t)\|_{L^2}^2 = \|u_0^\varepsilon\|_{L^2}^2 \quad (1.10)$$

$$\begin{aligned} & \|\nabla u^\varepsilon(t)\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} u^\varepsilon(t)\|_{L^2}^2 + \frac{1}{2} \|\nabla V(|u^\varepsilon(t)|^2)\|_{L^2}^2 \\ &= \|\nabla u_0^\varepsilon\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} u_0^\varepsilon\|_{L^2}^2 + \frac{1}{2} \|\nabla V(|u_0^\varepsilon|^2)\|_{L^2}^2. \end{aligned}$$

The scaling (1.5) of the confinement potential suggests that, if $\|u_0^\varepsilon\|_{L^2}^2$ is of order 1, it will be natural to start with an energy of order $\frac{1}{\varepsilon^2}$. Consequently, the solution of (1.4) will satisfy the following natural bounds:

$$\|u^\varepsilon\|_{L^2} \leq C, \quad \varepsilon \|u^\varepsilon\|_{\mathcal{B}^1} \leq C.$$

From Theorem 1.2, one can guess that these bounds will not be sufficient in order to analyze the limit of u as ε goes to zero. Consequently, we shall assume some regularity of the initial data with respect to the angular variable σ . To be more precise, let us introduce the spherical coordinates $(r, \sigma) \in \mathbb{R}_+^* \times \mathbb{S}^2$ defined for all $x \in \mathbb{R}^3 \setminus \{0\}$ by $r = |x|$ and $\sigma = \frac{x}{|x|}$. We recall that the Laplace operator has the following form in spherical coordinates:

$$\Delta = \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2} \Delta_\sigma.$$

In particular, a remarkable property is that this operator commutes with Δ_σ . This property will be crucial in our nonlinear analysis. In the sequel, we shall set $\langle \nabla_\sigma \rangle = (1 - \Delta_\sigma)^{1/2}$. We introduce a last notation. The transversal confinement operator is the following operator:

$$H_r = -\frac{1}{r^2} \partial_r (r^2 \partial_r) + V_c^\varepsilon(r), \quad (1.11)$$

with domain

$$D(H_r) = \left\{ u \in L^2(\mathbb{R}_+, r^2 dr) : \frac{1}{r^2} \partial_r (r^2 \partial_r u) \in L^2(\mathbb{R}_+, r^2 dr), V_c^\varepsilon u \in L^2(\mathbb{R}_+, r^2 dr) \right\}.$$

Assumption 1.3 implies that H_r is self-adjoint on $L^2(\mathbb{R}_+, r^2 dr)$, with a compact resolvent. We shall denote by $(E_p^\varepsilon)_{p \in \mathbb{N}}$, $(\psi_p^\varepsilon)_{p \in \mathbb{N}}$ its eigenvalues and its eigenfunctions.

We make the following assumption on the initial data.

Assumption 1.4. *The initial data u_0^ε satisfies*

$$\|\langle \nabla_\sigma \rangle u_0^\varepsilon\|_{L^2} + \varepsilon \|u_0^\varepsilon\|_{\mathcal{B}^1} \leq C, \quad (1.12)$$

where $C > 0$ is independent of $\varepsilon \in (0, 1]$. Moreover, we have

$$\limsup_{\varepsilon \rightarrow 0} (\|\mathbb{1}_{\varepsilon^2 H_r > R} \langle \nabla_\sigma \rangle u_0^\varepsilon\|_{L^2} + \|\mathbb{1}_{\langle \nabla_\sigma \rangle > R} \langle \nabla_\sigma \rangle u_0^\varepsilon\|_{L^2}) \rightarrow 0 \text{ as } R \rightarrow +\infty. \quad (1.13)$$

Let us comment on these assumptions. In (1.12), the L^2 bound of $\langle \nabla_\sigma \rangle u_0$ means a supplementary decay at the infinity for this quantity, compared to a H^1 norm which would only control the L^2 norm of $\frac{1}{r} \langle \nabla_\sigma \rangle u_0^\varepsilon$. The second assumption means that $\langle \nabla_\sigma \rangle u_0^\varepsilon$ is partially relatively compact in L^2 with respect to the σ variable ε^2 -oscillatory with respect to the operator H_r defined by (1.11) (see [18] for an abstract definition of ε -oscillatory sequences with respect to a self-adjoint operator).

Our third theorem is the following one.

Theorem 1.5. *Under Assumptions 1.3 and 1.4 on the confinement potential and the initial data, for all $\varepsilon > 0$, (1.4) admits a unique global solution $u^\varepsilon \in \mathcal{C}(\mathbb{R}, L^2 \cap \mathcal{B}_1)$. Moreover, the following equation admits a unique global solution $v^\varepsilon \in \mathcal{C}^0(\mathbb{R}, \mathcal{B}^1)$ such that $\langle \nabla_\sigma \rangle v^\varepsilon \in \mathcal{C}^0(\mathbb{R}, L^2)$:*

$$i\partial_t v^\varepsilon = H_r v^\varepsilon - \Delta_\sigma v^\varepsilon + \mathbb{G}(|v^\varepsilon|^2) v^\varepsilon, \quad v^\varepsilon(t=0) = u_0^\varepsilon, \quad (1.14)$$

where \mathbb{G} is the following generalization of the operator G defined in (1.1): for functions defined on \mathbb{R}^3 ,

$$\mathbb{G}(f)(r\sigma) = \frac{1}{4\pi} \int_0^{+\infty} \int_{\mathbb{S}^2} \frac{1}{|\sigma - \sigma'|} f(r'\sigma') r'^2 dr' d\sigma'. \quad (1.15)$$

Finally, we have the approximation of (1.4) by the limit system (1.14): for all $T > 0$,

$$\lim_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (u^\varepsilon - v^\varepsilon)\|_{L^\infty([-T, T], L^2)} = 0.$$

Notice that, in (1.14), the nonlinear Schrödinger dynamics operates only in the σ variable. Indeed, the function ω^ε defined by

$$\omega^\varepsilon(t, r, \sigma) = e^{itH_r} v^\varepsilon(t, r, \sigma)$$

solves the system

$$i\partial_t \omega^\varepsilon = -\Delta_\sigma \omega^\varepsilon + \mathbb{G}(|\omega^\varepsilon|^2) \omega^\varepsilon, \quad \omega^\varepsilon(t=0) = u_0^\varepsilon, \quad (1.16)$$

which is a "mixed-state" version of the scalar equation (1.2). Indeed, let us decompose the function ω^ε on the eigenvectors of the confinement operator H_r , setting

$$\omega^\varepsilon(t, r, \sigma) = \sum_{p=0}^{+\infty} \omega_p^\varepsilon(t, \sigma) \psi_p^\varepsilon(r),$$

i.e.

$$v^\varepsilon(t, r, \sigma) = \sum_{p=0}^{+\infty} e^{-itE_p^\varepsilon} \omega_p^\varepsilon(t, \sigma) \psi_p^\varepsilon(r).$$

Then each component ω_p^ε satisfies

$$i\partial_t\omega_p^\varepsilon = -\Delta_\sigma\omega_p^\varepsilon + G\left(\sum_{q=0}^{+\infty}|\omega_q^\varepsilon|^2\right)\omega_p^\varepsilon, \quad \omega_p^\varepsilon(t=0) = \int_0^{+\infty} u_0 \psi_p^\varepsilon r^2 dr.$$

Notice that the components ω_p^ε are only coupled through the selfconsistent potential. In particular, if the initial data is polarized on a single eigenmode, i.e. $u_0^\varepsilon = v_{p_0}\psi_{p_0}^\varepsilon$, then the solution of (1.14) remains polarized on the same mode p_0 , for all time: $v^\varepsilon(t, r, \sigma) = e^{-itE_{p_0}^\varepsilon/\varepsilon^2}\omega_{p_0}^\varepsilon(t, \sigma)\psi_{p_0}^\varepsilon(r)$ and $\omega_{p_0}^\varepsilon$ satisfies (1.2).

2. The Schrödinger-Poisson system on the sphere

This section is devoted to the analysis of the Schrödinger-Poisson system (1.2) on the two-dimensional sphere \mathbb{S}^2 . As announced in the introduction, we will prove that this system is locally well-posed in $H^s(\mathbb{S}^2)$ for all $s > 0$, but not for the critical case $L^2(\mathbb{S}^2)$. To make precise this statement, let us recall the notion of well-posedness that is meant here.

Definition 2.1. *Let $s \in \mathbb{R}$. We shall say that the equation (1.2) is, locally in time, uniform well-posed in $H^s(\mathbb{S}^2)$ if, for any bounded subset B of $H^s(\mathbb{S}^2)$, there exists $T > 0$ and a Banach space X_T continuously embedded into $C([-T, T], H^s(\mathbb{S}^2))$, such that*

- (i) *For every Cauchy data $u_0 \in B$, (1.2) admits a unique solution $u \in X_T$,*
- (ii) *If $u_0 \in H^\sigma(\mathbb{S}^2)$ for $\sigma > s$, then $u \in C([-T, T], H^\sigma(\mathbb{S}^2))$,*
- (iii) *The map $u_0 \in B \mapsto u \in X_T$ is uniformly continuous.*

The proof of Theorem 1.1 uses essentially the techniques developed in the works of Burq, Gérard, Tzvetkov [9, 10, 11, 12]. It can be decomposed into two main steps: the proof of well-posedness under the assumption of a multilinear estimate and the proof of the multilinear estimate itself. More precisely, we follow the work of Gérard and Pierfelice [20], who have adapted the methodology to the particular structure of Hartree-type nonlinearities as in (1.2). The analysis relies in a crucial way on a quadrilinear estimate. For the sake of completeness and readability, we sketch all the steps of this proof, although the new contribution in this proof essentially concerns the quadrilinear estimate proved in subsection 2.2.

More precisely, Theorem 1.1 can be seen as a direct corollary of the following Propositions 2.2 and 2.7, completed with the conservation laws (1.3) (which are very classical and will not be proved here).

2.1. Well-posedness via quadrilinear estimate. The following proposition reduces the study of wellposedness for (1.2) to a quadrilinear estimate.

Proposition 2.2. *Suppose that, for every $\chi \in \mathcal{C}_0^\infty(\mathbb{R})$ and for all $\varepsilon \in (0, \frac{1}{2})$, there exists $C_\varepsilon > 0$ such that, for any $f_1, f_2, f_3, f_4 \in L^2(\mathbb{S}^2)$ satisfying*

$$\mathbb{1}_{\sqrt{1-\Delta_\sigma} \in [N_j, 2N_j]}(f_j) = f_j, \quad j = 1, 2, 3, 4, \quad (2.1)$$

one has the following quadrilinear estimate:

$$\sup_{\tau \in \mathbb{R}} \left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} \chi(t) e^{it\tau} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| \leq C_\varepsilon (m(N_1, N_2, N_3, N_4))^\varepsilon \prod_{i=1}^4 \|f_j\|_{L^2(\mathbb{S}^2)} \quad (2.2)$$

where $u_j(t) = S(t)f_j$ for $j = 1, 2, 3, 4$, $S(t) = e^{it\Delta_\sigma}$ denoting the free evolution, and where $m(N_1, N_2, N_3, N_4)$ denotes the product of the smallest two numbers among N_1, N_2, N_3, N_4 . Then the Cauchy problem (1.2) is locally uniformly well-posed in $H^s(\mathbb{S}^2)$ for any $s > 0$.

Proof. We follow the steps of the proof of Theorem 1 in [10], see also [12, 20]. Proposition 2.2 will be proved thanks to a fixed point procedure formulated in the Bourgain spaces $X^{s,b}$.

Step 1. Reformulation of the problem in the Bourgain spaces.

Following the definitions in [7] and [12], we introduce the following family of Hilbert spaces

$$X^{s,b} = \left\{ u \in \mathcal{S}'(\mathbb{R} \times \mathbb{S}^2) : (1 + |i\partial_t + \Delta_\sigma|^2)^{b/2} (1 - \Delta_\sigma)^{s/2} u \in L^2(\mathbb{R} \times \mathbb{S}^2) \right\}, \quad (2.3)$$

for $s, b \in \mathbb{R}$. For any $T > 0$, we also denote by $X_T^{s,b}$ the space of restrictions of elements of $X^{s,b}$ to $(-T, T) \times \mathbb{S}^2$. We gather in the following lemma some interesting properties of these spaces.

Lemma 2.3. *The Bourgain space $X^{s,b}$ satisfies the following properties.*

- (i) $\forall f \in H^s(\mathbb{S}^2)$, $\forall b > 0$, the function $S(t)f$ belongs to $X^{s,b}$.
- (ii) $\forall b > \frac{1}{2}$, $X^{s,b} \hookrightarrow \mathcal{C}(\mathbb{R}, H^s(\mathbb{S}^2))$.
- (iii) Let b, b' such that $0 \leq b' < \frac{1}{2}$, $0 \leq b < 1 - b'$. There exists $C > 0$ such that, if $T \in (0, 1]$ and $w(t) = \int_0^t S(t - \tau) f(\tau) d\tau$, then

$$\|w\|_{X_T^{s,b}} \leq CT^{1-b-b'} \|f\|_{X_T^{s,-b'}}.$$

Proof. The first property (i) stems directly from the definition (2.3) of $X^{s,b}$. Remarking that

$$\|u\|_{X^{s,b}} = \|v\|_{H^b(\mathbb{R}, H^s(\mathbb{S}^2))}, \quad \text{where } v(t) = S(-t)u(t), \quad (2.4)$$

the second statement (ii) is obvious and the last statement (iii) appears to be a consequence of the following elementary result, proved e.g. in [21]:

$$\text{if } g(t) = \int_0^t h(\tau) d\tau \quad \text{then} \quad \|g\|_{H^b(-T, T)} \leq CT^{1-b-b'} \|f\|_{H^{-b'}(-T, T)}.$$

□

As a consequence of this lemma, it is easy to prove by a standard contraction argument that the Cauchy problem (1.2) is locally uniformly well-posed in $H^s(\mathbb{S}^2)$, $s > 0$, as soon as the following result holds true:

Lemma 2.4. *Assume that the quadrilinear estimate of Proposition 2.2 holds true. Then for all $s > 0$, one can find b, b' satisfying $0 < b' < 1/2 < b < 1 - b'$ and such that we have the estimate*

$$\|G(u_1 u_2) u_3\|_{X^{s,-b'}} \leq C \|u_1\|_{X^{s,b}} \|u_2\|_{X^{s,b}} \|u_3\|_{X^{s,b}}, \quad (2.5)$$

$$\|G(|u|^2)u\|_{X^{\sigma,-b'}} \leq C \|u\|_{X^{s,b}}^2 \|u\|_{X^{\sigma,b}} \quad \text{for } \sigma \geq s. \quad (2.6)$$

The end of this section consists in the proof of this lemma.

Step 2. Quadrilinear estimate in $X^{s,b}$.

Let us first give without proof two estimates on the operator G . These results can be obtained straightforwardly using Hardy-Littlewood-Sobolev inequalities in dimension 2.

Lemma 2.5. *The operator G defined by (1.1) satisfies the following estimates:*

$$\|G(f)\|_{L^q(\mathbb{S}^2)} \leq C \|f\|_{L^p(\mathbb{S}^2)}, \quad \text{for } 1 < p < 2 \text{ and } \frac{1}{q} = \frac{1}{p} - \frac{1}{2}, \quad (2.7)$$

$$\|G(f)\|_{L^\infty(\mathbb{S}^2)} \leq C \|f\|_{L^p(\mathbb{S}^2)}^\theta \|f\|_{L^1(\mathbb{S}^2)}^{1-\theta}, \quad \text{for } p > 2 \text{ and } \theta = \frac{p}{2p-2}. \quad (2.8)$$

We now reformulate the quadrilinear estimate (2.2) in the $X^{s,b}$ spaces.

Lemma 2.6. *Under the same assumption as in Proposition 2.2, let u_1, u_2, u_3, u_4 satisfy*

$$\mathbb{1}_{\sqrt{1-\Delta_\sigma} \in [N_j, 2N_j]}(u_j) = u_j, \quad j = 1, 2, 3, 4. \quad (2.9)$$

Then, for every $s > 0$, there exists $0 < b' < 1/2$ such that

$$\left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| \leq C m(N_1, N_2, N_3, N_4)^s \prod_{i=1}^4 \|u_i\|_{X^{0,b'}}. \quad (2.10)$$

Proof. The desired estimate (2.10) can be obtained by interpolation between the two following inequalities:

$$\left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| \leq C m(N_1, N_2, N_3, N_4)^{1/2} \prod_{i=1}^4 \|u_i\|_{X^{0,1/4}} \quad (2.11)$$

and, for any $b > 1/2$ and $0 < \varepsilon < 1/2$,

$$\left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| \leq C_\varepsilon m(N_1, N_2, N_3, N_4)^\varepsilon \prod_{i=1}^4 \|u_i\|_{X^{0,b}}. \quad (2.12)$$

Let us prove the first estimate (2.11). By symmetry and self-adjointness of G , only two cases have to be considered: $m(N_1, N_2, N_3, N_4) = N_1 N_2$ and $m(N_1, N_2, N_3, N_4) = N_1 N_3$. In the first case, we deduce from Hölder and from (2.8) that

$$\begin{aligned} \left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| &\leq \|G(u_1 u_2)\|_{L^2(\mathbb{R}, L^\infty(\mathbb{S}^2))} \|u_3 u_4\|_{L^2(\mathbb{R}, L^1(\mathbb{S}^2))} \\ &\leq C \|u_1 u_2\|_{L^2(\mathbb{R}, L^4(\mathbb{S}^2))}^{2/3} \|u_1 u_2\|_{L^2(\mathbb{R}, L^1(\mathbb{S}^2))}^{1/3} \|u_3 u_4\|_{L^2(\mathbb{R}, L^1(\mathbb{S}^2))} \\ &\leq C N_1^{1/2} N_2^{1/2} \prod_{i=1}^4 \|u_i\|_{L^4(\mathbb{R}, L^2(\mathbb{S}^2))}, \end{aligned}$$

since $\|u_i\|_{L^8(\mathbb{S}^2)} \leq C N_i^{3/4} \|u_i\|_{L^2(\mathbb{S}^2)}$ due to the spectral localization. Hence, to get (2.11) in this case, it suffices to remark that

$$\|u_j\|_{L^4(\mathbb{R}, L^2(\mathbb{S}^2))} = \|S(-t)u_j\|_{L^4(\mathbb{R}, L^2(\mathbb{S}^2))} \leq C \|S(-t)u_j\|_{H^{1/4}(\mathbb{R}, L^2(\mathbb{S}^2))} = C \|u_j\|_{X^{0,1/4}},$$

where we used the isometric action of $S(t)$, a Sobolev embedding in dimension one and (2.4).

The second case $m(N_1, N_2, N_3, N_4) = N_1 N_3$ can be treated as follows. Using (2.7), we have

$$\begin{aligned} \left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| &\leq \|G(u_1 u_2)\|_{L^2(\mathbb{R}, L^4(\mathbb{S}^2))} \|u_3 u_4\|_{L^2(\mathbb{R}, L^{4/3}(\mathbb{S}^2))} \\ &\leq C \|u_1 u_2\|_{L^2(\mathbb{R}, L^{4/3}(\mathbb{S}^2))} \|u_3 u_4\|_{L^2(\mathbb{R}, L^{4/3}(\mathbb{S}^2))} \\ &\leq C \|u_1\|_{L^4(\mathbb{R}, L^4(\mathbb{S}^2))} \|u_2\|_{L^4(\mathbb{R}, L^2(\mathbb{S}^2))} \|u_3\|_{L^4(\mathbb{R}, L^4(\mathbb{S}^2))} \|u_4\|_{L^4(\mathbb{R}, L^2(\mathbb{S}^2))} \\ &\leq C N_1^{1/2} N_3^{1/2} \prod_{i=1}^4 \|u_j\|_{L^4(\mathbb{R}, L^2(\mathbb{S}^2))} \leq C N_1^{1/2} N_2^{1/2} \prod_{i=1}^4 \|u_j\|_{X^{0,1/4}}, \end{aligned}$$

where we used $\|u_i\|_{L^4(\mathbb{S}^2)} \leq C N_i^{1/2} \|u_i\|_{L^2(\mathbb{S}^2)}$.

We now sketch the proof of the second estimate (2.12), for all $b > 1/2$, which is directly inspired from [12, 20]. Let us start by proving in a first step the result for the special case where the u_j are supported in time in an interval of size 1, say $(0, 1)$ after translation. Let $v_j(t) = S(-t)u_j(t)$ and let $f_j(\tau) = \hat{v}_j(\tau)$ denote the Fourier transform in time of v_j . Applying the inverse Fourier transform, we get

$$\begin{aligned} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma &= \frac{1}{(2\pi)^2} \iiint_{\mathbb{R}^4} e^{it(\tau_1 + \tau_2 + \tau_3 + \tau_4)} \times \\ &\quad \times G(S(t)f_1(\tau_1)S(t)f_2(\tau_2))S(t)f_3(\tau_3)S(t)f_4(\tau_4) \, d\sigma d\tau_1 d\tau_2 d\tau_3 d\tau_4. \end{aligned}$$

Hence, the quadrilinear estimate (2.2), applied pointwise in $\tau_1, \tau_2, \tau_3, \tau_4$ and after having chosen $\chi \in \mathcal{C}_0^\infty(\mathbb{R})$ such that $\chi = 1$ on $(0, 1)$, gives

$$\begin{aligned} \left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 \, d\sigma dt \right| &\leq C_\varepsilon m(N_1, N_2, N_3, N_4)^\varepsilon \prod_{i=1}^4 \int_{\mathbb{R}} \|\hat{v}_j\|_{L^2(\mathbb{S}^2)}(\tau) \, d\tau \\ &\leq C_\varepsilon m(N_1, N_2, N_3, N_4)^\varepsilon \prod_{i=1}^4 \|\langle \tau \rangle^b \hat{v}_j\|_{L^2(\mathbb{R} \times \mathbb{S}^2)} \\ &= C_\varepsilon m(N_1, N_2, N_3, N_4)^\varepsilon \prod_{i=1}^4 \|u_j\|_{X^{0,b}}, \end{aligned}$$

where the Cauchy-Schwarz inequality could be applied since $b > 1/2$.

To treat the general case, it suffices now to introduce a function $\psi \in \mathcal{C}_0^\infty(\mathbb{R})$ supported in $(0, 1)$ such that $\sum_{n \in \mathbb{Z}} \psi(t - \frac{n}{2}) = 1$, to decompose $u_j = \sum_{n \in \mathbb{Z}} u_{j,n}$, with $u_{j,n} = \psi(t - \frac{n}{2})u_j(t)$, $j = 1, 2, 3, 4$, and to apply the result of the first step to the functions $u_{j,n}$. The conclusion follows from the almost orthogonality property satisfied by the Bourgain spaces

$$\sum_{n \in \mathbb{Z}} \|u_{j,n}\|_{X^{0,b}}^2 \leq C \|u_j\|_{X^{0,b}}^2.$$

□

Step 3. Proof of Lemma 2.4 by dyadic decomposition.

We have now the tools to prove Lemma 2.4. We shall only prove (2.5), the proof of (2.6) being similar. By duality, we have to show that, for all $u_4 \in X^{-s,b'}$,

$$\left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_1 u_2) u_3 u_4 d\sigma dt \right| \leq C \left(\prod_{j=1}^3 \|u_j\|_{X^{s,b}} \right) \|u_4\|_{X^{-s,b'}}. \quad (2.13)$$

Let us perform a dyadic decomposition of the u_j 's, setting $u_{j,N} = \mathbb{1}_{\sqrt{1-\Delta_\sigma} \in [N, 2N]}(u_j)$. We have

$$\|u_j\|_{X^{s,b}}^2 \simeq \sum_N N^{2s} \|u_{j,N}\|_{X^{0,b}}^2 \simeq \sum_N \|u_{j,N}\|_{X^{s,b}}^2 \quad (2.14)$$

where N are dyadic integers. Proving (2.13) is equivalent to estimating the sum

$$\sum_{N_1, N_2, N_3, N_4} J(N_1, N_2, N_3, N_4),$$

where

$$J(N_1, N_2, N_3, N_4) = \int_{\mathbb{R}} \int_{\mathbb{S}^2} G(u_{1,N_1} u_{2,N_2}) u_{3,N_3} u_{4,N_4} d\sigma dt.$$

We now remark that in the above sum, all the terms with $N_4 > 4 \max(N_1, N_2, N_3)$ are zero. To prove this point, we first claim that the operator G is a function of Δ_σ , more precisely, that we have for all f

$$G(f) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \frac{1}{|x-y|} f(y) d\sigma(y) = (1 - 4\Delta_\sigma)^{-1/2}(f). \quad (2.15)$$

The proof of the claim is done below. Moreover, from spectral localization u_{j,N_j} is the sum of spherical harmonics of degrees between $\sqrt{N_j}/2$ and $\sqrt{2N_j}$. One can deduce from properties of products of spherical harmonics that the product $G(u_{1,N_1} u_{2,N_2}) u_{3,N_3}$ is a sum of spherical harmonics of degrees less than $\sqrt{2 \max(N_1, N_2, N_3)}$, so orthogonal to u_4 if $N_4 > 4 \max(N_1, N_2, N_3)$.

Hence, by symmetry, it suffices to consider the terms with $N_1 \leq N_2 \leq N_3$ and $N_4 \leq 4N_3$. Pick s' such that $0 < s' < s$. By Lemma 2.6, there exists $0 < b' < 1/2$ such that

$$\begin{aligned} \sum_{N_1, N_2, N_3, N_4} J(N_1, N_2, N_3, N_4) &\leq C \sum_{N_1, N_2, N_3} \sum_{N_4 \leq 4N_3} N_1^{s'} N_2^{s'} \prod_{i=1}^4 \|u_{j,N_j}\|_{X^{0,b'}}, \\ &\leq C \left(\prod_{j=1,2} \sum_{N_j} N_j^{s'-s} \|u_{j,N_j}\|_{X^{s,b'}} \right) \sum_{N_3} \sum_{N_4 \leq 4N_3} \left(\frac{N_3}{N_4} \right)^{-s} \|u_{3,N_3}\|_{X^{s,b'}} \|u_{4,N_4}\|_{X^{-s,b'}} \end{aligned}$$

where we used (2.14). The series in N_1 and N_2 converges easily. Denoting $N_4 = 2^n$ and $N_3/N_4 = 2^m$, the series in N_3, N_4 can be bounded by a simple Cauchy-Schwarz argument. Indeed, this series reads

$$\begin{aligned} &\sum_{m \geq -2} 2^{-ms} \sum_{n \geq \max(0, -m)} \|u_{3,2^{m+n}}\|_{X^{s,b'}} \|u_{4,2^n}\|_{X^{-s,b'}} \\ &\leq \left(\sum_{m \geq -2} 2^{-ms} \right) \left(\sum_n \|u_{3,2^n}\|_{X^{s,b'}}^2 \right)^{1/2} \left(\sum_n \|u_{4,2^n}\|_{X^{-s,b'}}^2 \right)^{1/2} \\ &\leq C \|u_3\|_{X^{s,b'}} \|u_4\|_{X^{-s,b'}}. \end{aligned}$$

The proof of Lemma 2.4 is complete, so Proposition 2.2 is proved.

Proof of the claim (2.15). We use a very old argument that can be found, e.g., in Poincaré's works [25]. Let us compute explicitly the action of G on a spherical harmonic Y_ℓ^m . We define the following function on \mathbb{R}^3 :

$$u(r, \sigma) = r^\ell Y_\ell^m(\sigma) \text{ for } r \leq 1, \quad u(r, \sigma) = r^{-\ell-1} Y_\ell^m(\sigma) \text{ for } r \geq 1.$$

It is readily seen that u is harmonic on $\mathbb{R}^3 \setminus \mathbb{S}^2$, so by computing explicitly the jump of $\partial_r u$ on \mathbb{S}^2 we obtain

$$-\Delta u = (2\ell + 1)Y_\ell^m d\sigma$$

in the distribution sense, where $d\sigma$ is the surface measure on \mathbb{S}^2 . Since u is decreasing at the infinity, this means that

$$u(r, \sigma) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \frac{1}{|r\sigma - \sigma'|} (2\ell + 1)Y_\ell^m d\sigma',$$

thus

$$u(1, \sigma) = (2\ell + 1)G(Y_\ell^m).$$

Denoting $\lambda = \ell(\ell + 1)$, we have $2\ell + 1 = \sqrt{1 + 4\lambda}$, so we have

$$G(Y_\ell^m) = \frac{1}{\sqrt{1 + 4\lambda}} Y_\ell^m$$

and (2.15) is proved. \square

2.2. Quadrilinear estimate. In this section, we prove the quadrilinear estimate (2.2) which is, thanks to Proposition 2.2, the core of the proof of local existence result in Theorem 1.1. The main result of this section is the following proposition.

Proposition 2.7. *There exists a constant $C > 0$ such that for any $f_1, f_2, f_3, f_4 \in L^2(\mathbb{S}^2)$ satisfying*

$$\mathbb{1}_{\sqrt{1-\Delta_\sigma} \in [N_j, 2N_j]}(f_j) = f_j, \quad j = 1, 2, 3, 4, \quad (2.16)$$

and for all $\varepsilon \in]0, \frac{1}{2}[$ one has the following quadrilinear estimate

$$\sup_{\tau \in \mathbb{R}} \left| \int_{\mathbb{R}} \int_{\mathbb{S}^2} \chi(t) e^{it\tau} G(u_1 u_2) u_3 u_4 d\sigma dt \right| \leq C_\varepsilon (m(N_1, N_2, N_3, N_4))^\varepsilon \prod_{i=1}^4 \|f_j\|_{L^2(\mathbb{S}^2)} \quad (2.17)$$

where $u_j(t) = S(t)f_j$ for $j = 1, 2, 3, 4$, $S(t) = e^{it\Delta_\sigma}$ denoting the free evolution, and $\chi \in \mathcal{C}_0^\infty(\mathbb{R})$ is arbitrary.

Proof. Let us decompose the f_j 's on spherical harmonics:

$$f_j = \sum_{n_j=N_j/2}^{2N_j} H_{n_j}^j \quad \text{with} \quad \Delta_\sigma H_{n_j}^j + n_j(n_j + 1)H_{n_j}^j = 0,$$

so that

$$u_j(t) = \sum_{n_j=N_j/2}^{2N_j} e^{-itn_j(n_j+1)} H_{n_j}^j.$$

The quantity to estimate can be written as

$$\int_{\mathbb{R}} \int_{\mathbb{S}^2} \chi(t) e^{it\tau} G(u_1 u_2) u_3 u_4 d\sigma dt = \sum_{n_1, n_2, n_3, n_4} \omega_{n_1, n_2, n_3, n_4}(\tau) I(n_1, n_2, n_3, n_4)$$

where

$$\omega_{n_1, n_2, n_3, n_4}(\tau) = \hat{\chi} \left(\tau - \sum_{j=1}^4 \varepsilon_j n_j (n_j + 1) \right) \quad \text{with } \varepsilon_j = (-1)^{j+1},$$

$\hat{\chi}$ being the Fourier transform of χ , and

$$I(n_1, n_2, n_3, n_4) = \int_{\mathbb{S}^2} G(H_{n_1}^1 H_{n_2}^2) H_{n_3}^3 H_{n_4}^4 d\sigma.$$

By symmetry and self-adjointness of G , assume for instance that $N_3 = \min(N_1, N_2, N_3, N_4)$. Let us estimate $I(n_1, n_2, n_3, n_4)$ by adapting the proof of multilinear estimates in [10, 11, 12, 20]. One can decompose $H_{n_j}^j$ using a microlocal partition of the unity with semiclassical cut-off of the form $\chi_j(x, h_j D)$ (with small support):

$$H_{n_j}^j = \varphi_j + \sum_k \chi_j^k(x, h_j D) H_{n_j}^j,$$

where φ_j is regular (for all s , $\|D^s \varphi_j\|_{L^2} \leq C_s \|H_{n_j}^j\|_{L^2}$) and where the sum is finite. Thus, we only have to estimate some terms of the form

$$\tilde{I} = \int_{\mathbb{S}^2} G(V_1 V_2) V_3 V_4 d\sigma,$$

where V_j is of the form $V_j = \chi_j(x, h_j D) H_{n_j}^j$. The key point, from Burq, Gérard, Tzvetkov [10, 11, 12] is that, using semiclassical Strichartz estimates in dimension one, for each choice of $(\chi_j)_{j=1,2,3,4}$ one can find a local system of linear coordinates (x, y) such that, for all $p, q \in [2, \infty]$ satisfying $\frac{2}{p} + \frac{1}{q} = \frac{1}{2}$, we have

$$\|V_j\|_{L_x^p L_y^q} \leq C N_j^{1/p} \|H_{n_j}^j\|_{L^2(\mathbb{S}^2)}, \quad \text{for } j = 1, 2, 3, 4. \quad (2.18)$$

The proof of this result uses the fact that the function $H_{n_j}^j$ satisfies

$$h_j^2 \Delta_\sigma H_{n_j}^j + H_{n_j}^j = 0, \quad \text{with } h_j^2 = \frac{1}{n_j(n_j + 1)} \sim N_j^2,$$

which can be locally seen as an evolution equation where the well-chosen variable x plays the role of a time.

In local coordinates we have

$$|\tilde{I}| \leq C \|G(V_1 V_2)\|_{L_x^\infty L_y^s} \|V_3\|_{L_x^{1/\varepsilon} L_y^r} \|V_4\|_{L_x^2 L_y^2}$$

with $s = \frac{1}{2\varepsilon}$ and $\frac{1}{r} = \frac{1}{2} - 2\varepsilon$. Since

$$|G(V_1 V_2)| \leq C \int \frac{1}{|x - x'| + |y - y'|} |V_1 V_2|(x', y') dx' dy',$$

a Hardy-Littlewood Sobolev inequality gives

$$\begin{aligned} \|G(V_1 V_2)\|_{L_x^\infty L_y^s} &\leq C \sup_x \int \frac{1}{|x - x'|^{1-2\varepsilon}} \|V_1 V_2(x', \cdot)\|_{L^1} dx' \\ &\leq C_\varepsilon \|V_1\|_{L_x^\infty L_y^2} \|V_2\|_{L_x^\infty L_y^2}, \end{aligned}$$

where we used the fact that all the functions are compactly supported. Then we apply (2.18) with the pairs $(p, q) = (\infty, 2)$ and $(\frac{1}{\varepsilon}, r)$:

$$\begin{aligned} |\tilde{I}| &\leq C_\varepsilon \|V_1\|_{L_x^\infty L_y^2} \|V_2\|_{L_x^\infty L_y^2} \|V_3\|_{L_x^{1/\varepsilon} L_y^r} \|V_4\|_{L_x^2 L_y^2} \\ &\leq C_\varepsilon (N_3)^\varepsilon \prod_{j=1}^4 \|H_{n_j}^j\|_{L^2}. \end{aligned}$$

To conclude, it remains to estimate uniformly with respect to τ the quantity

$$Q(\tau) = \sum_{n_1, n_2, n_3, n_4} |\omega_{n_1, n_2, n_3, n_4}(\tau)| \prod_{j=1}^4 \|H_{n_j}^j\|_{L^2}$$

where, in the sum, n_1, n_2, n_3, n_4 satisfy

$$\forall j \in \{1, 2, 3, 4\} \quad \frac{N_j}{2} \leq n_j \leq 2N_j. \quad (2.19)$$

This quantity has already been estimated in [20], but we reproduce again this proof here for the sake of completeness.

Denote by $\Lambda(k)$ the set of (n_1, n_2, n_3, n_4) satisfying (2.19) and

$$\sum_{j=1}^4 \varepsilon_j n_j (n_j + 1) = k.$$

We deduce from the decay of $\hat{\chi}$ at the infinity that

$$\begin{aligned} Q(\tau) &\leq C \sum_{\ell \in \mathbb{Z}} \frac{1}{1 + \ell^2} \sum_{\Lambda([\tau] + \ell)} \prod_{j=1}^4 \|H_{n_j}^j\|_{L^2} \\ &\leq C \sup_{k \in \mathbb{Z}} \sum_{\Lambda(k)} \prod_{j=1}^4 \|H_{n_j}^j\|_{L^2}. \end{aligned}$$

Let us denote by α, β the two indices such that $m(N_1, N_2, N_3, N_4) = (N_\alpha, N_\beta)$ and by γ, δ the two other indices: $\{\gamma, \delta\} = \{1, 2, 3, 4\} \setminus \{\alpha, \beta\}$. Then we introduce the set

$$\Gamma(k, i, j) = \left\{ (n_i, n_j) : \frac{N_i}{2} \leq n_i \leq 2N_i, \frac{N_j}{2} \leq n_j \leq 2N_j \right. \\ \left. \text{and } \varepsilon_i n_i (n_i + 1) + \varepsilon_j n_j (n_j + 1) = k \right\}.$$

Now, denoting

$$S(k, i, j) = \sum_{\Gamma(k, i, j)} \|H_{n_i}^i\|_{L^2} \|H_{n_j}^j\|_{L^2},$$

we split the sum as follows:

$$Q(\tau) \leq C \sup_{k \in \mathbb{Z}} \sum_{k' \in \mathbb{Z}} S(k, \alpha, \gamma) S(k - k', \beta, \delta).$$

Then we apply the following elementary result of number theory (see e.g. [10]).

Lemma 2.8. *Let $\sigma = \pm 1$. For all $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that, given $M \in \mathbb{Z}$ and $N \in \mathbb{N}^*$,*

$$\#\{(k_1, k_2) \in \mathbb{N}^2 : N \leq k_1 \leq 2N, k_1^2 + \sigma k_2^2 = M\} \leq C_\varepsilon N^\varepsilon.$$

Hence we get

$$\sup_{k'} \#\Gamma(k', \alpha, \gamma) \leq C_\varepsilon N_\alpha^\varepsilon, \quad \sup_{k, k'} \#\Gamma(k - k', \beta, \delta) \leq C_\varepsilon N_\beta^\varepsilon,$$

and by Cauchy-Schwarz

$$\begin{aligned} \sum_{k' \in \mathbb{Z}} S(k, \alpha, \gamma) S(k - k', \beta, \delta) &\leq C_\varepsilon (N_\alpha N_\beta)^\varepsilon \times \\ &\times \left(\sum_{k'} \sum_{\Gamma(k', \alpha, \gamma)} \|H_{n_\alpha}^\alpha\|_{L^2}^2 \|H_{n_\gamma}^\gamma\|_{L^2}^2 \right)^{1/2} \left(\sum_{k'} \sum_{\Gamma(k', \beta, \delta)} \|H_{n_\beta}^\beta\|_{L^2}^2 \|H_{n_\delta}^\delta\|_{L^2}^2 \right)^{1/2} \\ &\leq C_\varepsilon (N_\alpha N_\beta)^\varepsilon \prod_{j=1}^4 \|f_j\|_{L^2}, \end{aligned}$$

where we used the orthogonality of the spherical harmonics. The proof is complete. \square

2.3. An instability result in L^2 . In this section, we prove Theorem 1.2 and exhibit a high frequency instability for the Cauchy problem in $L^2(\mathbb{S}^2)$ for the Schrödinger-Poisson system (1.2). Consider the following spherical harmonic on \mathbb{S}^2

$$\psi_n(x) = (x_1 + ix_2)^n \quad (2.20)$$

where (x_1, x_2, x_3) are cartesian coordinates on \mathbb{R}^3 , $\mathbb{S}^2 = \{x_1^2 + x_2^2 + x_3^2 = 1\}$. Remark that this function concentrates on the equator $\{x_3 = 0\}$ as $n \rightarrow \infty$.

In the case of the cubic nonlinear Schrödinger equation on \mathbb{S}^d , an instability in some H^s space has been shown in [13] (see also [2] for a more precise result) by finding an ansatz for the solution of the equation with an initial data proportional to ψ_n and of order 1 in H^s . A different approach has been presented in [19] in order to exhibit the same instability. It consists in constructing a stationary solution for the nonlinear Schrödinger equation, rapidly oscillating in time, by minimizing the nonlinear energy of the problem. The point is to estimate precisely the corresponding nonlinear eigenvalue as n goes to infinity. Let us adapt this argument to our case.

Lemma 2.9. *The function ψ_n defined by (2.20) satisfies*

$$\|\psi_n\|_{L^2}^2 = \frac{C_1}{\sqrt{n}} + \mathcal{O}(n^{-3/2}), \quad \|\nabla_\sigma \psi_n\|_{L^2}^2 = n(n+1)\|\psi_n\|_{L^2}^2, \quad (2.21)$$

$$\int_{\mathbb{S}^2} G(|\psi_n|^2) |\psi_n|^2 d\sigma = C_2 \frac{\log n}{n} + \frac{C_3}{n} + \mathcal{O}\left(\frac{1}{n^2}\right), \quad (2.22)$$

where C_1, C_2, C_3 are constant real numbers.

Proof. The first estimates (2.21) are immediate. Let us prove (2.22), writing

$$\begin{aligned} \int_{\mathbb{S}^2} G(|\psi_n|^2) |\psi_n|^2 d\sigma &= \int \frac{(\cos \theta)^{2n+1} (\cos \theta')^{2n+1}}{(|\cos \theta e^{i\varphi} - \cos \theta' e^{i\varphi'}|^2 + (\sin \theta - \sin \theta')^2)^{1/2}} d\theta d\theta' d\varphi d\varphi' \\ &= \int (\cos \theta)^{2n+1/2} (\cos \theta')^{2n+1/2} \left(\frac{1 - \cos(\theta - \theta')}{\cos \theta \cos \theta'} + 1 - \cos(\varphi - \varphi') \right)^{-1/2} d\theta d\theta' d\varphi d\varphi', \end{aligned}$$

where the integration domain is $(\theta, \theta', \varphi, \varphi') \in [-\pi/2, \pi/2] \times [-\pi/2, \pi/2] \times [0, 2\pi] \times [0, 2\pi]$. As $n \rightarrow +\infty$, the main contribution in this integral is near the equator $\theta = 0, \theta' = 0$. Hence some elementary analysis using

$$\int_0^{2\pi} \int_0^{2\pi} (t + 1 - \cos(\varphi - \varphi'))^{-1/2} d\varphi d\varphi' = a \log t + b + \mathcal{O}(t) \quad \text{as } t \rightarrow 0+,$$

yields (2.22). \square

Now, for all $0 < \delta \leq 1$ we introduce the function $\phi_n = c_n \psi_n$ such that $\|\phi_n\|_{L^2} = \delta$. We deduce from Lemma 2.9 that

$$\|\nabla_\sigma \phi_n\|_{L^2}^2 = n(n+1)\delta^2, \quad (2.23)$$

$$\int_{\mathbb{S}^2} G(|\phi_n|^2)|\phi_n|^2 d\sigma = C_4 \delta^4 \log n + C_5 \delta^4 + \mathcal{O}\left(\frac{\delta^4}{n}\right), \quad (2.24)$$

for some constants C_4 and C_5 independent of n and δ . We will see that this factor $\log n$ is at the origin of the high frequency instability.

Let us define as in [19] the space

$$L_n^2 = \{f \in L^2(\mathbb{S}^2) : \forall \alpha \in \mathbb{R} \quad f \circ R_\alpha = e^{in\alpha} f\},$$

where R_α denotes the rotation of angle α around the x_3 axis. Since the spherical harmonics take the form

$$Y_\ell^m = c_{m,\ell} P_\ell^m(\sin \theta) e^{im\varphi},$$

in spherical coordinates (θ is the angle between the vector x and the equatorial plane $\{x_3 = 0\}$), where P_ℓ^m is a Legendre function and $c_{m,\ell}$ is a normalization factor, it is readily seen that the space L_n^2 is characterized by

$$L_n^2 = \text{span} \{Y_{n+k}^n, k \in \mathbb{N}\}, \quad (2.25)$$

where n is fixed. Therefore, we have a characterization of ϕ_n (up to a phase factor) as the minimizer of the Dirichlet energy on L_n^2 :

$$\phi_n = \text{argmin} \{ \|\nabla_\sigma u\|_{L^2}^2 \text{ for } u \in L_n^2 \text{ such that } \|u\|_{L^2} = \delta \}.$$

Let us now minimize the energy associated to (1.2):

$$\mathcal{E}(u) = \|\nabla_\sigma u\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{S}^2} G(|u|^2)|u|^2 d\sigma$$

on the sphere of radius δ of L_n^2 . It is easy to prove the compactness of minimizing sequences and to obtain the existence of a minimizer f_n to this problem. The Euler equation reads

$$-\Delta_\sigma f_n + G(|f_n|^2)f_n = \omega_n f_n,$$

with

$$\omega_n = \frac{1}{\delta^2} \left(\|\nabla_\sigma f_n\|_{L^2}^2 + \int_{\mathbb{S}^2} G(|f_n|^2)|f_n|^2 d\sigma \right) > 0, \quad (2.26)$$

and the function

$$u_n(t, x) = e^{-it\omega_n} f_n(x)$$

is a solution of (1.2). The key of the method is now to show that f_n is close to ϕ_n , up to a phase factor, and to give a precise estimate for ω_n .

Lemma 2.10. *There exists a constant C independent of $n \in \mathbb{N}^*$ and $\delta \in (0, 1]$ such that, for some $\alpha_n \in \mathbb{R}$,*

$$\|f_n - e^{i\alpha_n} \phi_n\|_{L^2} \leq C\delta^2 \left(\frac{\log n}{n}\right)^{1/2}, \quad (2.27)$$

$$\omega_n = n(n+1) + C_4\delta^2 \log n + C_5\delta^2 + \mathcal{O}\left(\delta^3 \frac{\log n}{n^{1/4}}\right). \quad (2.28)$$

Proof. Since f_n belongs to L_n^2 , let us decompose this function on the spherical harmonics, according to (2.25):

$$f_n = a_0\phi_n + \sum_{k=1}^{\infty} a_k Y_{n+k}^n, \quad (2.29)$$

the Y_ℓ^n being chosen normalized in L^2 . One can deduce from the normalization conditions that

$$|a_0|^2\delta^2 + \sum_{k=1}^{\infty} |a_k|^2 = \delta^2 \quad (2.30)$$

and the property $\mathcal{E}(f_n) \leq \mathcal{E}(\phi_n)$ reads

$$\begin{aligned} |a_0|^2\delta^2 n(n+1) + \sum_{k=1}^{\infty} |a_k|^2(n+k)(n+k+1) + \frac{1}{2} \int_{\mathbb{S}^2} G(|f_n|^2)|f_n|^2 d\sigma \\ \leq \delta^2 n(n+1) + \frac{1}{2} \int_{\mathbb{S}^2} G(|\phi_n|^2)|\phi_n|^2 d\sigma. \end{aligned} \quad (2.31)$$

By subtracting $n(n+1) \times (2.30)$ to (2.31), we obtain

$$\sum_{k=1}^{\infty} k(2n+k+1)|a_k|^2 \leq \frac{1}{2} \int_{\mathbb{S}^2} G(|\phi_n|^2)|\phi_n|^2 d\sigma \leq C\delta^4 \log n, \quad (2.32)$$

where we used (2.24). Hence we have

$$\|f_n - a_0\phi_n\|_{L^2}^2 = \sum_{k=1}^{\infty} |a_k|^2 \leq C\delta^4 \frac{\log n}{n}. \quad (2.33)$$

Inserting (2.33) in (2.30) leads to

$$0 \leq 1 - |a_0|^2 \leq C\delta^2 \frac{\log n}{n} \quad (2.34)$$

and (2.27) follows, setting $e^{i\alpha_n} = \frac{a_0}{|a_0|}$.

Let us now prove (2.28). By combining the two inequalities

$$\|\nabla_\sigma f_n\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{S}^2} G(|f_n|^2)|f_n|^2 d\sigma \leq \|\nabla_\sigma \phi_n\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{S}^2} G(|\phi_n|^2)|\phi_n|^2 d\sigma$$

and

$$\|\nabla_\sigma \phi_n\|_{L^2}^2 \leq \|\nabla_\sigma f_n\|_{L^2}^2,$$

and by using (2.26) it comes

$$\begin{aligned} 0 \leq \|\nabla_\sigma \phi_n\|_{L^2}^2 + \int_{\mathbb{S}^2} G(|\phi_n|^2)|\phi_n|^2 d\sigma - \delta^2 \omega_n \\ \leq \int_{\mathbb{S}^2} (G(|\phi_n|^2)|\phi_n|^2 d\sigma - G(|f_n|^2)|f_n|^2 d\sigma). \end{aligned} \quad (2.35)$$

Moreover, for all $M > 0$ and for all u, v in the centered ball of $L^{8/3}(\mathbb{S}^2)$ of radius M , we deduce from the estimate (2.7) on the operator G that

$$\left| \int_{\mathbb{S}^2} (G(|u|^2)|u|^2 - G(|v|^2)|v|^2) d\sigma \right| \leq CM^3 \|u - v\|_{L^{8/3}}. \quad (2.36)$$

Let us now estimate f_n, ϕ_n and the difference between these two functions in $L^{8/3}$. The key point will be the following estimate on spherical harmonics due to Sogge [27]:

$$\|Y_\ell^n\|_{L^{8/3}} \leq C\ell^{1/16} \|Y_\ell^n\|_{L^2}. \quad (2.37)$$

Therefore we have

$$\|\phi_n\|_{L^{8/3}} \leq C\delta n^{1/16}. \quad (2.38)$$

Moreover, from the decomposition (2.29) of f_n and by the Minkowski inequality,

$$\|f_n - e^{i\alpha_n} \phi_n\|_{L^{8/3}} \leq \left\| \left(a_0 - \frac{a_0}{|a_0|} \right) \phi_n \right\|_{L^{8/3}} + \sum_{k=1}^{\infty} |a_k| \|Y_{n+k}^n\|_{L^{8/3}}.$$

By (2.37), (2.32) and Cauchy-Schwarz, we have

$$\begin{aligned} \sum_{k=1}^{\infty} |a_k| \|Y_{n+k}^n\|_{L^{8/3}} &\leq C \left(\sum_{k=1}^{\infty} \frac{(n+k)^{1/8}}{k(2n+k+1)} \right)^{1/2} \left(\sum_{k=1}^{\infty} k(2n+k+1) |a_k|^2 \right)^{1/2} \\ &\leq C\delta^2 \frac{\log n}{n^{7/16}}. \end{aligned}$$

Hence, using also (2.34) and (2.38), we obtain

$$\|f_n - e^{i\alpha_n} \phi_n\|_{L^{8/3}} \leq C\delta^2 \frac{\log n}{n^{7/16}} \quad \text{and} \quad \|f_n\|_{L^{8/3}} \leq C\delta n^{1/16}.$$

Finally, (2.36) leads to

$$\left| \int_{\mathbb{S}^2} (G(|\phi_n|^2)|\phi_n|^2 - G(|f_n|^2)|f_n|^2) d\sigma \right| \leq C\delta^5 \frac{\log n}{n^{1/4}}.$$

By inserting this estimate in (2.35), then by using (2.23), (2.24), we obtain (2.28). \square

Proof of Theorem 1.2.

We now have the tools to prove the high frequency instability and Theorem 1.2. Remark first that, replacing f_n by $f_n e^{-i\alpha_n}$, one may assume $\alpha_n = 0$ in (2.27). As in [19], we consider two values of δ :

$$\delta_n = \delta_0, \quad \delta'_n = \kappa_n \delta_0,$$

where $0 < \delta_0 \leq 1$ and $\kappa_n \rightarrow 1$ as $n \rightarrow +\infty$ in a way that is defined below (see (2.39)), and we denote by f_n, ϕ_n and f'_n, ϕ'_n the corresponding functions.

By (2.27), we have

$$\|f_n - f'_n\|_{L^2} \leq \|f_n - \phi_n\|_{L^2} + \|f'_n - \phi'_n\|_{L^2} + |\delta_n - \delta'_n| \leq C \left(\frac{\log n}{n} \right)^{1/2} \delta_0 + |1 - \kappa_n| \delta_0,$$

whereas the corresponding solutions of (1.2)

$$u_n = e^{-it\omega_n} f_n, \quad u'_n = e^{-it\omega'_n} f'_n$$

satisfy

$$\begin{aligned} \|u_n - u'_n\|_{L^2} &\geq \left| e^{-it\omega_n} - e^{-it\omega'_n} \right| - \|f_n - f'_n\|_{L^2} \\ &\geq \left| \sin\left(\frac{t}{2}(\omega_n - \omega'_n)\right) \right| \delta_0 - C \left(\frac{\log n}{n}\right)^{1/2} \delta_0 - |1 - \kappa_n| \delta_0. \end{aligned}$$

By (2.28), we have

$$\omega_n - \omega'_n = C_4(1 - \kappa_n^2)\delta_0^2 \log n + C_5(1 - \kappa_n^2)\delta_0^2 + \mathcal{O}\left(\frac{\log n}{n^{1/4}}\right).$$

If now we define κ_n by

$$\kappa_n = \left(1 - (\log n)^{-1/2}\right)^{1/2} \quad (2.39)$$

and an observation time t_n by

$$t_n = \frac{\pi}{\omega_n - \omega'_n} = \frac{\pi}{C_4\delta_0^2(\log n)^{1/2}}(1 + o(1)),$$

we have clearly a sequence $t_n \rightarrow 0+$ such that

$$\|(u_n - u'_n)(t_n, \cdot)\|_{L^2} \geq \delta_0 - \varepsilon_n, \quad \|(u_n - u'_n)(0, \cdot)\|_{L^2} \leq \varepsilon_n,$$

where $\varepsilon_n \rightarrow 0$ as $n \rightarrow +\infty$. Since u_n and u'_n are in the ball of radius δ_0 of $L^2(\mathbb{S}^2)$, this contradicts Item (iii) of Definition 2.1, which means that (1.2) is not uniformly well-posed on $L^2(\mathbb{S}^2)$. \square

3. Asymptotic analysis

The aim of this second part of the paper is to prove Theorem 1.5, which justifies the use of (1.2) (in fact, of the mixed-state version (1.14) of this equation) as a model for quantum transport on a surface by means of asymptotic analysis, deriving this system from a well-established model, the three-dimensional Schrödinger-Poisson system (1.4). The Cauchy problem for this system without the confinement potential was studied in [8, 24] in the energy space, and in [14] in L^2 .

3.1. Estimating the Poisson nonlinearity. In this subsection, we obtain some estimates on the Poisson nonlinearity for functions confined near the sphere by the confinement operator V_c^ε . We will see that the following family of norms is well-adapted for the study of the singularly perturbed nonlinear problem (1.4):

$$\|\langle \nabla_\sigma \rangle^s u\|_{L^2} + \varepsilon \|u\|_{\mathcal{B}^1},$$

where s is a positive integer, whereas the following family of norms is well-adapted to the limit problem (1.14):

$$\|\langle \nabla_\sigma \rangle^{s_1} u\|_{L^2} + \|(1 + \varepsilon^2 H_r)^{s_2/2} u\|_{L^2},$$

where s_1 and s_2 are nonnegative integers and where we recall the definition (1.11) of the Hamiltonian H_r , which is nonnegative thanks to Assumption 1.3. The nonlinear analysis of (1.4) will be based on Lemmas 3.1 and 3.2.

Lemma 3.1. *Let $u \in \mathcal{B}^1$, such that $\langle \nabla_\sigma \rangle^s u \in L^2$ with $s \geq 1$. Then there exists $\gamma > 0$ and a constant $C > 0$ such that, for all $\varepsilon \in (0, 1]$, the nonlinearities $V(|u|^2)u$ and $\mathbb{G}(|u|^2)u$, respectively defined by (1.6) and (1.15), satisfy the tame estimate*

$$\|\langle \nabla_\sigma \rangle^s (\mathbb{G}(|u|^2)u)\|_{L^2} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 \|\langle \nabla_\sigma \rangle^s u\|_{L^2} \quad (3.1)$$

$$\|\langle \nabla_\sigma \rangle^s (V(|u|^2)u)\|_{L^2} \leq C \left(\|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + \varepsilon^{2(1+\gamma)} \|u\|_{\mathcal{B}^1}^2 \right) \|\langle \nabla_\sigma \rangle^s u\|_{L^2} \quad (3.2)$$

where C is independent of ε . Moreover, if $u, v \in \mathcal{B}^1$ are such that $\langle \nabla_\sigma \rangle u, \langle \nabla_\sigma \rangle v \in L^2$, then

$$\begin{aligned} & \|\langle \nabla_\sigma \rangle (\mathbb{G}(|u|^2)u - \mathbb{G}(|v|^2)v)\|_{L^2} \\ & \leq C \left(\|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + \|\langle \nabla_\sigma \rangle v\|_{L^2}^2 \right) \|\langle \nabla_\sigma \rangle (u - v)\|_{L^2}, \end{aligned} \quad (3.3)$$

$$\begin{aligned} & \|\langle \nabla_\sigma \rangle (V(|u|^2)u - V(|v|^2)v)\|_{L^2} \\ & \leq C \left(\|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + \varepsilon^{2(1+\gamma)} \|u\|_{\mathcal{B}^1}^2 + \|\langle \nabla_\sigma \rangle v\|_{L^2}^2 + \varepsilon^{2(1+\gamma)} \|v\|_{\mathcal{B}^1}^2 \right) \times \\ & \quad \times \left(\|\langle \nabla_\sigma \rangle (u - v)\|_{L^2} + \varepsilon^{2(1+\gamma)} \|u - v\|_{\mathcal{B}^1} \right). \end{aligned} \quad (3.4)$$

Finally, if $\langle \nabla_\sigma \rangle^2 u \in L^2$, then we have

$$\|\langle \nabla_\sigma \rangle (V(|u|^2)u - \mathbb{G}(|u|^2)u)\|_{L^2} \leq C\varepsilon^\gamma \left(\|\langle \nabla_\sigma \rangle^2 u\|_{L^2}^3 + \varepsilon^3 \|u\|_{\mathcal{B}^1}^3 \right). \quad (3.5)$$

Proof. Step 1. Sobolev embeddings. Let us first write some anisotropic Sobolev embeddings that will be useful several times in the proof. In spherical coordinates (recall that $x = r\sigma$), we shall denote

$$\begin{aligned} \forall p, q \in [1, +\infty) \quad & \|u\|_{L_r^p L_\sigma^q} = \left(\int_0^\infty \left(\int_{\mathbb{S}^2} |u|^q d\sigma \right)^{q/p} r^2 dr \right)^{1/q}, \\ \forall q \in [1, +\infty) \quad & \|u\|_{L_r^\infty L_\sigma^q} = \text{Sup}_{r>0} \text{ess} \|u(r, \cdot)\|_{L_\sigma^q}. \end{aligned}$$

By Sobolev embeddings in dimension 2, we have

$$\|u\|_{L_r^2 L_\sigma^p} \leq C_p \|\langle \nabla_\sigma \rangle u\|_{L^2} \quad (3.6)$$

for all $p \in [2, \infty)$. Moreover, since the H^1 norm reads in spherical coordinates

$$\|u\|_{H^1}^2 = \|u\|_{L^2}^2 + \|\partial_r u\|_{L^2}^2 + \left\| \frac{1}{r} \nabla_\sigma u \right\|_{L^2}^2,$$

we deduce from the Hardy inequality

$$\left\| \frac{u}{r} \right\|_{L^2} \leq C \|\nabla u\|_{L^2}$$

that

$$\left\| \langle \nabla_\sigma \rangle \left(\frac{u}{r} \right) \right\|_{L^2} \leq C \|u\|_{H^1}. \quad (3.7)$$

Hence, by Sobolev embeddings, we get

$$\left\| \frac{u}{r} \right\|_{L_r^2 L_\sigma^p} \leq C_p \|u\|_{H^1} \leq C_p \|u\|_{\mathcal{B}^1} \quad (3.8)$$

for all $p \in [2, \infty)$. Finally, we claim that

$$\left\| \frac{u}{r^{1/2}} \right\|_{L_r^2 L_\sigma^4} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2} + C\varepsilon^{1+\gamma} \|u\|_{\mathcal{B}^1}, \quad (3.9)$$

for some $\gamma > 0$ independent of ε . In order to prove (3.9), let us split the integral on $\{r > 1/2\} \cup \{r < 1/2\}$. We get

$$\begin{aligned} \left\| \frac{u}{r^{1/2}} \right\|_{L_r^2 L_\sigma^4} &= \left\| \frac{u}{r^{1/2}} \mathbb{1}_{r>1/2} \right\|_{L_r^2 L_\sigma^4} + \left\| \frac{u}{r^{1/2}} \mathbb{1}_{r<1/2} \right\|_{L_r^2 L_\sigma^4} \\ &\leq \sqrt{2} \left\| u \mathbb{1}_{r>1/2} \right\|_{L_r^2 L_\sigma^4} + C \left\| u \mathbb{1}_{r<1/2} \right\|_{L^2}^{1/2-\eta} \left\| u \mathbb{1}_{r<1/2} \right\|_{L_r^2 L_\sigma^{p_1}}^\eta \left\| \frac{u}{r} \mathbb{1}_{r<1/2} \right\|_{L_r^2 L_\sigma^{p_1}}^{1/2} \\ &\leq C \|\langle \nabla_\sigma \rangle u\|_{L^2} + C \left\| u \mathbb{1}_{r<1/2} \right\|_{L^2}^{1/2-\eta} \|\langle \nabla_\sigma \rangle u\|_{L^2}^\eta \|u\|_{\mathbb{B}^1}^{1/2} \end{aligned}$$

where we used a Hölder inequality for the second inequality and (3.6), (3.8) for the third one. Here $\eta \in (0, 1/2)$ is a parameter that will be fixed later and $p_1 = 2 + 1/\eta$. It remains to use the properties of the confinement operator. From Assumption 1.3, we deduce

$$\begin{aligned} \left\| u \mathbb{1}_{r<1/2} \right\|_{L^2} &\leq 2^{\alpha/2} \left\| (r-1)^{\alpha/2} u \right\|_{L^2} \leq C \varepsilon^{\alpha/2} \left\| V_c \left(\frac{r-1}{\varepsilon} \right)^{1/2} u \right\|_{L^2} \\ &\leq C \varepsilon^{1+\alpha/2} \|u\|_{\mathbb{B}^1}. \end{aligned} \quad (3.10)$$

Hence,

$$\left\| \frac{u}{r^{1/2}} \right\|_{L_r^2 L_\sigma^4} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2} + C \varepsilon^{(1/2-\eta)(1+\alpha/2)} \|u\|_{\mathbb{B}^1}^{1-\eta} \|\langle \nabla_\sigma \rangle u\|_{L^2}^\eta$$

To conclude, we choose η small enough such that $(1/2-\eta)(1+\alpha/2) > 1-\eta$. This is possible thanks to the assumption $\alpha > 2$. The Young inequality finally gives (3.9).

Step 2. Proof of (3.1). Let $s \geq 1$. From the Littlewood-Paley theory and the Mihlin-Hörmander multiplier theorem [1, 26] applied on the sphere, we have

$$\begin{aligned} \|\langle \nabla_\sigma \rangle^s (\mathbb{G}(|u|^2)u)\|_{L^2} &\leq C \|\mathbb{G}(|u|^2)\|_{L^\infty} \|\langle \nabla_\sigma \rangle^s u\|_{L^2} \\ &\quad + C \|\langle \nabla_\sigma \rangle^s \mathbb{G}(|u|^2)\|_{L_\sigma^p} \|u\|_{L_r^2 L_\sigma^q} \end{aligned} \quad (3.11)$$

for all pair (p, q) such that $p > 2$ and $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$. The L^∞ norm of $\mathbb{G}(|u|^2)$ is easy to estimate. Indeed, this function is independent of the variable r and we notice that \mathbb{G} is linked to the operator G defined by (1.1) thanks to the relation

$$\mathbb{G}(|u|^2) = \int_0^{+\infty} G(|u|^2(r', \cdot)) r'^2 dr',$$

Pick a real number $p > 2$. From (2.8) and the Hölder inequality, we deduce

$$\begin{aligned} 0 \leq \mathbb{G}(|u|^2) &\leq C \int_0^{+\infty} \left\| u(r' \cdot) \right\|_{L_\sigma^{2p}}^{2p/(2p-2)} \left\| u(r' \cdot) \right\|_{L_\sigma^2}^{(2p-4)/(2p-2)} r'^2 dr' \\ &\leq C \|u\|_{L_r^2 L_\sigma^{2p}}^{2p/(2p-2)} \|u\|_{L^2}^{(2p-4)/(2p-2)} \\ &\leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 \end{aligned} \quad (3.12)$$

where we used the Sobolev embedding (3.6).

Let us now estimate the $L_r^\infty L_\sigma^p$ norm of $\langle \nabla_\sigma \rangle^s \mathbb{G}(|u|^2)$, where $p > 2$ is given. Since, from (2.15), we have

$$\mathbb{G}(|u|^2) = \int_0^{+\infty} (1 - 4\Delta_\sigma)^{-1/2} (|u|^2) r^2 dr,$$

the operator $-\Delta_\sigma$ commutes with \mathbb{G} and therefore

$$\langle \nabla_\sigma \rangle^s \mathbb{G}(|u|^2) = \mathbb{G}(\langle \nabla_\sigma \rangle^s(|u|^2)).$$

Thus, by (2.7),

$$\begin{aligned} \|\langle \nabla_\sigma \rangle^s \mathbb{G}(|u|^2)\|_{L_\sigma^p} &\leq C \int_0^{+\infty} \|\langle \nabla_\sigma \rangle^s(|u|^2)(r \cdot)\|_{L_\sigma^{p_2}} r^2 dr \\ &\leq C \int_0^{+\infty} \|\langle \nabla_\sigma \rangle^s u(r \cdot)\|_{L_\sigma^2} \|u(r \cdot)\|_{L_\sigma^p} r^2 dr \\ &\leq C \|\langle \nabla_\sigma \rangle^s u\|_{L^2} \|u\|_{L_r^2 L_\sigma^p} \end{aligned} \quad (3.13)$$

where p_2 was chosen such that $\frac{1}{p_2} = \frac{1}{p} + \frac{1}{2}$ and where the Mihklin-Hörmander multiplier theorem on the sphere was used again. Finally, from (3.11), (3.12), (3.13) and (3.6), we deduce (3.1).

Step 3. Proof of (3.2). Let $s \geq 1$. By the Mihklin-Hörmander multiplier theorem, we have

$$\begin{aligned} \|\langle \nabla_\sigma \rangle^s (V(|u|^2)u)\|_{L^2} &\leq C \|V(|u|^2)\|_{L^\infty} \|\langle \nabla_\sigma \rangle^s u\|_{L^2} \\ &\quad + C \left\| r^{1/2} \langle \nabla_\sigma \rangle^s V(|u|^2) \right\|_{L_r^\infty L_\sigma^4} \left\| \frac{u}{r^{1/2}} \right\|_{L_r^2 L_\sigma^4} \end{aligned} \quad (3.14)$$

Let us first estimate the L^∞ norm of $V(|u|^2)$, written in spherical coordinates:

$$V(|u|^2)(r\sigma) = \frac{1}{4\pi} \int \frac{1}{|r\sigma - r'\sigma'|} |u(r'\sigma')|^2 r'^2 dr' d\sigma'. \quad (3.15)$$

Since σ and σ' are unitary, we have $(\sigma - \sigma') \cdot (\sigma + \sigma') = 0$, thus

$$|r\sigma - r'\sigma'|^2 = \left(\frac{r+r'}{2}\right)^2 |\sigma - \sigma'|^2 + \left(\frac{r-r'}{2}\right)^2 |\sigma + \sigma'|^2, \quad (3.16)$$

which yields

$$\frac{1}{|r\sigma - r'\sigma'|} \leq \frac{2}{\max(r, r')} \frac{1}{|\sigma - \sigma'|}. \quad (3.17)$$

This enables to estimate $V(|u|^2)$ by using again Lemma 2.5. For all $p > 2$, we have

$$\begin{aligned} 0 \leq V(|u|^2) &\leq \int_0^{+\infty} \frac{2}{r'} G(|u|^2(r', \cdot)) r'^2 dr' \\ &\leq C \int_0^{+\infty} \frac{1}{r'} \|u(r', \cdot)\|_{L_\sigma^{2p}}^{2p/(2p-2)} \|u(r', \cdot)\|_{L_\sigma^2}^{(2p-4)/(2p-2)} r'^2 dr'. \end{aligned}$$

Hence, by splitting the integral on $r > 1/2$ and $r < 1/2$ and using the Hölder inequality,

$$\begin{aligned} 0 \leq V(|u|^2) &\leq C \|u \mathbb{1}_{r>1/2}\|_{L_r^2 L_\sigma^{2p}}^{2p/(2p-2)} \|u \mathbb{1}_{r>1/2}\|_{L^2}^{(2p-4)/(2p-2)} \\ &\quad + C \left\| \frac{u}{r} \mathbb{1}_{r<1/2} \right\|_{L_r^2 L_\sigma^{2p}} \|u \mathbb{1}_{r<1/2}\|_{L_r^2 L_\sigma^{2p}}^{2/(2p-2)} \|u \mathbb{1}_{r<1/2}\|_{L^2}^{(2p-4)/(2p-2)} \\ &\leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + C \|u\|_{\mathcal{B}^1} \|\langle \nabla_\sigma \rangle u\|_{L^2}^{2/(2p-2)} \|u \mathbb{1}_{r<1/2}\|_{L^2}^{(2p-4)/(2p-2)} \end{aligned}$$

where we have used again the Sobolev embeddings (3.6) and (3.8). Finally, (3.10) yields

$$0 \leq V(|u|^2) \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + C \varepsilon^{(1+\alpha/2)(1-\frac{2}{2p-2})} \|u\|_{\mathcal{B}^1}^{2-\frac{2}{2p-2}} \|\langle \nabla_\sigma \rangle u\|_{L^2}^{2/(2p-2)}.$$

To conclude, we choose p large enough such that $(1 + \alpha/2)(1 - \frac{2}{2p-2}) > 2 - \frac{2}{2p-2}$. This is possible thanks to the assumption $\alpha > 2$ and we obtain finally, using the Young inequality,

$$0 \leq V(|u|^2) \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + C \varepsilon^{2(1+\gamma)} \|u\|_{\mathbb{B}^1}^2. \quad (3.18)$$

Let us now estimate the $L_r^\infty L_\sigma^4$ norm of $r^{1/2} \langle \nabla_\sigma \rangle^s V(|u|^2)$. Since $V(|u|^2) = (-\Delta)^{-1}(|u|^2)$, the operator $-\Delta_\sigma$ commutes with V and we have

$$\langle \nabla_\sigma \rangle^s V(|u|^2) = V(\langle \nabla_\sigma \rangle^s(|u|^2)).$$

Therefore, we deduce from (3.15), from (3.17), from (2.7), and finally from the Mihklin-Hörmander theorem on the sphere that

$$\begin{aligned} \left\| r^{1/2} \langle \nabla_\sigma \rangle^s V(|u|^2) \right\|_{L_r^\infty L_\sigma^4} &\leq C \int_0^{+\infty} \frac{1}{r^{1/2}} \|\langle \nabla_\sigma \rangle^s(|u|^2)(r, \cdot)\|_{L_\sigma^{4/3}} r^2 dr \\ &\leq C \int_0^{+\infty} \|\langle \nabla_\sigma \rangle^s u(r, \cdot)\|_{L_\sigma^2} \left\| \frac{u(r, \cdot)}{r^{1/2}} \right\|_{L_\sigma^4} r^2 dr \\ &\leq C \|\langle \nabla_\sigma \rangle^s u\|_{L^2} \left\| \frac{u}{r^{1/2}} \right\|_{L_r^2 L_\sigma^4}. \end{aligned} \quad (3.19)$$

Finally, we deduce (3.2) from (3.14), (3.18), (3.19) and (3.9). The proofs of (3.3) and (3.4) are very similar to the proofs of (3.1) and (3.2), with $s = 1$ and we leave the details to the reader.

Step 4. Proof of (3.5). In order to estimate the L^2 norm of $\langle \nabla_\sigma \rangle (V(|u|^2)u - \mathbb{G}(|u|^2)u)$, let us introduce

$$\delta(r, r', \sigma, \sigma') := \frac{1}{|\sigma - \sigma'|} - \frac{1}{|r\sigma - r'\sigma'|},$$

such that, for all function w , we have

$$(V(w) - \mathbb{G}(w))(r\sigma) = - \int_0^{+\infty} \int_{\mathbb{S}^2} \delta(r, r', \sigma, \sigma') w(r'\sigma') r'^2 dr' d\sigma'. \quad (3.20)$$

From (3.17), one deduces directly that

$$|\delta(r, r', \sigma, \sigma')| \leq \frac{1}{|\sigma - \sigma'|} \left(1 + \frac{2}{\max(r, r')} \right). \quad (3.21)$$

Moreover, by using (3.16), one can decompose $\delta(r, r', \sigma, \sigma') = \delta_1 + \delta_2$ with

$$\delta_1 = \frac{((r + r')^2 - 4)|\sigma - \sigma'|^2}{4|r\sigma - r'\sigma'| |\sigma - \sigma'| (|r\sigma - r'\sigma'| + |\sigma - \sigma'|)}$$

and

$$\delta_2 = \frac{(r - r')^2 |\sigma + \sigma'|^2}{4|r\sigma - r'\sigma'| |\sigma - \sigma'| (|r\sigma - r'\sigma'| + |\sigma - \sigma'|)}.$$

Let χ be a continuous function on \mathbb{R} , positive for $z \neq 0$, such that $\chi(z) \sim |z|$ as $z \rightarrow 0$ and $\chi(z) = 1$ for $|z| \geq 2$. From (3.17), one deduces that

$$|\delta_1| \leq C \frac{(\chi(r-1) + \chi(r'-1))}{|\sigma - \sigma'|} \left(1 + \frac{1}{\max(r, r')} \right).$$

Furthermore, since by (3.16) we have

$$|r - r'| |\sigma + \sigma'| \leq 2|r\sigma - r'\sigma'|,$$

we can estimate δ_2 as follows:

$$|\delta_2| \leq \frac{|r - r'| |\sigma + \sigma'|}{2|r\sigma - r'\sigma'| |\sigma - \sigma'|} \leq C \frac{(\chi(r-1) + \chi(r'-1))}{|\sigma - \sigma'|^2} \left(1 + \frac{1}{\max(r, r')}\right).$$

We have thus proved that

$$|\delta(r, r', \sigma, \sigma')| \leq C \frac{(\chi(r-1) + \chi(r'-1))}{|\sigma - \sigma'|^2} \left(1 + \frac{1}{\max(r, r')}\right).$$

By interpolating between this inequality and (3.21), one gets finally, for all $\eta \in (0, 1)$,

$$|\delta(r, r', \sigma, \sigma')| \leq C \frac{\kappa(r, r')}{|\sigma - \sigma'|^{1+\eta}}, \quad (3.22)$$

where we have set

$$\kappa(r, r') = ((\chi(r-1))^\eta + (\chi(r'-1))^\eta) \left(1 + \frac{1}{\max(r, r')}\right).$$

Next, we claim that one can adapt Steps 2 and 3 in order to prove that, for $\eta > 0$ small enough,

$$\begin{aligned} & \|\langle \nabla_\sigma \rangle (V(|u|^2)u - \mathbb{G}(|u|^2)u)\|_{L^2} \\ & \leq C \|\chi(r-1)\langle \nabla_\sigma \rangle u\|_{L^2}^\eta \|\langle \nabla_\sigma \rangle u\|_{L^2}^{3-\eta} + C\varepsilon^\gamma (\|\langle \nabla_\sigma \rangle u\|_{L^2}^3 + \varepsilon^3 \|u\|_{\mathcal{B}^1}^3). \end{aligned} \quad (3.23)$$

Let us prove this claim. We have

$$\begin{aligned} & \|\langle \nabla_\sigma \rangle (V(|u|^2)u - \mathbb{G}(|u|^2)u)\|_{L^2} \\ & \leq C \left(\int_0^{+\infty} \|V(|u|^2) - \mathbb{G}(|u|^2)\|_{L^\infty_\sigma}^2 \|\langle \nabla_\sigma \rangle u\|_{L^2_\sigma}^2 r^2 dr \right)^{1/2} \\ & \quad + C \left(\int_0^{+\infty} \|V(\langle \nabla_\sigma \rangle(|u|^2)) - \mathbb{G}(\langle \nabla_\sigma \rangle(|u|^2))\|_{L^4_\sigma}^2 \|u\|_{L^4_\sigma}^2 r^2 dr \right)^{1/2} \end{aligned} \quad (3.24)$$

By (3.20) and (3.22), we have

$$\begin{aligned} |V(|u|^2) - \mathbb{G}(|u|^2)| & \leq C \int_0^{+\infty} \int_{\mathbb{S}^2} \frac{\kappa(r, r')}{|\sigma - \sigma'|^{1+\eta}} |u(r'\sigma')|^2 d\sigma' r'^2 dr' \\ & \leq C \int_0^{+\infty} ((\chi(r-1))^\eta + (\chi(r'-1))^\eta) \left(1 + \frac{1}{r'}\right) \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1+\eta)p}{p-1}} \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1-\eta)p-2}{p-1}} r'^2 dr' \\ & \leq C \int_{1/2}^{+\infty} ((\chi(r-1))^\eta + (\chi(r'-1))^\eta) \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1+\eta)p}{p-1}} \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1-\eta)p-2}{p-1}} r'^2 dr' \\ & \quad + C \int_0^{1/2} \frac{1}{r'} \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1+\eta)p}{p-1}} \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1-\eta)p-2}{p-1}} r'^2 dr' \end{aligned}$$

where we used the Hölder inequality and where $p > \frac{2}{1-\eta}$. The first integral in the last inequality can be bounded as in Step 2, by using the Sobolev embedding (3.6):

$$\begin{aligned} & \int_{1/2}^{+\infty} ((\chi(r-1))^\eta + (\chi(r'-1))^\eta) \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1+\eta)p}{p-1}} \|u(r'\cdot)\|_{L^{2p}_\sigma}^{\frac{(1-\eta)p-2}{p-1}} r'^2 dr' \\ & \leq (\chi(r-1))^\eta \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + \|\chi(r'-1)\langle \nabla_\sigma \rangle u\|_{L^2} \|\langle \nabla_\sigma \rangle u\|_{L^2}. \end{aligned}$$

The second integral can be bounded as in Step 3: by using (3.6), (3.8) and (3.10), we get, for η small enough and p large enough,

$$\int_0^{1/2} \frac{1}{r'} \|u(r'\cdot)\|_{L_\sigma^{2p}}^{\frac{(1+\eta)p}{p-1}} \|u(r'\cdot)\|_{L_\sigma^2}^{\frac{(1-\eta)p-2}{p-1}} r'^2 dr' \leq C\varepsilon^\gamma (\|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + \varepsilon^2 \|u\|_{\mathcal{B}^1}^2).$$

Finally, the first term in (3.24) can be estimated as follows:

$$\begin{aligned} & \left(\int_0^{+\infty} \|V(|u|^2) - \mathbb{G}(|u|^2)\|_{L_\sigma^\infty}^2 \|\langle \nabla_\sigma \rangle u\|_{L_\sigma^2}^2 r^2 dr \right)^{1/2} \\ & \leq C \|\chi(r' - 1)\|^\eta \|\langle \nabla_\sigma \rangle u\|_{L^2} \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + C\varepsilon^\gamma (\|\langle \nabla_\sigma \rangle u\|_{L^2}^3 + \varepsilon^3 \|u\|_{\mathcal{B}^1}^3). \end{aligned} \quad (3.25)$$

Let us now estimate the second term in (3.24). Setting

$$w(r', \sigma) = \int_{\mathbb{S}^2} \frac{1}{|\sigma - \sigma'|^{1+\eta}} \langle \nabla_\sigma \rangle (|u|^2)(r'\sigma') d\sigma',$$

we deduce from (3.20), (3.22) and from the Minkowski inequality that

$$\begin{aligned} & \int_0^{+\infty} \|V(\langle \nabla_\sigma \rangle (|u|^2)) - \mathbb{G}(\langle \nabla_\sigma \rangle (|u|^2))\|_{L_\sigma^4}^2 \|u\|_{L_\sigma^4}^2 r^2 dr \\ & \leq C \left(\int_0^{+\infty} \left(1 + \frac{1}{\sqrt{r'}}\right) \chi(r' - 1)^\eta \|w(r', \cdot)\|_{L_\sigma^4} r'^2 dr' \right)^2 \left(\int_0^{+\infty} \left(1 + \frac{1}{\sqrt{r}}\right)^2 \|u(r \cdot)\|_{L_\sigma^4}^2 r^2 dr \right) \\ & + C \left(\int_0^{+\infty} \left(1 + \frac{1}{\sqrt{r'}}\right) \|w(r', \cdot)\|_{L_\sigma^4} r'^2 dr' \right)^2 \left(\int_0^{+\infty} \left(1 + \frac{1}{\sqrt{r}}\right)^2 \chi(r - 1)^{2\eta} \|u(r \cdot)\|_{L_\sigma^4}^2 r^2 dr \right). \end{aligned}$$

By Hardy-Littlewood-Sobolev, one has

$$\|w(r', \cdot)\|_{L_\sigma^4} \leq C \|\langle \nabla_\sigma \rangle (|u|^2)\|_{L_\sigma^{\frac{4}{3-2\eta}}} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2} \|u\|_{L_\sigma^{\frac{4}{1-2\eta}}}$$

and, by adapting the proof of (3.9), one can prove that there exists $\gamma > 0$ such that, for η small enough,

$$\left\| \frac{u}{\sqrt{r}} \right\|_{L_r^2 L_\sigma^{\frac{4}{1-2\eta}}} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2} + C\varepsilon^{1+\gamma} \|u\|_{\mathcal{B}^1}. \quad (3.26)$$

Finally, using (3.6) and (3.26), one gets

$$\begin{aligned} & \left(\int_0^{+\infty} \|V(\langle \nabla_\sigma \rangle (|u|^2)) - \mathbb{G}(\langle \nabla_\sigma \rangle (|u|^2))\|_{L_\sigma^4}^2 \|u\|_{L_\sigma^4}^2 r^2 dr \right)^{1/2} \\ & \leq C \|\chi(r - 1)\|^\eta \|\langle \nabla_\sigma \rangle u\|_{L^2} (\|\langle \nabla_\sigma \rangle u\|_{L^2}^2 + C\varepsilon^{2+2\gamma} \|u\|_{\mathcal{B}^1}^2) \end{aligned} \quad (3.27)$$

and the claim (3.23) can be deduced from (3.24), (3.25), (3.27), the Young inequality and the Hölder inequality.

We are in position to conclude the proof of (3.5). By using an interpolation inequality on the sphere and the properties of the truncation function χ , we obtain

$$\begin{aligned} \|\chi(r - 1)\langle \nabla_\sigma \rangle u\|_{L^2} & \leq C \|\chi(r - 1)u\|_{L^2}^{1/2} \|\langle \nabla_\sigma \rangle^2 u\|_{L^2}^{1/2} \\ & \leq C \| |r - 1|^{\alpha/2} u \|_{L^2}^{1/\alpha} \|u\|_{L^2}^{1/2-1/\alpha} \|\langle \nabla_\sigma \rangle^2 u\|_{L^2}^{1/2} \\ & \leq C\varepsilon^{1/2} \left\| V_c \left(\frac{r-1}{\varepsilon} \right)^{1/2} u \right\|^{1/\alpha} \|u\|_{L^2}^{1/2-1/\alpha} \|\langle \nabla_\sigma \rangle^2 u\|_{L^2}^{1/2} \\ & \leq C\varepsilon^{1/2} (\varepsilon \|u\|_{\mathcal{B}^1})^{1/\alpha} \|\langle \nabla_\sigma \rangle^2 u\|_{L^2}^{1-1/\alpha} \end{aligned}$$

where we used Assumption 1.3 and the fact that $\alpha \geq 2$. Finally, inserting this estimate in (3.23) and using the Young inequality leads to (3.5), up to changing the value of γ . \square

Let us now state another estimate, where we recall that the operator H_r was defined in (1.11).

Lemma 3.2. *Let u be such that $\langle \nabla_\sigma \rangle u \in L^2$ and $H_r^{s/2} \langle \nabla_\sigma \rangle u \in L^2$ with $s \geq 1$ an integer, then*

$$\left\| (1 + \varepsilon^2 H_r)^{s/2} \langle \nabla_\sigma \rangle (\mathbb{G}(|u|^2)u) \right\|_{L^2} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 \|(1 + \varepsilon^2 H_r)^{s/2} \langle \nabla_\sigma \rangle u\|_{L^2}. \quad (3.28)$$

Proof. Since the operator H_r only acts on the variable r and since \mathbb{G} is independent of the variable r , we have, for all $s \geq 0$,

$$\begin{aligned} \|(1 + \varepsilon^2 H_r)^{s/2} \langle \nabla_\sigma \rangle (\mathbb{G}(|u|^2)u)\|_{L^2} &= \|\langle \nabla_\sigma \rangle (\mathbb{G}(|u|^2)(1 + \varepsilon^2 H_r)^{s/2} u)\|_{L^2} \\ &\leq \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 \|(1 + \varepsilon^2 H_r)^{s/2} \langle \nabla_\sigma \rangle u\|_{L^2}, \end{aligned}$$

where we used (3.12) and the following inequality, that can be proved as (3.1) with $s = 1$:

$$\|\langle \nabla_\sigma \rangle (\mathbb{G}(|u|^2)v)\|_{L^2} \leq C \|\langle \nabla_\sigma \rangle u\|_{L^2}^2 \|\langle \nabla_\sigma \rangle v\|_{L^2}.$$

\square

We end this section with the following lemma, which will enable to deal with more convenient norms than $\|(1 + \varepsilon^2 H_r)^{s/2} u\|_{L^2}$.

Lemma 3.3. *Assume that the confinement potential satisfies Assumption 1.3 and let H_r be defined by (1.11). Then H_r is a positive selfadjoint operator on $L^2(\mathbb{R}_+, r^2 dr)$ and, for every integer $s \geq 1$,*

$$D((1 + H_r)^{s/2}) = \left\{ u \in L^2(\mathbb{R}_+, r^2 dr) : (V_c^\varepsilon)^{\frac{s-k}{2}} \partial_r^k(ru) \in L^2(\mathbb{R}_+, dr), 0 \leq k \leq s \right\}.$$

Moreover, the following norms are equivalent, with constants independent of ε :

$$\begin{aligned} &\left\| (1 + \varepsilon^2 H_r)^{\frac{s}{2}} u \right\|_{L^2}, \\ &\|u\|_{L^2} + \varepsilon^s \left\| \frac{1}{r} \partial_r^s(ru) \right\|_{L^2} + \varepsilon^s \|(V_c^\varepsilon)^{\frac{s}{2}} u\|_{L^2}, \\ &\|u\|_{L^2} + \varepsilon^s \sum_{k=0}^s \left\| (V_c^\varepsilon)^{\frac{s-k}{2}} \frac{1}{r} \partial_r^k(ru) \right\|_{L^2} \quad \text{and} \quad \|u\|_{L^2} + \varepsilon^s \sum_{k=0}^s \left\| \frac{1}{r} \partial_r^k \left(r (V_c^\varepsilon)^{\frac{s-k}{2}} u \right) \right\|_{L^2}, \end{aligned}$$

where $\|\cdot\|_{L^2}$ denotes the $L^2(\mathbb{R}_+, r^2 dr)$ norm.

Note that proving this result is equivalent to proving the equivalence of the following norms in dimension one:

$$\begin{aligned} &\left\| (1 + \varepsilon^2 (-\partial_r^2 + V_c^\varepsilon))^{\frac{s}{2}} u \right\|_{L^2(\mathbb{R}_+, dr)}, \\ &\|u\|_{L^2(\mathbb{R}_+, dr)} + \varepsilon^s \|\partial_r^s u\|_{L^2(\mathbb{R}_+, dr)} + \varepsilon^s \|(V_c^\varepsilon)^{\frac{s}{2}} u\|_{L^2(\mathbb{R}_+, dr)}, \\ &\|u\|_{L^2(\mathbb{R}_+, dr)} + \varepsilon^s \sum_{k=0}^s \left\| (V_c^\varepsilon)^{\frac{s-k}{2}} \partial_r^k u \right\|_{L^2(\mathbb{R}_+, dr)}, \end{aligned}$$

$$\|u\|_{L^2(\mathbb{R}_+, dr)} + \varepsilon^s \sum_{k=0}^s \left\| \partial_r^k \left(V_c^\varepsilon \right)^{\frac{s-k}{2}} u \right\|_{L^2(\mathbb{R}_+, dr)}.$$

Now, recalling that $V_c^\varepsilon(r) = \frac{1}{\varepsilon^2} V_c\left(\frac{r-1}{\varepsilon}\right)$, let us apply the dilation $r' = \frac{r-1}{\varepsilon}$. The above norms become

$$\begin{aligned} & \left\| (1 - \partial_r^2 + V_c)^{\frac{s}{2}} u \right\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)}, \\ & \|u\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)} + \|\partial_r^s u\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)} + \|(V_c)^{\frac{s}{2}} u\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)}, \\ & \|u\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)} + \sum_{k=0}^s \left\| (V_c)^{\frac{s-k}{2}} \partial_r^k u \right\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)}, \\ & \|u\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)} + \sum_{k=0}^s \left\| \partial_r^k \left(V_c \right)^{\frac{s-k}{2}} u \right\|_{L^2\left(-\frac{1}{\varepsilon}, +\infty\right)}. \end{aligned}$$

Hence, Lemma 3.3 is a consequence of Proposition A.1 proved in Appendix A.

3.2. Approximation by an intermediate model. In this subsection, we make a first step towards Theorem 1.5. We obtain a priori estimates for the singularly perturbed system 1.4 and prove that it can be approximated by the following system, where we only pass to the limit in the nonlinear term:

$$i\partial_t w^\varepsilon = -\Delta w^\varepsilon + V_c^\varepsilon w^\varepsilon + \mathbb{G}(|w^\varepsilon|^2) w^\varepsilon, \quad w^\varepsilon(t=0) = u_0^\varepsilon, \quad (3.29)$$

where \mathbb{G} is defined by (1.15).

Proposition 3.4. *Assume that the confinement potential and the initial data satisfy Assumptions 1.3 and 1.4. Then the following holds true.*

(i) *For all $\varepsilon \in (0, 1]$, (1.4) admits a unique solution u^ε in the energy space $\mathcal{C}^0(\mathbb{R}, \mathcal{B}^1)$. Moreover, there exists $T > 0$ such that*

$$\sup_{\varepsilon \in (0, 1]} (\varepsilon \|u^\varepsilon\|_{L^\infty([-T, T], \mathcal{B}^1)} + \|\langle \nabla_\sigma \rangle u^\varepsilon\|_{L^\infty([-T, T], L^2)}) < +\infty. \quad (3.30)$$

(ii) *There exists $T > 0$ such that, for all $\varepsilon \in (0, 1]$, (3.29) admits a unique solution $w^\varepsilon \in \mathcal{C}^0([-T, T], \mathcal{B}^1)$ with $\langle \nabla_\sigma \rangle w^\varepsilon \in \mathcal{C}^0([-T, T], L^2)$, and with a uniform bound:*

$$\sup_{\varepsilon \in (0, 1]} (\varepsilon \|w^\varepsilon\|_{L^\infty([-T, T], \mathcal{B}^1)} + \|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T, T], L^2)}) < +\infty. \quad (3.31)$$

(iii) *Assume that $\varepsilon_0 \in (0, 1]$ and $T > 0$ are such that*

$$\sup_{\varepsilon \in (0, \varepsilon_0]} (\varepsilon \|w^\varepsilon\|_{L^\infty([-T, T], \mathcal{B}^1)} + \|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T, T], L^2)}) < +\infty. \quad (3.32)$$

Then, one has

$$\lim_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (u^\varepsilon - w^\varepsilon)(t)\|_{L^\infty([-T, T], L^2)} = 0. \quad (3.33)$$

Proof. The well-posedness of the Cauchy problem in the energy space \mathcal{B}^1 for (1.4), for all fixed $\varepsilon > 0$, is very standard. It can be proved by using standard techniques: local in time existence by a Banach fixed-point procedure, then global existence thanks to the conservation laws (1.10) and (1.11).

Step 1: a priori estimate (3.30). Let us now prove the existence of $T > 0$ such that the a priori estimate (3.30) holds. Thanks to Assumption 1.4, we have

$$\|\nabla u_0^\varepsilon\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} u_0^\varepsilon\|_{L^2}^2 \leq \|u_0^\varepsilon\|_{\mathcal{B}^1}^2 \leq \frac{C}{\varepsilon^2} \quad (3.34)$$

and

$$\|\nabla V(|u_0^\varepsilon|^2)\|_{L^2}^2 = \int_{\mathbb{R}^3} V(|u_0^\varepsilon|^2)|u_0^\varepsilon|^2 dx \leq \|V(|u_0^\varepsilon|^2)\|_{L^\infty} \|u_0^\varepsilon\|_{L^2}^2 \leq C,$$

where we used (3.18). Thus, the energy conservation law (1.11) yields, for all t ,

$$\varepsilon^2 \|u^\varepsilon(t)\|_{\mathcal{B}^1}^2 \leq \varepsilon^2 \|u_0\|_{\mathcal{B}^1}^2 + \frac{\varepsilon^2}{2} \|\nabla V(|u_0^\varepsilon|^2)\|_{L^2}^2 \leq C. \quad (3.35)$$

Apply now the operator $\langle \nabla_\sigma \rangle = \sqrt{1 - \Delta_\sigma}$ to the Schrödinger equation (1.4). Since this operator commutes with the Hamiltonian, we get for all t

$$\begin{aligned} \|\langle \nabla_\sigma \rangle u^\varepsilon(t)\|_{L^2} &\leq \|\langle \nabla_\sigma \rangle u_0^\varepsilon\|_{L^2} + \int_{-|t|}^{|t|} \|\langle \nabla_\sigma \rangle (V(|u^\varepsilon|^2)u^\varepsilon)\|_{L^2}(\tau) d\tau \\ &\leq C + C \int_{-|t|}^{|t|} (\|\langle \nabla_\sigma \rangle u^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|u^\varepsilon\|_{\mathcal{B}^1}^2) \|\langle \nabla_\sigma \rangle u^\varepsilon\|_{L^2} d\tau, \end{aligned}$$

where we used Assumption 1.4 and (3.2). Using the estimate (3.35) and a standard bootstrap lemma, this yields a local in time estimate: there exist T and C_T independent of ε such that

$$\forall t \in [-T, T] \quad \|\langle \nabla_\sigma \rangle u^\varepsilon(t)\|_{L^2} \leq C_T. \quad (3.36)$$

The proof of Item (i) of Proposition 3.4 is complete.

Step 2: the Cauchy problem for the intermediate model. Let us now consider the Cauchy problem for (3.29). By using (3.1) with $s = 1$, it is easy to prove by a fixed-point procedure that, for all $\varepsilon > 0$, (3.29) admits a unique maximal solution w^ε such that $\langle \nabla_\sigma \rangle w^\varepsilon \in \mathcal{C}^0((-T^\varepsilon, T^\varepsilon), L^2)$. Note that, by a bootstrap argument similar as above, one can prove that T^ε is bounded from below independently of ε : there exists $T > 0$ such that

$$\sup_{\varepsilon \in (0, \varepsilon_0]} \|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T, T], L^2)} < +\infty.$$

Moreover, this solution w^ε also belongs to the energy space for all time and satisfies the mass and energy conservation laws:

$$\|w^\varepsilon(t)\|_{L^2}^2 = \|u_0^\varepsilon\|_{L^2}^2 \quad (3.37)$$

$$\begin{aligned} \|\nabla w^\varepsilon(t)\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} w^\varepsilon(t)\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{R}^3} \mathbb{G}(|w^\varepsilon(t, x)|^2) |w^\varepsilon(t, x)|^2 dx \\ = \|\nabla u_0^\varepsilon\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} u_0^\varepsilon\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{R}^3} \mathbb{G}(|u_0^\varepsilon|^2) |u_0^\varepsilon|^2 dx. \end{aligned}$$

Notice that this energy is finite and of order $\frac{1}{\varepsilon^2}$ at the initial time. Indeed, the two first terms in the right-hand side are bounded by (3.34) and the third term is bounded thanks to Assumption 1.4 and (3.12):

$$\int_{\mathbb{R}^3} \mathbb{G}(|u_0^\varepsilon|^2) |u_0^\varepsilon|^2 dx \leq \|\mathbb{G}(|u_0^\varepsilon|^2)\|_{L^\infty} \|u_0^\varepsilon\|_{L^2}^2 \leq C.$$

Notice also that, to solve this problem (3.29), it was crucial to bound the L^∞ norm of $\mathbb{G}(|u|^2)$, at the initial time and during the evolution, locally in time. Thanks to (3.12), such estimate is available as soon as $\langle \nabla_\sigma \rangle u$ belongs to L^2 . This is the reason why we introduced Assumption (1.12) on the initial data. It is not clear whether the Cauchy problem for (3.29) is well-posed on H^1 (or \mathcal{B}^1) only.

Step 3: regularization and approximation result. Let us now prove (3.33). Let $\varepsilon_0 > 0$ and $T > 0$ be such that a uniform bound (3.32) holds. We set

$$M := 1 + \sup_{\varepsilon \in (0, \varepsilon_0]} (\varepsilon \|w^\varepsilon\|_{L^\infty([-T, T], \mathbb{B}^1)} + \|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T, T], L^2)}) < +\infty.$$

Remark in particular that $\|\langle \nabla_\sigma \rangle u_0^\varepsilon\|_{L^2} \leq M < 2M$. Let T_1^ε be defined as follows:

$$T_1^\varepsilon = \sup \{ \tau \in (0, T] : \|\langle \nabla_\sigma \rangle u^\varepsilon\|_{L^\infty([- \tau, \tau], L^2)} < 2M \}.$$

From Step 1, we know that T_1^ε is bounded from below: there exists $T_1 > 0$ such that $T_1^\varepsilon > T_1$ for all $\varepsilon \in (0, \varepsilon_0]$. Moreover, by a continuity argument, one can prove that if $T_1^\varepsilon < T$, then

$$\|\langle \nabla_\sigma \rangle u^\varepsilon\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} = 2M.$$

Let $\delta > 0$. From Assumption (1.13) on the initial data, one can define a regularized subsequence $u_0^{\varepsilon, \delta}$ of the initial data such that

$$\sup_\varepsilon \left(\|\langle \nabla_\sigma \rangle^2 u_0^{\varepsilon, \delta}\|_{L^2} + \varepsilon \|u_0^{\varepsilon, \delta}\|_{\mathbb{B}^1} \right) < +\infty \quad (3.38)$$

and

$$\limsup_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (u_0^\varepsilon - u_0^{\varepsilon, \delta})\| \leq \delta. \quad (3.39)$$

Let $u^{\varepsilon, \delta}$, $w^{\varepsilon, \delta}$ be the solutions of (1.4) and (3.29) with $u_0^{\varepsilon, \delta}$ as initial data. By standard arguments, using (3.3) and (3.4), one can prove that these solutions depend continuously on the initial data: for δ small enough, (3.38), (3.39) yield

$$\limsup_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (w^{\varepsilon, \delta} - w^\varepsilon)\|_{L^\infty([-T, T], L^2)} \leq C\delta, \quad (3.40)$$

$$\limsup_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (u^{\varepsilon, \delta} - u^\varepsilon)\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} \leq C\delta, \quad (3.41)$$

where C is independent of ε and where we used that $\varepsilon \|u^{\varepsilon, \delta}\|_{\mathbb{B}^1}$ and $\varepsilon \|u^\varepsilon\|_{\mathbb{B}^1}$ are uniformly bounded. In particular, if ε_0 and δ have been initially chosen small enough, we have the estimates

$$\sup_{\varepsilon \in (0, \varepsilon_0]} \|\langle \nabla_\sigma \rangle u^{\varepsilon, \delta}\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} + \sup_{\varepsilon \in (0, \varepsilon_0]} \|\langle \nabla_\sigma \rangle w^{\varepsilon, \delta}\|_{L^\infty([-T, T], L^2)} \leq 4M. \quad (3.42)$$

Next, applying the operator $\langle \nabla_\sigma \rangle^2$ to (1.4) and (3.29), then using the tame estimates (3.2) and (3.1), with $s = 2$, one deduces thanks to (3.42) and the Gronwall lemma that, for all $\varepsilon \in (0, \varepsilon_0]$,

$$\|\langle \nabla_\sigma \rangle^2 u^{\varepsilon, \delta}\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} + \|\langle \nabla_\sigma \rangle^2 w^{\varepsilon, \delta}\|_{L^\infty([-T, T], L^2)} \leq C_M \|\langle \nabla_\sigma \rangle^2 u_0^{\varepsilon, \delta}\|_{L^2} \leq C_{M, \delta} \quad (3.43)$$

where C_M and $C_{M, \delta}$ are generic constants which only depend, respectively, on M and on (M, δ) , and where we used the bound (3.38).

Let us now write the equation satisfied by the difference $z = u^{\varepsilon, \delta} - w^{\varepsilon, \delta}$:

$$\begin{aligned} i\partial_t z &= -\Delta z + V_c^\varepsilon z + V(|u^{\varepsilon, \delta}|^2)u^{\varepsilon, \delta} - \mathbb{G}(|u^{\varepsilon, \delta}|^2)u^{\varepsilon, \delta} \\ &\quad + \mathbb{G}(|u^{\varepsilon, \delta}|^2)u^{\varepsilon, \delta} - \mathbb{G}(|w^{\varepsilon, \delta}|^2)w^{\varepsilon, \delta}. \end{aligned} \quad (3.44)$$

From (3.5) and (3.43) (and using also the energy estimates), we deduce that

$$\|\langle \nabla_\sigma \rangle (V(|u^{\varepsilon, \delta}|^2)u^{\varepsilon, \delta} - \mathbb{G}(|u^{\varepsilon, \delta}|^2)u^{\varepsilon, \delta})\|_{L^2} \leq \varepsilon^\gamma C_{M, \delta}.$$

Moreover, since $\langle \nabla_\sigma \rangle u^{\varepsilon, \delta}$ and $\langle \nabla_\sigma \rangle w^{\varepsilon, \delta}$ are uniformly bounded in L^2 , one deduces from (3.3) that, for all t and ε ,

$$\|\langle \nabla_\sigma \rangle (\mathbb{G}(|u^{\varepsilon, \delta}|^2)u^{\varepsilon, \delta} - \mathbb{G}(|w^{\varepsilon, \delta}|^2)w^{\varepsilon, \delta})\|_{L^2} \leq C_M \|\langle \nabla_\sigma \rangle z(t)\|_{L^2}.$$

Finally, since $z(t=0) = 0$, applying $\langle \nabla_\sigma \rangle$ to (3.44) leads to

$$\|\langle \nabla_\sigma \rangle z(t)\|_{L^2} \leq \varepsilon^\gamma C_{M, \delta} T + C_M \int_{-|t|}^{|t|} \|\langle \nabla_\sigma \rangle z(\tau)\|_{L^2} d\tau,$$

and a Gronwall lemma enables to conclude that

$$\|\langle \nabla_\sigma \rangle z\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} \leq \varepsilon^\gamma C_{M, \delta} T e^{C_M T}.$$

From this inequality and from (3.40), (3.41), letting δ tend to zero, one deduces that

$$\lim_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (u^\varepsilon - w^\varepsilon)\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} = 0.$$

Finally, by fixing $\varepsilon_1 \in (0, \varepsilon_0]$ small enough such that

$$\sup_{\varepsilon \in (0, \varepsilon_1]} \|\langle \nabla_\sigma \rangle (u^\varepsilon - w^\varepsilon)\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} < \frac{M}{2},$$

we ensure that

$$\sup_{\varepsilon \in (0, \varepsilon_1]} \|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T_1^\varepsilon, T_1^\varepsilon], L^2)} \leq \frac{3M}{2} < 2M$$

(since $M > 0$). We deduce from this inequality that, for all $\varepsilon \in (0, \varepsilon_1]$, $T_1^\varepsilon = T$. The proof of the proposition is complete. \square

3.3. Proof of Theorem 1.5. In order to prove Theorem 1.5, it remains to prove that the solution of the intermediate model (3.29) can be approximated by the solution of the limit model (1.14). Notice that (3.29) reads

$$i\partial_t w^\varepsilon = H_r w^\varepsilon - \frac{1}{r^2} \Delta_\sigma w^\varepsilon + \mathbb{G}(|w^\varepsilon|^2) w^\varepsilon, \quad w^\varepsilon(t=0) = u_0^\varepsilon.$$

Proposition 3.5. *Assume that the confinement potential and the initial data satisfy Assumptions 1.3 and 1.4. Then the following holds true.*

(i) *For all $\varepsilon > 0$, the limit system (1.14) admits a unique global solution v^ε such that $\langle \nabla_\sigma \rangle v^\varepsilon \in C^0(\mathbb{R}, L^2)$. Moreover, the following conservation laws are satisfied for all $t \in \mathbb{R}$:*

$$\|v^\varepsilon(t)\|_{L^2}^2 = \|u_0^\varepsilon\|_{L^2}^2, \quad (3.45)$$

$$\|\partial_r v^\varepsilon(t)\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} v^\varepsilon(t)\|_{L^2}^2 = \|\partial_r u_0^\varepsilon\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2} u_0^\varepsilon\|_{L^2}^2, \quad (3.46)$$

$$\|\nabla_\sigma v^\varepsilon\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{R}^3} \mathbb{G}(|v^\varepsilon|^2) |v^\varepsilon|^2 dx = \|\nabla_\sigma u_0^\varepsilon\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{R}^3} \mathbb{G}(|u_0^\varepsilon|^2) |u_0^\varepsilon|^2 dx. \quad (3.47)$$

(ii) *Let $T > 0$. Then there exists $\varepsilon_0 > 0$ such that the intermediate model (3.29) admits a unique solution w^ε on $[-T, T]$ with a uniform bound (3.32). Moreover, one has*

$$\lim_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (w^\varepsilon - v^\varepsilon)(t)\|_{L^\infty([-T, T], L^2)} = 0. \quad (3.48)$$

Proof. Thanks to (3.1), (3.3) and (3.28) (with $s = 1$) it is easy to prove that the Cauchy problem (1.14) is well-posed locally in time. In fact, the solution will be global thanks to (3.45), (3.46) and (3.47). Let us now prove these conservation laws. The first one (3.45) is the standard conservation of the L^2 norm. The second one (3.46) is the conservation of the L^2 norm for the equation satisfied by $H_r^{1/2}v^\varepsilon$:

$$i\partial_t H_r^{1/2}v^\varepsilon = H_r H_r^{1/2}v^\varepsilon - \Delta_\sigma H_r^{1/2}v^\varepsilon + \mathbb{G}(|v^\varepsilon|^2) H_r^{1/2}v^\varepsilon, \quad H_r^{1/2}v^\varepsilon(t=0) = H_r^{1/2}u_0^\varepsilon,$$

recalling that $\mathbb{G}(\cdot)$ is independent of r and that

$$H_r = -\frac{1}{r^2}\partial_r(r^2\partial_r) + V_c^\varepsilon, \quad \|(H_r)^{1/2}u\|_{L^2}^2 = \|\partial_r u\|_{L^2}^2 + \|(V_c^\varepsilon)^{1/2}v^\varepsilon\|_{L^2}^2.$$

The third identity (3.47) is obtained by multiplying (1.14) by $\partial_t \overline{v^\varepsilon}$, integrating on \mathbb{R}^3 , taking the real part of the equation and finally using (3.46).

Let us now prove Item (ii). Let $T > 0$ and denote

$$M_0 = 1 + \sup_{\varepsilon \in (0,1]} \left(\|u_0^\varepsilon\|_{L^2}^2 + \|\nabla_\sigma u_0^\varepsilon\|_{L^2}^2 + \frac{1}{2} \int_{\mathbb{R}^3} \mathbb{G}(|u_0^\varepsilon|^2) |u_0^\varepsilon|^2 dx \right) < +\infty.$$

By (3.45) and (3.47), we have, for all $\varepsilon > 0$,

$$\|\langle \nabla_\sigma \rangle v^\varepsilon\|_{L^\infty([-T,T],L^2)} \leq M_0. \quad (3.49)$$

By Proposition 3.4, the Cauchy problem for the intermediate model (3.29) is locally well-posed for $0 < \varepsilon < 1$, with a uniform bound of the form (3.31). Denote by w^ε its solution and set

$$T_0^\varepsilon = \sup \{ \tau \in (0, T] : \|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([- \tau, \tau], L^2)} < 2M_0 \}. \quad (3.50)$$

We already know that T_0^ε is bounded from below: there exists $T_0 > 0$ such that $T_0^\varepsilon > T_0$ for all $\varepsilon \in (0, 1]$. Moreover, by a continuity argument, one can prove that if $T_0^\varepsilon < T$, then

$$\|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T_0^\varepsilon, T_0^\varepsilon], L^2)} = 2M_0.$$

Let us now regularize the initial data as follows. From (1.13) again, we deduce that, for all $\delta > 0$, there exists a subsequence of the initial data u_0^ε , still denoted $u_0^{\varepsilon,\delta}$, such that

$$\sup_\varepsilon \left(\|\langle \nabla_\sigma \rangle^6 u_0^{\varepsilon,\delta}\|_{L^2} + \|(1 + \varepsilon H_r)^{3/2} \langle \nabla_\sigma \rangle u_0^{\varepsilon,\delta}\|_{L^2} \right) < +\infty \quad (3.51)$$

and

$$\limsup_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (u_0^\varepsilon - u_0^{\varepsilon,\delta})\| \leq \delta. \quad (3.52)$$

We denote by $v^{\varepsilon,\delta}$ the corresponding solution of (1.14). By standard arguments, using (3.51), (3.52) and (3.3), one can prove that

$$\limsup_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (v^{\varepsilon,\delta} - v^\varepsilon)\|_{L^\infty([-T,T],L^2)} \leq C\delta. \quad (3.53)$$

In particular, for δ and ε_0 small enough we have the estimate

$$\sup_{\varepsilon \in (0, \varepsilon_0]} \|\langle \nabla_\sigma \rangle v^{\varepsilon,\delta}\|_{L^\infty([-T,T],L^2)} \leq 4M_0. \quad (3.54)$$

Next, applying the operator $\langle \nabla_\sigma \rangle^6$ to (1.14), then using the tame estimate (3.1) with $s = 6$,

$$\begin{aligned} \|\langle \nabla_\sigma \rangle^6 v^{\varepsilon, \delta}(t)\|_{L^2} &\leq C \|\langle \nabla_\sigma \rangle^6 u_0^{\varepsilon, \delta}\|_{L^2} + C \int_{-|t|}^{|t|} \|\langle \nabla_\sigma \rangle v^{\varepsilon, \delta}(\tau)\|_{L^2}^2 \|\langle \nabla_\sigma \rangle^6 v^{\varepsilon, \delta}(\tau)\| d\tau \\ &\leq C_{M_0, \delta} + C_{M_0} \int_{-|t|}^{|t|} \|\langle \nabla_\sigma \rangle^6 v^{\varepsilon, \delta}(\tau)\| d\tau. \end{aligned}$$

Hence the Gronwall lemma yields

$$\|\langle \nabla_\sigma \rangle^6 v^{\varepsilon, \delta}\|_{L^\infty([-T, T], L^2)} \leq C_{M_0, \delta}. \quad (3.55)$$

Similarly, applying the operator $(1 + \varepsilon H_r)^{3/2} \langle \nabla_\sigma \rangle$ to (1.14), then using the estimate (3.28), with $s = 4$ leads to

$$\begin{aligned} \|(1 + \varepsilon^2 H_r)^{3/2} \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}(t)\|_{L^2} &\leq C \|(1 + \varepsilon^2 H_r)^{3/2} \langle \nabla_\sigma \rangle u_0^{\varepsilon, \delta}\|_{L^2} \\ &\quad + C \int_{-|t|}^{|t|} \|\langle \nabla_\sigma \rangle v^{\varepsilon, \delta}(\tau)\|_{L^2}^2 \|(1 + \varepsilon^2 H_r)^{3/2} \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}(\tau)\| d\tau \\ &\leq C_{M_0, \delta} + C_{M_0, \delta} \int_{-|t|}^{|t|} \|(1 + \varepsilon^2 H_r)^{3/2} \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}(\tau)\| d\tau. \end{aligned}$$

The crucial point for this estimate was that the operators $(1 + \varepsilon H_r)^{3/2} \langle \nabla_\sigma \rangle$ and $H_r - \Delta_\sigma$ commute together (whereas $(1 + \varepsilon H_r)^{3/2} \langle \nabla_\sigma \rangle$ does not commute with the complete Laplace operator Δ that appears in the intermediate model (3.29)). Hence the Gronwall lemma yields

$$\|(1 + \varepsilon^2 H_r)^{3/2} \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}\|_{L^\infty([-T, T], L^2)} \leq C_{M_0, \delta}. \quad (3.56)$$

With these estimates, we are now ready to conclude. Let us introduce a smooth function χ , defined on \mathbb{R}_+ , such that $0 \leq \chi \leq 1$, $\chi(r) = 0$ for $r \leq 1/3$ and $\chi(r) = 1$ for $r \geq 2/3$. Since the support of $1 - \chi$ is $\{r \leq 2/3\}$, one has

$$\begin{aligned} \|(1 - \chi) \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}\|_{L^2} &\leq C \| |r - 1|^\alpha \langle \nabla_\sigma \rangle v^{\varepsilon, \delta} \|_{L^2} \\ &\leq C \varepsilon^\alpha \|\varepsilon^2 V_c^\varepsilon \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}\|_{L^2} \\ &\leq C \varepsilon^\alpha \|(1 + \varepsilon^2 H_r) \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}\|_{L^2} \leq \varepsilon^\alpha C_{M_0, \delta} \end{aligned} \quad (3.57)$$

where we used Assumption (1.7), Lemma 3.3 and (3.56). Moreover, the function $\chi v^{\varepsilon, \delta}$ satisfies the equation

$$i \partial_t (\chi v^{\varepsilon, \delta}) = H_r (\chi v^{\varepsilon, \delta}) - \Delta_\sigma (\chi v^{\varepsilon, \delta}) + \mathbb{G}(|v^{\varepsilon, \delta}|^2) (\chi v^{\varepsilon, \delta}) + R^\varepsilon$$

where the remainder

$$R^\varepsilon = -2\chi' \partial_r v^{\varepsilon, \delta} - \left(\chi'' + \frac{2}{r} \chi' \right) v^{\varepsilon, \delta}$$

can be estimated as follows:

$$\begin{aligned} \|\langle \nabla_\sigma \rangle R^\varepsilon\|_{L^2} &\leq C \left\| |r - 1|^\alpha \langle \nabla_\sigma \rangle \frac{1}{r} \partial_r (r v^{\varepsilon, \delta}) \right\|_{L^2} + C \| |r - 1|^\alpha \langle \nabla_\sigma \rangle v^{\varepsilon, \delta} \|_{L^2} \\ &\leq C \varepsilon^{\alpha-1} \left\| \varepsilon^3 V_c^\varepsilon \langle \nabla_\sigma \rangle \frac{1}{r} \partial_r (r v^{\varepsilon, \delta}) \right\|_{L^2} + C \varepsilon^\alpha \|\varepsilon^2 V_c^\varepsilon \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}\|_{L^2} \\ &\leq C \varepsilon^{\alpha-1} \|(1 + \varepsilon^2 H_r)^{3/2} \langle \nabla_\sigma \rangle v^{\varepsilon, \delta}\|_{L^2} \end{aligned} \quad (3.58)$$

$$\leq \varepsilon^{\alpha-1} C_{M_0, \delta} \quad (3.59)$$

where we used again Assumption (1.7), Lemma 3.3 and (3.56) and the fact that the supports of the functions χ' and χ'' are $\{1/3 \leq r \leq 2/3\}$.

Let us now estimate the difference $y = \chi v^{\varepsilon, \delta} - w^\varepsilon$. This function satisfies the equation

$$\begin{aligned} i\partial_t \langle \nabla_\sigma \rangle y &= H_r \langle \nabla_\sigma \rangle y - \frac{1}{r^2} \Delta_\sigma \langle \nabla_\sigma \rangle y - \chi(r) \frac{r^2 - 1}{r^2} \Delta_\sigma \langle \nabla_\sigma \rangle v^{\varepsilon, \delta} \\ &\quad + \langle \nabla_\sigma \rangle \left(\mathbb{G}(|v^{\varepsilon, \delta}|^2) \chi v^{\varepsilon, \delta} - \mathbb{G}(|w^{\varepsilon, \delta}|^2) w^\varepsilon \right) + \langle \nabla_\sigma \rangle R^\varepsilon \end{aligned} \quad (3.60)$$

with $y(0) = u_0^{\varepsilon, \delta} - u_0^\varepsilon$. By (3.3), (3.50), (3.54) and (3.57), for $t \leq T_0^\varepsilon$ we have

$$\left\| \langle \nabla_\sigma \rangle \left(\mathbb{G}(|v^{\varepsilon, \delta}|^2) \chi v^{\varepsilon, \delta} - \mathbb{G}(|w^\varepsilon|^2) w^{\varepsilon, \delta} \right) (t) \right\|_{L^2} \leq C_{M_0} \|\langle \nabla_\sigma \rangle y(t)\|_{L^2} + \varepsilon^\alpha C_{M_0, \delta}.$$

Moreover, by interpolating and using that $\chi(r)$ vanishes near 0, one gets

$$\begin{aligned} \left\| \chi(r) \frac{r^2 - 1}{r^2} \Delta_\sigma \langle \nabla_\sigma \rangle v^{\varepsilon, \delta} \right\|_{L^2} &\leq C \left\| (r - 1) v^{\varepsilon, \delta} \right\|_{L^2}^{1/2} \left\| \langle \nabla_\sigma \rangle^6 v^{\varepsilon, \delta} \right\|_{L^2}^{1/2} \\ &\leq \varepsilon^{1/2} C_{M_0, \delta} \left\| V_c \left(\frac{r - 1}{\varepsilon} \right)^{1/\alpha} v^{\varepsilon, \delta} \right\|_{L^2}^{1/2} \\ &\leq \varepsilon^{1/2} C_{M_0, \delta} \left\| V_c \left(\frac{r - 1}{\varepsilon} \right)^{1/2} v^{\varepsilon, \delta} \right\|_{L^2}^{1/\alpha} \|v^{\varepsilon, \delta}\|_{L^2}^{1/2 - 1/\alpha} \\ &\leq \varepsilon^{1/2} C_{M_0, \delta} \left(\|(1 + \varepsilon^2 H_r)^{1/2} v^{\varepsilon, \delta}\|_{L^2} \right)^{1/\alpha} \|v^{\varepsilon, \delta}\|_{L^2}^{1/2 - 1/\alpha} \\ &\leq \varepsilon^{1/2} C_{M_0, \delta}. \end{aligned}$$

In this series of inequalities, we used (3.55), Assumption (1.7), a Hölder inequality (note that $\alpha > 2$) and, finally, the conservation laws (3.45) and (3.46) for the regularized function $v^{\varepsilon, \delta}$:

$$\|(1 + \varepsilon^2 H_r)^{1/2} v^{\varepsilon, \delta}\|_{L^2}^2 = \|(1 + \varepsilon^2 H_r)^{1/2} u_0^{\varepsilon, \delta}\|_{L^2}^2 \leq C.$$

Finally, the L^2 estimate for (3.60) yields

$$\begin{aligned} \|\langle \nabla_\sigma \rangle y(t)\|_{L^2} &\leq \|\langle \nabla_\sigma \rangle (u_0^{\varepsilon, \delta} - u_0^\varepsilon)\|_{L^2} + \varepsilon^{1/2} C_{M_0, \delta} + \varepsilon^{\alpha-1} C_{M_0, \delta} \\ &\quad + C_{M_0} \int_{-|t|}^{|t|} \|\langle \nabla_\sigma \rangle y(\tau)\|_{L^2} d\tau. \end{aligned}$$

We conclude by using the Gronwall lemma. We obtain

$$\limsup_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle y\|_{L^\infty([-T_0^\varepsilon, T_0^\varepsilon], L^2)} \leq C\delta.$$

Hence, using (3.53), (3.57) and letting δ tend to zero yields

$$\lim_{\varepsilon \rightarrow 0} \|\langle \nabla_\sigma \rangle (v^\varepsilon - w^\varepsilon)\|_{L^\infty([-T_0^\varepsilon, T_0^\varepsilon], L^2)} = 0.$$

In particular, from (3.49), we deduce that, for ε small enough, we have

$$\|\langle \nabla_\sigma \rangle w^\varepsilon\|_{L^\infty([-T_0^\varepsilon, T_0^\varepsilon], L^2)} \leq \frac{2M_0}{3},$$

which implies that $T_0^\varepsilon = T$. The proof of Proposition 3.5 is complete. \square

Proof of Theorem 1.5.

Theorem 1.5 is a direct consequence of Proposition 3.5, combined with Proposition 3.4. Indeed, Prop. 3.5, (i), states that the limit system (1.14) is globally well-posed. Let $T > 0$. Prop. 3.5, (ii), says that the intermediate system (3.29) is well-posed on $[-T, T]$ (for ε small enough), is uniformly bounded and converges to (1.14) as $\varepsilon \rightarrow 0$. Therefore, Prop. 3.4 (iii) can be applied thanks to this uniform bound: (1.4) is asymptotically close to (3.29), thus also converges to (1.14). \square

Appendix A. Proof of Lemma 3.3

In this appendix we identify the norm on the domains of iterates of the operator H_r used in Section 3. The lemma is a consequence of the following result.

Proposition A.1. *Let $W \in C^\infty(\mathbb{R}_+)$ be a real valued potential such that $W(x) \geq 1$ for every $x \in \mathbb{R}$ and satisfying*

$$\forall k \in \mathbb{N}, \exists C_k > 0 : \forall x \in \mathbb{R}_+, |W^{(k)}(x)| \leq C_k W(x) . \quad (\text{A.1})$$

Consider the following unbounded operator on $L^2(\mathbb{R}_+)$,

$$D(A) = \{u \in H^2(\mathbb{R}_+) : u(0) = 0, Wu \in L^2(\mathbb{R}_+)\} ; \quad Au := -u'' + Wu .$$

Then A is a positive selfadjoint operator and, for every integer $s \geq 1$,

$$D(A^{s/2}) = \left\{ u \in L^2(\mathbb{R}_+) : W^{\frac{s-k}{2}} u^{(k)} \in L^2(\mathbb{R}_+), 0 \leq k \leq s \right. \\ \left. \text{and } \left(-\frac{d^2}{dx^2} + W \right)^p u(0) = 0, 0 \leq p \leq \left\lfloor \frac{s-1}{2} \right\rfloor \right\}$$

and, on this space, the norm $\|A^{s/2}u\|_{L^2}$ is equivalent to

$$\sum_{k=0}^s \|W^{\frac{s-k}{2}} u^{(k)}\|_{L^2} ,$$

with constants only depending on the constants C_k in (A.1), for k in a finite set.

Proof. For simplicity, we denote by $\|f\|$ the norm of f in $L^2(\mathbb{R}_+)$, and by $(f|g)$ the corresponding inner product. We shall proceed in several steps.

Step 1. The case $s = 2$ and selfadjointness. In view of the definition of A , the symmetry identity

$$(Au_1|u_2) = (u_1|Au_2) , \quad u_1, u_2 \in D(A)$$

is merely an integration by parts. We now pass to a priori estimates. Firstly, integration by parts also implies

$$(Au|u) = \|u'\|^2 + \|\sqrt{W}u\|^2 , \quad u \in D(A) . \quad (\text{A.2})$$

In view of the assumption on W , this implies in particular

$$\|Au\| \geq \|u\| , \quad u \in D(A) . \quad (\text{A.3})$$

Next we derive a more precise estimate on $\|Au\|$ by computing

$$\|Au\|^2 = \|u''\|^2 + \|Wu\|^2 - 2\text{Re}(u''|Wu) .$$

Introducing $\chi \in C_0^\infty(\mathbb{R})$, $\chi \geq 0$, $\chi = 1$ near 0, we get

$$-(u''|Wu) = - \lim_{R \rightarrow \infty} \int_{\mathbb{R}_+} \chi\left(\frac{x}{R}\right) u''(x) W(x) \overline{u(x)} dx$$

and, after an integration by parts,

$$-(u''|Wu) = \lim_{R \rightarrow \infty} \int_{\mathbb{R}_+} \left(\frac{1}{R} \chi' \left(\frac{x}{R} \right) W(x) + \chi \left(\frac{x}{R} \right) W'(x) \right) u'(x) \overline{u(x)} + \chi \left(\frac{x}{R} \right) W(x) |u'(x)|^2 dx$$

Since $u' \in L^2$ and $Wu \in L^2$, the first term in the right hand side tends to 0 as R tends to infinity. Since moreover $W' = O(W)$, the second term has a limit. Consequently, the third term also has a limit. By Fatou's lemma, we conclude that $\sqrt{W}u' \in L^2$, and finally

$$\|Au\|^2 = \|u''\|^2 + \|Wu\|^2 + 2\|\sqrt{W}u'\|^2 + 2\text{Re}(u'|W'u). \quad (\text{A.4})$$

Because of the Cauchy–Schwarz inequality and of (A.1),

$$\begin{aligned} 2\text{Re}(u'|W'u) &\geq -2C_1\|u'\| \|Wu\| \geq -2C_1^2\|u'\|^2 - \frac{1}{2}\|Wu\|^2 \\ &\geq -\frac{1}{2}(\|u''\|^2 + \|Wu\|^2) - C'_1\|u\|^2, \end{aligned}$$

where C'_1 only depends on C_1 . Combining this inequality with (A.3), we infer

$$\|Au\| \geq c_1(\|u''\| + \|Wu\| + \|\sqrt{W}u'\|), \quad (\text{A.5})$$

where $c_1 > 0$ only depends on C_1 . We therefore have proved the statement for $s = 2$. Let us use this inequality for proving that A is selfadjoint. Recall that

$$D(A^*) = \{\psi \in L^2(\mathbb{R}_+) : \exists C > 0, \forall u \in D(A), |(Au|\psi)| \leq C\|u\|\}.$$

The symmetry of A already implies that $D(A) \subset D(A^*)$ and that $A^*u = Au$ for every $u \in D(A)$. Therefore we just have to prove that $D(A^*) \subset D(A)$. We claim that it is enough to prove that $\ker A^* = \{0\}$. Indeed, from estimate (A.5), it is easy to prove that the range of A is a closed subspace of L^2 . Since its orthogonal is $\ker A^*$, the cancellation of $\ker A^*$ would imply that A is onto. Consequently, for every $\psi \in D(A^*)$, there would exist $u \in D(A)$ such that $A^*\psi = Au$, namely $\psi - u \in \ker A^*$, hence $\psi = u \in D(A)$.

We now prove that $\ker A^* = \{0\}$. First of all, we observe that, for every $\psi \in D(A^*)$, in the distributional sense $\psi'' - W\psi \in L^2(\mathbb{R}_+)$, hence $\psi \in H^2((0, R))$ for every $R > 0$. Moreover, by integration by parts, for every $u \in D(A)$ supported into $[0, R]$ for some $R > 0$, we have

$$(Au|\psi) = u'(0)\overline{\psi(0)} + (u| - \psi'' + W\psi) = u'(0)\overline{\psi(0)} + O(\|u\|)$$

and testing the information $(Au|\psi) = O(\|u\|)$ on $u(x) = x\chi(nx)$ for large n imposes $\psi(0) = 0$. Assume moreover that $\psi \in \ker A^*$, namely that

$$\psi'' - W\psi = 0.$$

By the Sobolev embedding, we infer that $\psi \in C^\infty(\mathbb{R}_+)$. Set

$$v = |\psi|^2.$$

Then $v \in C^\infty(\mathbb{R}_+) \cap L^1(\mathbb{R}_+)$ and $v(0) = 0$. Plugging the differential equation satisfied by ψ , we get

$$v'' = 2Wv + 2|\psi'|^2 \geq 0.$$

In other word, v is a convex function. Since v is integrable at infinity, this implies that v is non increasing and tends to 0 at infinity. Since $v(0) = 0$, we conclude $v = 0$ and hence $\psi = 0$.

Step 2. The case $s = 1$. The domain of \sqrt{A} is characterized as the subspace of vectors $u \in L^2(\mathbb{R}_+)$ such that there exists a sequence (u_n) of $D(A)$ which tends to u in L^2 and which is a Cauchy sequence for the norm

$$N_1(v) := \sqrt{(Av|v)} \simeq \|v'\| + \|\sqrt{W}v\| .$$

This clearly implies that, if $u \in D(\sqrt{A})$, then $u \in H_0^1(\mathbb{R}_+)$ and $\sqrt{W}u \in L^2(\mathbb{R}_+)$. Conversely, if $u \in H_0^1(\mathbb{R}_+)$ and $\sqrt{W}u \in L^2(\mathbb{R}_+)$, a simple cutoff shows that u can be approximated in the N_1 norm by elements of $H_0^1(\mathbb{R}_+)$ with bounded supports. Then the claim reduces to the standard characterization of $H_0^1(\mathbb{R}_+)$ as the closure of $C_0^\infty((0, \infty))$ for the H^1 norm.

Step 3. The general case. We just prove the description of $D(A^{\frac{s}{2}})$, the corresponding equivalence of norms being proved in the same way, by keeping track the constants. We proceed by induction on s . Let $s \geq 3$ such that the claim is proved for every $s' \leq s - 1$. Then $u \in D(A^{\frac{s}{2}})$ if and only if $u \in D(A)$ and $Au \in D(A^{\frac{s-2}{2}})$. Using the induction hypothesis, the latter condition is equivalent to the following two conditions :

- $W^{\frac{s-2-k}{2}}(Au)^{(k)} \in L^2$ for every $k \leq s - 2$. Expanding $(Au)^{(k)}$ and using (A.1), we observe that this is equivalent to

$$W^{\frac{s-k}{2}}u^{(k)} - W^{\frac{s-2-k}{2}}u^{(k+2)} \in L^2 , \quad k \leq s - 2 ,$$

since the error terms are controlled by the fact that $u \in D(A^{\frac{s-1}{2}})$. In the special case $k = 0$, using again (A.1) and $u \in D(A^{\frac{s-1}{2}})$, we observe that this condition is equivalent to

$$-v'' + Wv \in L^2$$

for $v := W^{\frac{s-2}{2}}u$. Since $u \in D(A)$, $v \in H^2(0, R)$ for every $R > 0$ and $v(0) = 0$. Moreover, since $u \in D(A^{\frac{s-1}{2}})$ and by the induction hypothesis, $v \in H^1(\mathbb{R}_+)$. Hence, by computing $(Au|v)$ for $u \in D(A)$, we have $v \in D(A^*)$, which, by the first step, implies $Wv \in L^2$, or $W^{\frac{s}{2}}u \in L^2$. Combining with the other conditions for $k \leq s-2$, we eventually obtain $W^{\frac{s-k}{2}}u^{(k)} \in L^2$ for $k \leq s$.

- The boundary conditions

$$\left(-\frac{d^2}{dx^2} + W\right)^p (Au)(0) = 0 , \quad 0 \leq p \leq \left\lfloor \frac{s-3}{2} \right\rfloor .$$

Since $Au = -u'' + Wu$ and since $u(0) = 0$ from $u \in D(A)$, this leads to the claimed boundary conditions at rank s .

The proof is complete. □

Acknowledgements. The authors were supported by the Agence Nationale de la Recherche, ANR project EDP dispersives. F. Méhats was also supported by the ANR project QUATRIN and by the INRIA project IPSO.

References

- [1] S. Alinhac, P. Gérard, Pseudo-differential operators and the Nash-Moser theorem. Graduate Studies in Mathematics, **82**. American Mathematical Society, Providence, RI, 2007.
- [2] V. Banica, On the nonlinear Schrödinger dynamic on S^2 . J. Math. Pures Appl. **83** (2004), 77-98.
- [3] N. Ben Abdallah, F. Castella, F. Delebecque-Fendt, F. Méhats, The strongly confined Schrödinger-Poisson system for the transport of electrons in a nanowire. SIAM J. Appl. Math. **69** (2009), no. 4, 1162-1173.
- [4] N. Ben Abdallah, F. Castella, F. Méhats, Time averaging for the strongly confined nonlinear Schrödinger equation, using almost periodicity. J. Diff. Eq. **245** (2008), no. 1, 154-200..
- [5] N. Ben Abdallah, F. Méhats, O. Pinaud, Adiabatic approximation of the Schrödinger-Poisson system with a partial confinement. SIAM J. Math. Anal. **36** (2005), no. 3, 986-1013.
- [6] J.-M. Bony, J.-Y. Chemin, Espaces fonctionnels associés au calcul de Weyl-Hörmander. Bull. Soc. Math. France, Vol. 122, N. 1, 77-118 (1994).
- [7] J. Bourgain, Fourier transform restriction phenomena for certain lattice subsets and application to nonlinear equations I. Schrödinger equations. Geom. and Funct. Anal. **3** (1993), 107-156.
- [8] F. Brezzi, P. A. Markowich, The three dimensional Wigner -Poisson Problem : existence, uniqueness and approximation. Math. Methods Appl. Sci. **14** (1991), no. 1, 35-61.
- [9] N. Burq, P. Gérard, N. Tzvetkov, Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds. Amer. J. Math. **126** (2004), no. 3, 569-605.
- [10] N. Burq, P. Gérard, N. Tzvetkov, N. Bilinear eigenfunction estimates and the nonlinear Schrödinger equation on surfaces. Invent. math. **159** (2005), 187-223.
- [11] N. Burq, P. Gérard, N. Tzvetkov, N., Multilinear eigenfunction estimates and global existence for the three dimensional nonlinear Schrödinger equations. Ann. Sci. École Norm. Sup. (4) **38** (2005), no. 2, 255-301.
- [12] N. Burq, P. Gérard, N. Tzvetkov, N. The Cauchy problem for the nonlinear Schrödinger equation on compact manifolds. Phase space analysis of partial differential equations. Vol. I, 21-52, Publ. Cent. Ric. Mat. Ennio Giorgi, Scuola Norm. Sup., Pisa, 2004.
- [13] N. Burq, P. Gérard, N. Tzvetkov, An instability property of the nonlinear Schrödinger equation on S^d . Math. Res. Lett. **9** (2002), 323-335.
- [14] F. Castella, L^2 solutions to the Schrödinger-Poisson system: Existence, uniqueness, time behaviour, and smoothing effects. Math. Models Methods Appl. Sci., **7** (1997), 1051-1083.
- [15] R. C. T. da Costa, Quantum mechanics for a constraint particle. Phys. Rev. A **23** (1981), no. 4, 1982-1987.
- [16] F. Delebecque-Fendt, F. Méhats, An effective mass theorem for the bidimensional electron gas in a strong magnetic field. Comm. Math. Phys. **292** (2009), 829-870.
- [17] R. Froese, I. Herbst, Realizing Holonomic Constraints in Classical and Quantum Mechanics. Comm. Math. Phys. **220** (2001), no. 3, 489-535.
- [18] I. Gallagher, P. Gérard, Profile decomposition for the wave equation outside a convex obstacle. J. Math. Pures Appl. (9) **80** (2001), no. 1, 1-49.
- [19] P. Gérard, Nonlinear Schrödinger equations on compact manifolds. In European Congress of Mathematics (Stockholm, 2004), ed. by Ari Laptev, EMS Publishing House, Zürich, 2005, 121-139.
- [20] P. Gérard, V. Pierfelice, Nonlinear Schrödinger equation on four-dimensional compact manifolds. Bull. Soc. Math. France **138** (2010), no. 1, 119-151.
- [21] J. Ginibre, Le problème de Cauchy pour des EDP semi-linéaires périodiques en variable d'espace (d'après Bourgain). Séminaire Bourbaki, exp. 796, Astérisque **237** (1996), 163-187.
- [22] B. Helffer, F. Nier, Hypocoelliptic estimates and spectral theory for Fokker-Planck operators and Witten-Laplacians. Lecture Notes in Mathematics, Vol. 1862. Springer-Verlag, Berlin (2005).
- [23] L. Hörmander, The analysis of linear partial differential operators. III. Pseudodifferential operators. Chapter XVIII. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 274. Springer-Verlag, Berlin, 1985.
- [24] R. Illner, P. F. Zweifel, H. Lange, Global Existence, Uniqueness and Asymptotic Behaviour of Solutions of the Wigner-Poisson and Schrödinger-Poisson Systems. Math. Methods Appl. Sci. **17** (1994), no. 5, 349-376.

- [25] H. Poincaré, La méthode de Neumann et le problème de Dirichlet. Acta Math. **20** (1897), no. 1, 59–142.
- [26] W. Schlag, Harmonic analysis notes, available on <http://math.uchicago.edu/~schlag/>.
- [27] C. Sogge, Fourier integrals in classical analysis. Cambridge Tracts in Mathematics, 1993.
- [28] S. Teufel, Adiabatic Perturbation Theory in Quantum Dynamics. Lecture Notes in Mathematics 1821, Springer-Verlag, Berlin, Heidelberg, New York, 2003.

UNIVERSITÉ PARIS-SUD 11, LABORATOIRE DE MATHÉMATIQUES D'ORSAY, CNRS, UMR 8628, ET INSTITUT UNIVERSITAIRE DE FRANCE

E-mail address: `patrick.gerard@math.u-psud.fr`

IRMAR, UNIVERSITÉ RENNES 1, FRANCE

E-mail address: `florian.mehats@univ-rennes1.fr`

URL: <http://perso.univ-rennes1.fr/florian.mehats/>