

Amenability of covers and critical exponents

Dynamics of group actions

Cetraro, mai 2019

60 ans d'Yves Benoist


Work in collaboration with
Rémi Coulon, Rhiannon Dougall and Samuel Tapie

Critical exponents

A discrete group Γ acts on a hyperbolic space X (for ex. $X = \mathbb{H}^n$).

The **critical exponent** of this action is

$$\delta_\Gamma = \limsup_{R \rightarrow \infty} \frac{1}{R} \log \#\{\gamma \in \Gamma, d(o, \gamma o) \leq R\}.$$



Critical exponents

Coincides with

→ the **dimension** of the (radial) limit set $\Lambda_{rad}(\Gamma)$ inside ∂X

→ the **entropy** of the geodesic flow on $T^1(X/\Gamma)$

Of course, if $\Gamma' < \Gamma$, $\delta_{\Gamma'} \leq \delta_{\Gamma}$.

Question : When do we have equality $\delta_{\Gamma'} = \delta_{\Gamma}$?

Our result

Theorem : (Coulon-Dougall-Sch.-Tapie 2018) Let Γ be a discrete group acting on a proper hyperbolic space X , with *entropy gap at infinity* $\delta_\Gamma^\infty < \delta_\Gamma$. Let $\Gamma' < \Gamma$ be a subgroup. Then

$$\delta_{\Gamma'} = \delta_\Gamma \quad \text{iff} \quad \Gamma/\Gamma' \text{ amenable.}$$

Result already known in particular cases :

- Brooks (81,85) : convex-cocompact actions on \mathbb{H}^n , with $\delta_\Gamma > n/2$
- Grigorchuk, Cohen (80, 82) : action of a free group on its Cayley graph
- Stadlbauer: convex-cocompact (and some geom. finite) actions on \mathbb{H}^n
- Dougall-Sharp : convex-cocompact actions in variable neg. curvature
- Coulon-Dal'bo-Sambusetti : cocompact actions on $CAT(-1)$ -spaces
- Roblin (03) : amenability implies equality when $\Gamma' \triangleleft \Gamma$

Amenability

The group Γ' is **coamenable** in Γ if the regular representation


$$\rho : \Gamma \rightarrow \mathcal{U}(\ell^2(\Gamma/\Gamma'))$$

defined by $\rho(\gamma)(\varphi)(\cdot) = \varphi(\gamma \cdot)$ **almost admits invariant vectors**:
for all $\varepsilon > 0$ and $S \subset \Gamma$ finite, there exists $\varphi \in \ell^2(\Gamma/\Gamma')$ such that
for $\gamma \in S$, $\|\rho(\gamma)\varphi - \varphi\| < \varepsilon\|\varphi\|$.


Typical amenable group: \mathbb{Z}, \mathbb{Z}^d

Typical nonamenable group: \mathbb{F}^n

(Non)-Amenable covers



And an attempt to draw a \mathbb{F}_2 -cover of a compact hyperbolic surface



Entropy gap at infinity

The group Γ acts on X (for ex. $X = \mathbb{H}^n$)

Let $K \subset X$ be a compact set, $o \in K$. Define

$$\Gamma_K = \{\gamma \in \Gamma, [o, \gamma o] \cap \Gamma K \subset K \cup \gamma K\} \subset \Gamma.$$

The **entropy at infinity** is


$$\delta_\Gamma^\infty = \inf_{K \subset X} \delta_{\Gamma_K} \leq \delta_\Gamma.$$

The action of Γ on X admits a **entropy gap at infinity** when $\delta_\Gamma^\infty < \delta_\Gamma$.


We call these actions **strongly positively recurrent actions**.

Manifolds with(out) entropy gap at infinity

Typical examples **without** entropy gap : infinite covers.



Typical examples **with** entropy gap : compact or convex-cocompact manifolds, geometrically finite locally symmetric manifolds, Schottky products, Ancona surfaces.



Optimality of the assumptions

Result **false without hyperbolicity** :

Γ amenable group with exponential growth, $X = \text{Cay}(\Gamma)$, $\Gamma' = \{1\}$.
Then $\delta_\Gamma > 0$ whereas $\delta_{\Gamma'} = 0$, and $\Gamma/\Gamma' = \Gamma$ is amenable

On **higher rank symmetric spaces**, if Γ/Γ' is amenable, then
 $\delta_\Gamma = \delta_{\Gamma'}$ (Glorieux-Tapie).

Result **false without critical gap** :

Let $S = X/\Gamma$ be a negatively curved surface without critical gap.
For example, S is a \mathbb{Z} -cover of a compact hyperbolic surface. Build
a \mathbb{F}_2 -cover $S' = X/\Gamma'$ of S by cutting S along two disjoint
nonseparating closed curves. Then there is no critical gap :

$$\delta_\Gamma \geq \delta_{\hat{\Gamma}} \geq \delta_{\hat{\Gamma}}^\infty = \delta_{\Gamma'}^\infty = \delta_{\Gamma'}.$$

Recall the Patterson-Sullivan construction

The **Poincaré series**

$$P(s) = \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)}$$

has **critical exponent** δ_Γ . For $s > \delta_\Gamma$, build a measure on $X \cup \partial X$

$$\nu^s = \frac{1}{P(s)} \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} \mathcal{D}_{\gamma o}.$$

When $s \rightarrow \delta_\Gamma$, get ν on ∂X as any weak limit of ν^s . The measure ν on ∂X is quasi-invariant under Γ .

The unit tangent bundle satisfies $T^1X \simeq \partial X \times \partial X \setminus \text{Diag} \times \mathbb{R}$

Build a Γ -invariant product equivalent to $\nu \times \nu \times dt$

Get *Bowen-Margulis measure* m_{BM} on T^1X/Γ (ergodic, mixing...)

Strategy of the proof

Step 1: Twisted Poincaré series $A(s) = \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} \rho(\gamma)$. It has

a critical exponent $\delta_\rho \in [\delta_{\Gamma'}, \delta_\Gamma]$ such that for $s > \delta_\rho$, $A(s) \in \mathcal{B}(\ell^2(\Gamma/\Gamma'))$.

Step 2: Build a twisted Patterson-Sullivan measure a^ρ on ∂X with (nonzero) values in $\mathcal{B}(\ell^2(\Gamma/\Gamma'))$, by taking limits of

$$\frac{1}{\|A(s)\|} \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} \rho(\gamma) \mathcal{D}_{\gamma o}.$$

Step 3: When $\delta_\rho = \delta_\Gamma$ and $\delta_\Gamma^\infty < \delta_\Gamma$, get absolute continuity of a^ρ w.r.t. the classical Patterson-Sullivan measure ν .

Step 4: By an ergodicity argument, deduce that $a^\rho = \Psi \cdot \nu$ where $\Psi \in \mathcal{B}(\ell^2(\Gamma/\Gamma'))$ is a "multiplicative constant".

Step 5: By construction of a^ρ and ν , Ψ "takes values" in the set of almost invariant vectors.

Step 1: The twisted Poincaré series

Study $A(s) = \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} \rho(\gamma)$.

The Hilbert space $\mathcal{H} = \ell^2(\Gamma/\Gamma', \mathbb{R})$ has a **partial order** compatible with the norm: $\phi \geq 0$ if for all $y \in \Gamma/\Gamma'$, $\phi(y) \geq 0$.

Define the associated **positive cone** \mathcal{H}_+ . A bounded operator on \mathcal{H} is **positive** if it preserves \mathcal{H}_+ . All $\rho(\gamma)$ are positive.

The Poincaré series $A(s)$ is **bounded** if $\exists M > 0$, s.t. for all finite $S \subset \Gamma$, $\|\sum_{\gamma \in S} e^{-sd(o, \gamma o)} \rho(\gamma)\| \leq M$. The **critical exponent**

$$\delta_\rho = \inf\{s \in \mathbb{R}, A(s) \text{ is bounded}\}$$

is well defined. Easy to check that

$$\delta_{\Gamma'} \leq \delta_\rho \leq \delta_\Gamma.$$

The **assumption** $\delta_{\Gamma'} = \delta_\Gamma$ is only used to **guarantee that** $\delta_\rho = \delta_\Gamma$.

Step 2: The twisted Patterson-Sullivan measure

Let $\mathcal{H} = \ell^2(\Gamma/\Gamma')$. Use a nonprincipal **ultrafilter** $\omega : \mathcal{P}(\mathbb{N}) \rightarrow \{0, 1\}$.

Build a **larger Hilbert space** $\mathcal{H}_\omega = \lim_\omega \mathcal{H}$, and extend ρ to

$\rho_\omega : \Gamma \rightarrow \mathcal{U}(\mathcal{H}_\omega)$. We still have a **partial order** on \mathcal{H}_ω .

A sequence $\Phi = (\phi_n)$ of **almost invariant vectors** in $\mathcal{H}^{\mathbb{N}}$ becomes an **invariant vector** Φ under ρ_ω on \mathcal{H}_ω .

Choose $s_n \rightarrow \delta_\rho$. Define

$$a_n^\rho = \frac{1}{\|A(s_n)\|} \sum_{\gamma \in \Gamma} e^{-s_n d(o, \gamma o)} \rho(\gamma) \mathcal{D}_{\gamma o}.$$

For $f \in C(X \cup \partial X)$, $\int f da_n^\rho$ belongs to $\mathcal{B}(\mathcal{H})$, with norm uniformly bounded in n .


Define $a^\rho : C(X \cup \partial X) \rightarrow \mathcal{B}(\mathcal{H}_\omega)$ positive, linear, continuous by

$$a^\rho(f) := \lim_\omega \int f da_n^\rho \in \mathcal{B}(\mathcal{H}_\omega).$$

Nonzero measure on ∂X with values in $\mathcal{B}(\mathcal{H}_\omega)$

Step 3: Absolute continuity (I)

A **shadow** is $\mathcal{O}_o(B(y, r)) = \{\xi \in \partial X, (o\xi) \cap B(y, r) \neq \emptyset\}$.



The classical **Patterson-Sullivan measure** ν is a weak limit of $\frac{1}{P(s)} \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} \mathcal{D}_{\gamma o}$.

Sullivan's Shadow lemma says that $\nu(\mathcal{O}_o(B(\gamma o, r))) \asymp e^{-\delta_{\Gamma} d(o, \gamma o)}$.

A **half-Shadow lemma** for a^{ρ} : $\|a^{\rho}(\mathcal{O}_o(B(\gamma o, r)))\| \leq e^{-\delta_{\Gamma} d(o, \gamma o)}$.

Step 3: Absolute continuity (II)

Absolute continuity on shadows

$$\|a^\rho(\mathcal{O}_o(B(\gamma o, r)))\| \leq \nu(\mathcal{O}_o(B(\gamma o, r))).$$

The entropy gap allows to show that points lying in infinitely many shadows have full ν and a^ρ -measure.

Only but crucial place where we need the entropy gap.

By a Vitali type argument, we deduce that $0 \neq a^\rho \ll \nu$.

For all $\phi \in \mathcal{H}$, $a^\rho \cdot \phi \ll \nu$.

There exists a Radon-Nikodym derivative $D(\phi) \in L^\infty(X \cup \partial X, \mathcal{H})$, such that

$$\int f d(a^\rho \cdot \phi) = \int f D(\phi) d\nu.$$

Step 4: Ergodicity

* The map $\phi \in \mathcal{H} \rightarrow D(\phi) \in L^\infty((\partial X, \nu), \mathcal{H}_\omega)$ is linear and satisfies

$$\rho(\gamma) \circ D(\phi) \circ \gamma^{-1} = D(\phi).$$

* The map $(\xi, \eta) \in (\partial X)^2 \rightarrow \langle D(\phi)(\xi), D(\phi)(\eta) \rangle_{\mathcal{H}_\omega} \in \mathbb{R}$ is a Γ -invariant real-valued map.

→ The measure $\nu \times \nu$ on $\partial X \times \partial X$ is ergodic w.r.t. the Γ -action

Hint: $T^1X \simeq \partial X \times \partial X \times \mathbb{R}$. The PS measure ν on ∂X allows to build the Bowen-Margulis measure $m_{BM} \sim \nu \times \nu \times dt$ on T^1X/Γ . By Hopf argument, when X is a $CAT(-1)$ -space, it is an ergodic invariant measure for the geodesic flow. Also true for X Gromov-hyperbolic (Bader-Furman strategy).

→ We deduce $D(\phi)$ is $\nu \times \nu$ -a.s. constant.

Step 5 : Conclusion

We already know that (for any $\phi \in \mathcal{H}_\omega$, say with $\|\phi\| = 1$)

$$\rho(\gamma) \circ D(\phi) \circ \gamma^{-1} = D(\phi).$$

Moreover, as a map defined on $X \cup \partial X$, it is a.s. constant.

Therefore, for all $\gamma \in \Gamma$, we get the equality in \mathcal{H}_ω

$$\rho(\gamma).D(\phi) = D(\phi)$$

We got it !! $D(\phi)$ is our ρ -invariant vector in \mathcal{H}_ω .


More on the entropy gap

Show that ν and a^p are supported on $\Lambda_{rad}(\Gamma)$ (same proof).

$$\Lambda_{rad}(\Gamma) \supset \Lambda_{rad}^K(\Gamma) = \{\xi \in \Lambda(\Gamma), [o\xi) \text{ returns i.o. in } \Gamma.K\}$$

Define

$$U_K(T) = \{y \in X \cup \partial X, [oy) \text{ does not return in } K \text{ until time } T\}.$$



Entropy gap $\delta_\Gamma^\infty < \delta_\Gamma$ allows to show $\nu(U_K^T) \leq e^{(\delta_{\Gamma_K} - \delta_\Gamma)T}$

Deduce

$$\nu(\cap_{T>0} U_K^T) = 0 \quad \text{and} \quad \nu(\Gamma.(\cap_{T>0} U_K^T)) = 0$$

The "easy" direction

Kesten Criterion : any random walk on Γ/Γ' has spectral radius = 1.

Build a sequence of random walks w.r.t. uniform spherical measures on the spheres $S(e, n)$.

Barta's trick : estimate spectral radius on positive functions.

Estimate uniformly from above their spectral radius by $\exp(n(\delta_{\Gamma'} - \delta_{\Gamma}))$.

Roblin needed $\Gamma' \triangleleft \Gamma$. We remove this assumption, but use $\delta_{\Gamma'}^{\infty} < \delta_{\Gamma}$

Thank you!

And

Joyeux Anniversaire!