

Dimension of ergodic measures and currents on $\mathbb{CP}(2)$

Christophe Dupont and Axel Rogue

October 19, 2018

Abstract

Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. We estimate the local directional dimensions of closed positive currents S with respect to ergodic dilating measures ν . We infer several applications. The first one is an upper bound for the lower pointwise dimension of the equilibrium measure, towards a Binder-DeMarco's formula for this dimension. The second one shows that every current S containing a measure of entropy $h_\nu > \log d$ has a directional dimension > 2 , which answers a question of de Thélin-Vigny in a directional way. The last one estimates the dimensions of the Green current of Dujardin's semi-extremal endomorphisms.

Key words: dimension theory, Lyapunov exponents, normal forms.

MSC 2010: 37C45, 37F10, 32H50, 32U40.

1 Introduction

We study the dimension of ergodic measures for holomorphic endomorphisms of \mathbb{P}^2 and the dimension of currents containing such measures.

Let us recall dynamical properties of these mappings, see [18] for a detailed exposition. The topological entropy of an endomorphism of \mathbb{P}^2 of degree $d \geq 2$ is equal to $2 \log d$. Its Green current is defined by $T := \lim_n \frac{1}{d^n} f^{n*} \omega$, where ω is the Fubini-Study $(1,1)$ -form of \mathbb{P}^2 . The support of T is the subset of points $x \in \mathbb{P}^2$ such that $\{f^n, n \geq 1\}$ is not equicontinuous in any neighbourhood of x . The equilibrium measure of f is $\mu := T \wedge T$. This is a mixing invariant measure, it equidistributes the repulsive cycles of f , and its Lyapunov exponents are $\geq \frac{1}{2} \log d$. This is also the unique measure of maximal entropy $h_\mu = 2 \log d$.

The ergodic measures ν satisfying $\log d < h_\nu \leq 2 \log d$ are called *measures of large entropy*. Their support is contained in the support of μ [11, 16] and their exponents are larger than or equal to $\frac{1}{2}(h_\nu - \log d)$ [12, 21]. Examples are constructed in [21] by using coding techniques. The ergodic measures ν with positive exponents are called *dilating*.

1.1 Dimension of dilating measures

An open problem is to find a formula for the dimension of dilating measures, see [23, Question 2.17] for μ . Difficulties are due to the facts that f is not invertible and not

conformal. The dimension of a probability measure ν is defined by (see [26, 29]):

$$\dim_H(\nu) := \inf \left\{ \dim_H(A), A \text{ Borel set of } \mathbb{P}^2, \nu(A) = 1 \right\}.$$

The lower and upper pointwise dimensions of ν at x are

$$\underline{d}_\nu(x) := \liminf_{r \rightarrow 0} \frac{\log(\nu(B_x(r)))}{\log r}, \quad \overline{d}_\nu(x) := \limsup_{r \rightarrow 0} \frac{\log(\nu(B_x(r)))}{\log r}.$$

These functions are ν -almost everywhere constant when ν is ergodic. If $a \leq d_\nu \leq \overline{d}_\nu \leq b$, then $a \leq \dim_H(\nu) \leq b$. If ν is dilating and $\lambda_1 \geq \lambda_2$ denote its exponents, then

$$\frac{h_\nu}{\lambda_1} \leq \underline{d}_\nu \leq \overline{d}_\nu \leq \frac{h_\nu}{\lambda_2}. \quad (1.1)$$

Binder-DeMarco [5] conjectured the following formula for the measure μ

$$\dim_H(\mu) = \frac{\log d}{\lambda_1} + \frac{\log d}{\lambda_2}, \quad (1.2)$$

which generalizes the one-dimensional Mañé's formula [25]. Let us note that by (1.1), that formula is true when $\lambda_1 = \lambda_2$. In view of that Conjecture, the following upper bound was proved in [5] for polynomial mappings

$$\dim_H(\mu) \leq 4 - \frac{2(\lambda_1 + \lambda_2) - \log d^2}{\lambda_1}, \quad (1.3)$$

which was extended in [17] to meromorphic mappings. Moreover, for every dilating measure ν , the article [20] established:

$$\underline{d}_\nu \geq \frac{\log d}{\lambda_1} + \frac{h_\nu - \log d}{\lambda_2}, \quad (1.4)$$

which yields half of the Conjecture:

$$\dim_H(\mu) \geq \underline{d}_\mu \geq \frac{\log d}{\lambda_1} + \frac{\log d}{\lambda_2}. \quad (1.5)$$

At this stage, by combining (1.3) and (1.4), the Conjecture is true for every endomorphism of \mathbb{P}^2 satisfying $\frac{1}{2} \log d = \lambda_2 \leq \lambda_1$, for which we have

$$\dim_H(\mu) = \frac{\log d}{\lambda_1} + 2. \quad (1.6)$$

In this article we prove lower and upper estimates for the directional dimensions of currents containing dilating measures. Our techniques allow to show the following upper bound for the lower dimension of such measures, towards a Binder-DeMarco's formula for \underline{d}_μ . The exponents *do not resonate* if $\lambda_1 \neq k\lambda_2$ for every $k \geq 2$.

Theorem 1.1. *Let f be an endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Let ν be a dilating measure of exponents $\lambda_1 \geq \lambda_2$ whose support is contained in the support of μ . Then*

$$\underline{d}_\nu \leq \frac{\log d}{\lambda_1} + \frac{\log d}{\lambda_2} + 2 \left(1 - \frac{\lambda_2}{\lambda_1}\right).$$

Moreover, if the exponents do not resonate, then:

$$\underline{d}_\nu \leq \frac{\log d}{\lambda_1} + \frac{\log d}{\lambda_2} + 2 \min \left(1 - \frac{\lambda_2}{\lambda_1}; \frac{\lambda_1}{\lambda_2} - 1\right).$$

The next sections deal with the dimension of currents. We recall in Section 1.2 properties of the trace measure. The directional dimensions are more precise than the dimension of the trace measure, we present our estimates in Sections 1.3, 1.4 and 1.5.

1.2 Currents and the dimension of their trace measure

A positive closed current S of bidegree $(1,1)$ on a complex manifold is locally equal (in the sense of distributions) to $i\partial\bar{\partial}u$ where u is a plurisubharmonic function, see [14, Chapitre 3]. In particular S is locally equal to

$$S_{1,1} \frac{i}{2} dz \wedge d\bar{z} + S_{1,2} \frac{i}{2} dz \wedge d\bar{w} + S_{2,1} \frac{i}{2} dw \wedge d\bar{z} + S_{2,2} \frac{i}{2} dw \wedge d\bar{w},$$

where $S_{1,1}, S_{2,2}$ are positive measures and $S_{1,2} = \bar{S}_{2,1}$ is a complex measure dominated by $S_{1,1} + S_{2,2}$. The *trace measure* of S on \mathbb{P}^2 is the positive measure $S \wedge \omega$, where ω is the Fubini-Study form on \mathbb{P}^2 . This is a probability measure (up to a multiplicative positive constant) equivalent to $S_{1,1} + S_{2,2}$ in every local coordinates as before. It is known that $S \wedge \omega(B_x(r)) \leq c(x)r^2$ for every $x \in \mathbb{P}^2$, hence the pointwise dimension of $S \wedge \omega$ satisfies

$$\forall x \in \mathbb{P}^2, \underline{d}_S(x) := \liminf_{r \rightarrow 0} \frac{\log S \wedge \omega(B_x(r))}{\log r} \geq 2.$$

This bound can be improved for the Green current of endomorphisms f of \mathbb{P}^2 . Indeed, if we set

$$d_\infty := \lim_n \|Df^n\|_\infty^{1/n} \text{ and } \gamma_0 := \min\{1, \log d / \log d_\infty\},$$

then T has local γ -Hölder potentials u for every $\gamma < \gamma_0$, see [18, Proposition 1.18]. This implies that $T \wedge \omega(B_x(r)) \leq c_\gamma(x)r^{2+\gamma}$ for every $x \in \mathbb{P}^2$ and every $\gamma < \gamma_0$, see [27, Théorème 1.7.3]. Hence

$$\forall x \in \mathbb{P}^2, \underline{d}_T(x) \geq 2 + \gamma_0. \tag{1.7}$$

The geometric structure of the Green current is not well understood. A way to get informations is to study its dimension with respect to ergodic measures. For positive closed currents, there are dimensional constraints to contain measures of large entropy. For instance, any current of integration on an algebraic subset of \mathbb{P}^2 can not contain any such measure: this comes from Gromov's iterated graph argument and the relative variational principle, see [8] and [18, Section 1.7]. Further, de Thélin-Vigny [13] proved

that if a current S contains a measure of large entropy (or even dilating) with exponents $\lambda_1 \geq \lambda_2$, then for every $\epsilon > 0$ there exists x in the support of ν such that

$$\overline{d}_S(x) \geq 2 \frac{\lambda_2}{\lambda_1} + \frac{h_\nu - \log d}{\lambda_2} - \epsilon. \quad (1.8)$$

This inequality quantifies in an ergodic way (with exponents and entropy) the thickness of currents containing large entropy measures. Our results will in particular improve (1.7) and (1.8) in a directional way, when the exponents do not resonate.

1.3 Directional dimensions of currents

Let (Z, W) be holomorphic coordinates near x in \mathbb{P}^2 . The lower pointwise directional dimension of S with respect to Z is defined by

$$\underline{d}_{S,Z}(x) := \liminf_{r \rightarrow 0} \frac{\log \left[S \wedge \left(\frac{i}{2} dZ \wedge d\overline{Z} \right) (B_x(r)) \right]}{\log r}.$$

We use a similar definition for the upper pointwise dimension by taking a lim sup. Geometrically, the positive measure $S \wedge \left(\frac{i}{2} dZ \wedge d\overline{Z} \right)$ is the average of the slices of the current S transversally to the coordinate Z , see Section A.2. Moreover we have

$$\underline{d}_S(x) = \min \left\{ \underline{d}_{S,Z}(x), \underline{d}_{S,W}(x) \right\}. \quad (1.9)$$

In this article we shall work with coordinates (Z, W) coming from a normal form Theorem for the inverse branches $f_{\hat{x}}^{-n}$ along generic orbits \hat{x} . We shall call them Oseledec-Poincaré coordinates. When the exponents $\lambda_1 \geq \lambda_2$ do not resonate, this Theorem provides coordinates $(Z_{\hat{x}}^\epsilon, W_{\hat{x}}^\epsilon)$ near x such that

$$Z_{f_{\hat{x}}^{-n}(\hat{x})}^\epsilon \circ f_{\hat{x}}^{-n} \simeq e^{-n\lambda_1 \pm n\epsilon} \times Z_{\hat{x}}^\epsilon, \quad W_{f_{\hat{x}}^{-n}(\hat{x})}^\epsilon \circ f_{\hat{x}}^{-n} \simeq e^{-n\lambda_2 \pm n\epsilon} \times W_{\hat{x}}^\epsilon.$$

In the next statements, the functions $O(\epsilon)$ are of the form $\epsilon M(\epsilon)$, where M is a positive function which depends on the exponents and the entropy of ν , and on the degree of f . Concerning Theorem 1.2 and Corollaries 1.4 and 1.5, the functions $\underline{d}_{T,Z}$, $\underline{d}_{T,W}$, $\overline{d}_{T,Z}$, $\overline{d}_{T,W}$ are ν -almost everywhere constant.

1.4 Estimates for directional dimensions of currents

We begin with lower estimates for the directional dimensions of the Green current T with respect to dilating measures ν contained in μ (examples are measures of large entropy).

Theorem 1.2. *Let f be an endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Let ν be a dilating measure whose exponents $\lambda_1 > \lambda_2$ do not resonate and whose support is contained in the support of μ . Then for every $\epsilon > 0$ and for ν -almost every x , there exist holomorphic coordinates (Z, W) near x such that*

$$\underline{d}_{T,Z}(x) \leq 2 \frac{\lambda_1}{\lambda_2} + \frac{\log d}{\lambda_2} + O(\epsilon),$$

$$\underline{d}_{T,W}(x) \leq 2 + \frac{\log d}{\lambda_2} + O(\epsilon).$$

The proof relies on pluripotential theory (μ is the Monge-Ampère mass $T \wedge T$) and on a study of the jacobians of $T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon$ and $T \wedge \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\overline{W}_{\hat{x}}^\epsilon$ with respect to f . This study is given by Proposition 3.3, which will be also crucial to prove Theorem 1.1.

The next result concerns upper estimates for currents S containing dilating measures.

Theorem 1.3. *Let f be an endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Let S be a $(1, 1)$ -closed positive current on \mathbb{P}^2 . We assume that the support of S contains a measure of large entropy ν whose exponents satisfy $\lambda_1 > \lambda_2$ and do not resonate. Then for every $\epsilon > 0$, there exist $x \in \text{Supp } \nu$ and holomorphic coordinates (Z, W) near x such that:*

$$\begin{aligned} \overline{d}_{S,Z}(x) &\geq 2 + \frac{h_\nu - \log d}{\lambda_2} - O(\epsilon), \\ \overline{d}_{S,W}(x) &\geq 2 \frac{\lambda_2}{\lambda_1} + \frac{h_\nu - \log d}{\lambda_2} - O(\epsilon). \end{aligned}$$

In particular, S has a local directional dimension > 2 at some $x \in \text{Supp } \nu$.

Theorem 1.3 specifies in a directional way Theorem 2 of de Thélin-Vigny [13]. Indeed, the estimate concerning the Z -coordinate improves (1.8) by replacing $\frac{\lambda_2}{\lambda_1}$ by 1, which answers a question of [13] in a directional way. Our proof follows the strategy of [13] by taking into account the normal form Theorem. We obtain more precise lower bounds depending on \underline{d}_ν and we deduce Theorem 1.3 from the lower estimate (1.4) on \underline{d}_ν . We shall begin with a proof for the Green current T (Theorem 4.6), in this case the exposition is simpler because the directional dimensions of T are constant ν -almost everywhere. The case of general currents S needs a localization at some point in the support of S (Theorem 4.7).

Theorems 1.2 and 1.3 immediately imply the following result. Let us set

$$\gamma_1 := \frac{h_\mu - \log d}{\lambda_2} = \frac{\log d}{\lambda_2}.$$

Corollary 1.4. *Assume that the exponents of μ satisfy $\lambda_1 > \lambda_2$ and do not resonate. Then for every $\epsilon > 0$ and for μ -almost every x , there exist holomorphic coordinates (Z, W) near x such that*

$$2 + \gamma_1 - O(\epsilon) \leq \overline{d}_{T,Z}(x) \quad , \quad \underline{d}_{T,W}(x) \leq 2 + \gamma_1 + O(\epsilon).$$

This shows that, modulo $O(\epsilon)$, either the directional dimensions $\overline{d}_{T,Z}(x)$ and $\underline{d}_{T,W}(x)$ are distinct and separated by $2 + \gamma_1$, or they are equal to $2 + \gamma_1$. Let us observe that the first estimate improves (1.7), since $\lambda_1 \leq \log d_\infty$ and $\lambda_1 > \lambda_2$ imply $\gamma_1 > \gamma_0$.

We note that a bound > 2 for the dimension of the trace measure of the Green currents T^\pm is proved in [10] for invertible and meromorphic mappings of Kähler surfaces, the proof relies on the laminar properties of T^\pm and coding techniques.

1.5 Semi extremal endomorphisms

An endomorphism f is *extremal* if the exponents of μ satisfy $\lambda_1 = \lambda_2 = \frac{1}{2} \log d$. The articles [1, 4, 17] characterize these endomorphisms by several equivalent properties which are $\dim_H(\mu) = 4$, $\mu \ll \text{Leb}_{\mathbb{P}^2}$, T is a positive smooth form on some open subset of \mathbb{P}^2 , and f is a Lattès map. The articles [3, 15] show other characterizations.

An endomorphism f is *semi-extremal* if the exponents of μ satisfy $\lambda_1 > \lambda_2 = \frac{1}{2} \log d$. Formula (1.6) of Section 1.1 implies that these endomorphisms satisfy Conjecture (1.2):

$$\dim_H(\mu) = \frac{\log d}{\lambda_1} + 2.$$

Dujardin [19] proved that if $\mu \ll T \wedge \omega$ then f is semi-extremal. Examples satisfying this condition are suspensions of one-dimensional Lattès maps, and more generally endomorphisms with an invariant pencil of lines on which f induces a one dimensional Lattès map [22]. One may ask if there exist other examples. In view of a possible characterization, the next result provides necessary conditions about the dimension of those endomorphisms, it follows from Theorem 1.3 and (1.6).

Corollary 1.5. *Assume that $\mu \ll T \wedge \omega$, that $\overline{d_\mu} = \underline{d_\mu}$ and that the exponents $\lambda_1 > \lambda_2$ of μ do not resonate. For every $\epsilon > 0$ and for μ -almost every $x \in \mathbb{P}^2$, there exist holomorphic coordinates (Z, W) near x such that*

$$4 - O(\epsilon) \leq \overline{d_{T,Z}}(x) \quad \text{and} \quad 2 + \frac{\log d}{\lambda_1} - O(\epsilon) \leq \overline{d_{T,W}}(x) \leq 2 + \frac{\log d}{\lambda_1}.$$

The first estimate provides a maximal dimension for T , which could be explained by the presence of a one dimensional Lattès map inside the dynamics of f . However, it seems difficult to produce such a Lattès map, as well as an invariant pencil of lines. Note that Corollary 1.5 provides a situation where the dimensions $\overline{d_{T,Z}}(x)$ and $\overline{d_{T,W}}(x)$ are not equal. To outline the proof of Corollary 1.5, we shall see that (1.6) implies $\min \left\{ \overline{d_{T,Z}}(x), \overline{d_{T,W}}(x) \right\} \leq 2 + \frac{\log d}{\lambda_1}$, then we use Theorem 1.3 to verify that the minimum concerns the W -coordinate.

1.6 Organization of the article

Section 2 is devoted to normal forms and to the geometry of inverse branches. Theorem 1.2 is proved in Section 3. We show Theorem 1.3 and Corollary 1.5 in Section 4, and we show Theorem 1.1 in Section 5. Technical results are put together in an Appendix.

Acknowledgements: This article is part of the PhD thesis of the second author. We warmly thank Eric Bedford and Johan Taffin for their numerous comments which allowed to improve the first version of this work. We also thank the referee for his careful reading and for numerous remarks. We got the supports of Lambda (ANR-13-BS01-0002) and Centre Henri Lebesgue (ANR-11-LABX-0020-01).

2 Normal forms and Oseledec-Poincaré coordinates

2.1 Natural extension and normal forms

Let f be an endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Let \mathcal{C}_f be its critical set, this is an algebraic subset of \mathbb{P}^2 . If ν is a dilating measure, then $x \mapsto \log |Jac f(x)| \in L^1(\nu)$, which implies $\nu(\mathcal{C}_f) = 0$. Let X be the f -invariant Borel set $\text{Supp}(\nu) \setminus \cup_{n \in \mathbb{Z}} f^n(\mathcal{C}_f)$ and let

$$\hat{X} := \left\{ \hat{x} = (x_n)_{n \in \mathbb{Z}} \in X^{\mathbb{Z}}, \quad x_{n+1} = f(x_n) \right\}.$$

Let \hat{f} be the left shift on \hat{X} and $\pi_0(\hat{x}) := x_0$. There exists a unique \hat{f} -invariant measure $\hat{\nu}$ on \hat{X} such that $(\pi_0)_* \hat{\nu} = \nu$. We set $\hat{x}_n := \hat{f}^n(\hat{x})$ for every $n \in \mathbb{Z}$. A function $\alpha : \hat{X} \rightarrow]0, +\infty]$ is ϵ -tempered if $\alpha(\hat{f}^{\pm 1}(\hat{x})) \geq e^{-\epsilon} \alpha(\hat{x})$. For every $\hat{x} \in X$ we denote by $f_{\hat{x}}^{-n}$ the inverse branch of f^n defined in a neighbourhood of x_0 with values in a neighbourhood of x_{-n} . The articles [2] and [24] provide normal forms for these mappings. Let $d(\cdot, \cdot)$ be the Fubini-Study distance on \mathbb{P}^2 .

Theorem 2.1. ([2, Prop. 4.3]) *Let ν be a dilating measure with exponents $\lambda_1 > \lambda_2$ and let $\epsilon > 0$. There exists an \hat{f} -invariant Borel set $\hat{F} \subset \hat{X}$ such that $\hat{\nu}(\hat{F}) = 1$ and satisfying the following properties. There exist ϵ -tempered functions $\eta_\epsilon, \rho_\epsilon : \hat{F} \rightarrow]0, 1]$ and $\beta_\epsilon, L_\epsilon, M_\epsilon : \hat{F} \rightarrow [1, +\infty[$ and for every $\hat{x} \in \hat{F}$, there exists a holomorphic mapping*

$$\xi_{\hat{x}}^\epsilon : B_{x_0}(\eta_\epsilon(\hat{x})) \rightarrow \mathbb{D}^2(\rho_\epsilon(\hat{x}))$$

such that the following diagram commutes for every $n \geq n_\epsilon(\hat{x})$:

$$\begin{array}{ccc} B_{x_{-n}}(\eta_\epsilon(\hat{x}_{-n})) & \xleftarrow{f_{\hat{x}}^{-n}} & B_{x_0}(\eta_\epsilon(\hat{x})) \\ \downarrow \xi_{\hat{x}_{-n}}^\epsilon & & \xi_{\hat{x}}^\epsilon \downarrow \\ \mathbb{D}^2(\rho_\epsilon(\hat{x}_{-n})) & \xleftarrow{R_{n, \hat{x}}} & \mathbb{D}^2(\rho_\epsilon(\hat{x})) \end{array}$$

and such that

1. $\forall (p, q) \in B_{x_0}(\eta_\epsilon(\hat{x})), \quad \frac{1}{2}d(p, q) \leq |\xi_{\hat{x}}^\epsilon(p) - \xi_{\hat{x}}^\epsilon(q)| \leq \beta_\epsilon(\hat{x})d(p, q)$.
2. $\text{Lip}(f_{\hat{x}}^{-n}) \leq L_\epsilon(\hat{x})e^{-n\lambda_2+n\epsilon}$ on $B_{x_0}(\eta_\epsilon(\hat{x}))$.
3. if $\lambda_1 \notin \{k\lambda_2, k \geq 2\}$, $R_{n, \hat{x}}(z, w) = (\alpha_{n, \hat{x}}z, \beta_{n, \hat{x}}w)$,
if $\lambda_1 = k\lambda_2$ where $k \geq 2$, $R_{n, \hat{x}}(z, w) = (\alpha_{n, \hat{x}}z, \beta_{n, \hat{x}}w) + (\gamma_{n, \hat{x}}w^k, 0)$, with
 - (a) $e^{-n\lambda_1-n\epsilon} \leq |\alpha_{n, \hat{x}}| \leq e^{-n\lambda_1+n\epsilon}$ and $|\gamma_{n, \hat{x}}| \leq M_\epsilon(\hat{x})e^{-n\lambda_1+n\epsilon}$,
 - (b) $e^{-n\lambda_2-n\epsilon} \leq |\beta_{n, \hat{x}}| \leq e^{-n\lambda_2+n\epsilon}$.

Remark 2.2. *The diagram commutes for every $n \in \llbracket 1, \dots, n_\epsilon(\hat{x}) \rrbracket$ for the germs of the mappings, see [2]. The integer $n_\epsilon(\hat{x})$ is actually the smallest integer such that $L_\epsilon(\hat{x})e^{-n\lambda_2+n\epsilon} \leq e^{-n\epsilon}$, so that $L_\epsilon(\hat{x})e^{-n\lambda_2+n\epsilon}\eta_\epsilon(\hat{x}) \leq e^{-n\epsilon}\eta_\epsilon(\hat{x}) \leq \eta_\epsilon(\hat{x}_{-n})$. Item 2 thus ensures that $f_{\hat{x}}^{-n}(B_{x_0}(\eta_\epsilon(\hat{x}))) \subset B_{x_{-n}}(\eta_\epsilon(\hat{x}_{-n}))$ as indicated in the diagram.*

We shall need the following Lemma. Let $n_1(L)$ be the smallest integer n satisfying $L/4 \leq e^{n\epsilon}$. The first item uses the upper bound for $\text{Lip}(f_{\hat{x}}^{-n})$ provided by Theorem 2.1. The second item comes from [17, Proposition 3.1].

Lemma 2.3. *Let $\hat{x} \in \hat{F}$ such that $\eta_\epsilon(\hat{x}) \geq \eta$ and $L_\epsilon(\hat{x}) \leq L$. If $n \geq n_1(L)$ and $r \leq \eta$,*

1. $f_{\hat{x}_n}^{-n}(B_{x_n}(r/4)) \subset B_{x_0}(re^{-n\lambda_2+3n\epsilon})$ and $f_{\hat{x}}^{-n}(B_{x_0}(r/4)) \subset B_{x_{-n}}(re^{-n\lambda_2+3n\epsilon})$.
2. $B_{x_0}(re^{-n\lambda_1-4n\epsilon}) \subset f_{\hat{x}_n}^{-n}(B_{x_n}(\frac{r}{4}e^{-2n\epsilon}))$.

2.2 Oseledec-Poincaré coordinates

Let ν be a dilating measure with exponents $\lambda_1 > \lambda_2$. Let $\epsilon > 0$ and let us apply Theorem 2.1. For every $\hat{x} \in \hat{F}$ we denote by $(Z_{\hat{x}}^\epsilon, W_{\hat{x}}^\epsilon)$ the coordinates of $\xi_{\hat{x}}^\epsilon$. If the exponents do not resonate then the commutative diagram given by Theorem 2.1 implies:

$$Z_{\hat{x}_{-n}}^\epsilon \circ f_{\hat{x}}^{-n} = \alpha_{n,\hat{x}} \times Z_{\hat{x}}^\epsilon, \quad W_{\hat{x}_{-n}}^\epsilon \circ f_{\hat{x}}^{-n} = \beta_{n,\hat{x}} \times W_{\hat{x}}^\epsilon. \quad (2.1)$$

Hence, $f_{\hat{x}}^{-n}$ multiplies the first coordinate by $e^{-n\lambda_1 \pm n\epsilon}$ and the second coordinate by $e^{-n\lambda_2 \pm n\epsilon}$. The second property in (2.1) also holds in the resonant case $\lambda_1 \in \{k\lambda_2, k \geq 2\}$. We shall name the collection of local holomorphic coordinates

$$(Z, W)_\epsilon := (Z_{\hat{x}}^\epsilon, W_{\hat{x}}^\epsilon)_{\hat{x} \in \hat{F}}$$

Oseledec-Poincaré coordinates for (f, ν) . If S is $(1, 1)$ positive closed current we define

$$\underline{d}_{S,Z}(\hat{x}) := \liminf_{r \rightarrow 0} \frac{\log \left[S \wedge \left(\frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z_{\hat{x}}^\epsilon} \right) (B_x(r)) \right]}{\log r},$$

with similar definitions for $\overline{d}_{S,W}(\hat{x})$, $\overline{d}_{S,Z}(\hat{x})$, $\overline{d}_{S,W}(\hat{x})$. Using (2.1) and the fact that the Green current T is f -invariant, we obtain the following Proposition.

Proposition 2.4. *Let ν be a dilating measure of exponents $\lambda_1 > \lambda_2$. Let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . Then there exists a \hat{f} -invariant Borel set $\hat{\Lambda}_T \subset \hat{F}$ of $\hat{\nu}$ -measure 1 such that*

1. $\hat{x} \mapsto \overline{d}_{T,W}(\hat{x})$ and $\hat{x} \mapsto \underline{d}_{T,W}(\hat{x})$ are \hat{f} -invariant on $\hat{\Lambda}_T$.
2. $\hat{x} \mapsto \overline{d}_{T,Z}(\hat{x})$ and $\hat{x} \mapsto \underline{d}_{T,Z}(\hat{x})$ are \hat{f} -invariant on $\hat{\Lambda}_T$ if $\lambda_1 \notin \{k\lambda_2, k \geq 2\}$.

In particular, if the exponents do not resonate, these functions are constant $\hat{\nu}$ -almost everywhere. We shall denote them by

$$\overline{d}_{T,Z}(\nu), \underline{d}_{T,Z}(\nu), \overline{d}_{T,W}(\nu), \underline{d}_{T,W}(\nu).$$

Proof. We prove the invariance of $\overline{d_{T,W}}(\hat{x})$, the same arguments hold for the other functions. For every $z \in \mathbb{P}^2 \setminus \mathcal{C}_f$ we denote

$$a(z) := \frac{1}{2} \|(D_z f)^{-1}\|^{-1}, \quad \gamma(z) := \min\{a(z) \|f\|_{\mathcal{C}^2, \mathbb{P}^2}^{-1}, 1\}.$$

Then [7, Lemme 2] asserts that f is injective on $B_z(\gamma(z))$ and

$$\forall r \in [0, \gamma(z)] , \quad B_{f(z)}(a(z)r) \subset f(B_z(r)).$$

Let $\hat{x} \in \hat{F}$. Since $x_n \notin \mathcal{C}_f$ for every $n \in \mathbb{Z}$, we obtain for every $r \leq \gamma(x_0)$:

$$T \wedge \left(\frac{i}{2} dW_{\hat{f}(\hat{x})}^\epsilon \wedge d\overline{W_{\hat{f}(\hat{x})}^\epsilon} \right) [B_{f(x_0)}(a(x_0)r)] \leq T \wedge \left(\frac{i}{2} dW_{\hat{f}(\hat{x})}^\epsilon \wedge d\overline{W_{\hat{f}(\hat{x})}^\epsilon} \right) [f(B_{x_0}(r))]. \quad (2.2)$$

Since f is injective on $B_{x_0}(r)$, we can change the variables to get:

$$T \wedge \left(\frac{i}{2} dW_{\hat{f}(\hat{x})}^\epsilon \wedge d\overline{W_{\hat{f}(\hat{x})}^\epsilon} \right) [f(B_{x_0}(r))] = \int_{B_{x_0}(r)} f^* T \wedge f^* \left(\frac{i}{2} dW_{\hat{f}(\hat{x})}^\epsilon \wedge d\overline{W_{\hat{f}(\hat{x})}^\epsilon} \right). \quad (2.3)$$

Now let us recall that

$$f^* T = dT \quad \text{and} \quad f^* \left(\frac{i}{2} dW_{\hat{f}(\hat{x})}^\epsilon \wedge d\overline{W_{\hat{f}(\hat{x})}^\epsilon} \right) = |c(\hat{x})|^2 \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\overline{W_{\hat{x}}^\epsilon}, \quad (2.4)$$

where the second equality comes from (2.1) by setting $c(\hat{x})^{-1} := \beta_{1, \hat{f}(\hat{x})}$, it is valid near x_0 according to Remark 2.2. By combining (2.2), (2.3) and (2.4) we deduce:

$$T \wedge \left(\frac{i}{2} dW_{\hat{f}(\hat{x})}^\epsilon \wedge d\overline{W_{\hat{f}(\hat{x})}^\epsilon} \right) [B_{f(x_0)}(a(x_0)r)] \leq d|c(\hat{x})|^2 T \wedge \left(\frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\overline{W_{\hat{x}}^\epsilon} \right) [B_{x_0}(r)]$$

for every r small enough. Taking the logarithm and dividing by $\log(a(x_0)r) < 0$, we get $\overline{d_{T,W}}(\hat{f}(\hat{x})) \geq \overline{d_{T,W}}(\hat{x})$ by taking limits. Since $\hat{\nu}$ is ergodic, the function $\overline{d_{T,W}}(\hat{x})$ is constant on a Borel set $\hat{\Lambda}_T$ of $\hat{\nu}$ -measure 1 (see [28, Chapter 1.5]). One can replace it by $\bigcap_{n \in \mathbb{Z}} \hat{f}^n(\hat{\Lambda}_T)$ to obtain an invariant set. \square

Similar arguments yield the following Proposition for the trace measure.

Proposition 2.5. *If ν is an ergodic measure, the functions $x \mapsto \underline{d}_T(x)$ and $x \mapsto \overline{d}_T(x)$ are invariant, hence ν -almost everywhere constant. We denote them by $\underline{d}_T(\nu)$ and $\overline{d}_T(\nu)$.*

Proof. The arguments follow the proof of Proposition 2.4. In this case we study the measure $T \wedge \omega$, and we replace the second equality in (2.4) by $f^* \omega \leq \rho(x_0) \omega$ on $B_{x_0}(\gamma(x_0))$, where $\rho(x_0) > 0$ is a large enough positive constant. \square

2.3 Pullback of the Fubini-Study form ω

Let ν be a dilating measure of exponents $\lambda_1 > \lambda_2$. Let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . Let $n_2(\beta)$ be the smallest integer n such that $e^{-n\epsilon} \leq \beta^{-1}$.

Proposition 2.6. *Let $\hat{x} \in \hat{F}$ such that $\eta_\epsilon(\hat{x}) \geq \eta$ and $\beta_\epsilon(\hat{x}) \leq \beta$. If $n \geq \max\{n_2(\beta), n_\epsilon(\hat{x}_n)\}$ and if $r \leq \eta$, then we have on $f_{\hat{x}_n}^{-n}(B_{x_n}(re^{-n\epsilon}))$:*

1. $(f^n)^*\omega \geq e^{-6n\epsilon+2n\lambda_1} \left(\frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon \right)$ if the exponents do not resonate.
2. $(f^n)^*\omega \geq e^{-6n\epsilon+2n\lambda_2} \left(\frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\overline{W}_{\hat{x}}^\epsilon \right)$.

The proof follows from the next two lemmas. First observe that Theorem 2.1 gives

$$(f^n)^*\omega = (\xi_{\hat{x}}^\epsilon)^*((R_{n, \hat{x}_n})^{-1})^*((\xi_{\hat{x}_n})^{-1})^*\omega$$

on $f_{\hat{x}_n}^{-n}(B_{x_n}(re^{-n\epsilon}))$. Let $\omega_0 := \frac{i}{2} dz \wedge d\bar{z} + \frac{i}{2} dw \wedge d\bar{w}$ be the standard $(1, 1)$ -form on \mathbb{D}^2 .

Lemma 2.7. *Let $\hat{x} \in \hat{F}$ such that $\eta_\epsilon(\hat{x}) \geq \eta$ and $\beta_\epsilon(\hat{x}) \leq \beta$. For every $n \geq n_2(\beta)$ and $r \leq \eta$, we have on $\xi_{\hat{x}_n}^\epsilon(B_{x_n}(re^{-n\epsilon}))$:*

$$e^{-4n\epsilon}\omega_0 \leq ((\xi_{\hat{x}_n}^\epsilon)^{-1})^*\omega \leq 4\omega_0.$$

Proof. For every $p = (z, w)$ and $p' = (z', w')$ in $\xi_{\hat{x}_n}^\epsilon(B_{x_n}(re^{-n\epsilon}))$, we have

$$e^{-n\epsilon}\beta^{-1}|p - p'| \leq d((\xi_{\hat{x}_n}^\epsilon)^{-1}(p), (\xi_{\hat{x}_n}^\epsilon)^{-1}(p')) \leq 2|p - p'|.$$

This implies for every $n \geq n_2(\beta)$ and $(z, w) \in \xi_{\hat{x}_n}^\epsilon(B_{x_n}(re^{-n\epsilon}))$:

$$\forall u \in \mathbb{C}^2, \quad e^{-2n\epsilon}|u| \leq \left| D_{(z, w)}(\xi_{\hat{x}_n}^\epsilon)^{-1}(u) \right| \leq 2|u|.$$

This provides the desired estimates. □

Lemma 2.8. *Let $\hat{x} \in \hat{F}$. If $n \geq n_\epsilon(\hat{x}_n)$, then*

1. $((R_{n, \hat{x}_n})^{-1})^*\omega_0 \geq e^{2(n\lambda_1 - n\epsilon)} \frac{i}{2} dz \wedge d\bar{z}$ if the exponents do not resonate.
2. $((R_{n, \hat{x}_n})^{-1})^*\omega_0 \geq e^{2(n\lambda_2 - n\epsilon)} \frac{i}{2} dw \wedge d\bar{w}$.

Proof. We use the fact that the linear part of R_{n, \hat{x}_n} is diagonal with coefficients $e^{-n\epsilon - n\lambda_1} \leq |\alpha_{n, \hat{x}_n}| \leq e^{n\epsilon - n\lambda_1}$ and $e^{-n\epsilon - n\lambda_2} \leq |\beta_{n, \hat{x}_n}| \leq e^{n\epsilon - n\lambda_2}$ (see Theorem 2.1) and the fact that the $(1, 1)$ -forms $\frac{i}{2} dz \wedge d\bar{z}$ and $\frac{i}{2} dw \wedge d\bar{w}$ are positive. □

To end the proof of Proposition 2.6, we observe that for every $\hat{x} \in \hat{F}$:

$$(\xi_{\hat{x}}^\epsilon)^*\left(\frac{i}{2} dz \wedge d\bar{z}\right) = \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon, \quad (\xi_{\hat{x}}^\epsilon)^*\left(\frac{i}{2} dw \wedge d\bar{w}\right) = \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\overline{W}_{\hat{x}}^\epsilon$$

which follows from the definitions of $Z_{\hat{x}}^\epsilon$ and $W_{\hat{x}}^\epsilon$.

3 Upper bounds for the directional dimensions of T

Let ν be a dilating measure such that $\text{Supp}(\nu) \subset \text{Supp}(\mu)$ and whose exponents $\lambda_1 > \lambda_2$ do not resonate. Let $\epsilon > 0$ and let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . In this section we establish Theorem 1.2 stated in the introduction, that is

$$\underline{d_{T,Z}}(\nu) \leq 2 \frac{\lambda_1}{\lambda_2} + \frac{\log d}{\lambda_2} + O(\epsilon) \quad \text{and} \quad \underline{d_{T,W}}(\nu) \leq 2 + \frac{\log d}{\lambda_2} + O(\epsilon).$$

The proof relies on the Monge-Ampère equation $\mu = T \wedge T$ and on a study of the jacobians of $T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\bar{Z}_{\hat{x}}^\epsilon$ and $T \wedge \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\bar{W}_{\hat{x}}^\epsilon$ with respect to f .

3.1 Dimensions of the Green current on the equilibrium measure

The following Proposition is proved in Section 3.3.

Proposition 3.1. *Let f be an endomorphism of \mathbb{P}^2 of degree $d \geq 2$ and let T be its Green current. Let $x \in \text{Supp} \mu$ and let Z be a local holomorphic coordinate (submersion) in a neighbourhood V of x . Then $T \wedge (\frac{i}{2} dZ \wedge d\bar{Z})$ is not the zero measure on V .*

This implies:

Proposition 3.2. *Let ν be a dilating measure with exponents $\lambda_1 > \lambda_2$ whose support is contained in the support of μ . Let $\epsilon > 0$ and let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . Then, for every $0 < r < \eta_\epsilon(\hat{x})$,*

$$\left[T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\bar{Z}_{\hat{x}}^\epsilon \right] (B_x(r)) > 0 \quad \text{and} \quad \left[T \wedge \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\bar{W}_{\hat{x}}^\epsilon \right] (B_x(r)) > 0.$$

In particular, for every $\delta > 0$, there exist $m_0 \geq 1$, $L_0 \geq 1$ and $q_0 \geq 1$ such that

$$\hat{\Omega}_\epsilon := \left\{ \eta_\epsilon(\hat{x}) \geq \frac{1}{4m_0}, \quad L(\hat{x}) \leq L_0, \quad \left[T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\bar{Z}_{\hat{x}}^\epsilon \right] (B_x(\frac{1}{4m_0})) \geq \frac{1}{q_0}, \quad \left[T \wedge \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\bar{W}_{\hat{x}}^\epsilon \right] (B_x(\frac{1}{4m_0})) \geq \frac{1}{q_0} \right\}$$

satisfies $\hat{\nu}(\hat{\Omega}_\epsilon) \geq 1 - \delta$.

Proof. The first part immediately follows from Proposition 3.1. To prove the second part, let $m_0 \geq 1$ and $L_0 \geq 1$ be such that $\hat{\nu} \left\{ \eta_\epsilon \geq \frac{1}{4m_0} \right\} \cap \{L \leq L_0\} \geq 1 - \delta/2$. Then we choose q_0 large enough so that $\hat{\nu}(\hat{\Omega}_\epsilon) \geq 1 - \delta$. \square

We define for every $n \geq 1$:

$$\hat{\Omega}_\epsilon^n := \hat{\Omega}_\epsilon \cap \hat{f}^{-n}(\hat{\Omega}_\epsilon).$$

Since $\hat{\nu}$ is invariant, we have:

$$\hat{\nu}(\hat{\Omega}_\epsilon^n) \geq 1 - 2\delta. \tag{3.1}$$

The following Proposition is crucial to prove Theorems 1.1 and 1.2. L_0 is defined in Proposition 3.2 and $n_1(L_0) \geq 1$ is defined before Lemma 2.3.

Proposition 3.3. *Let ν be a dilating measure with exponents $\lambda_1 > \lambda_2$ and such that $\text{Supp}(\nu) \subset \text{Supp}(\mu)$. For every $n \geq n_1(L_0)$ and $\hat{x} \in \hat{\Omega}_\epsilon^n$, we have:*

$$\begin{aligned} \left[T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon \right] \left(B_x \left(\frac{1}{m_0} e^{-n\lambda_2 + 3n\epsilon} \right) \right) &\geq \frac{1}{d^n} e^{-2n\lambda_1 - 2n\epsilon} \frac{1}{q_0} \quad \text{if } \lambda_1 \notin \{k\lambda_2, k \geq 2\}, \\ \left[T \wedge \frac{i}{2} dW_{\hat{x}}^\epsilon \wedge d\overline{W}_{\hat{x}}^\epsilon \right] \left(B_x \left(\frac{1}{m_0} e^{-n\lambda_2 + 3n\epsilon} \right) \right) &\geq \frac{1}{d^n} e^{-2n\lambda_2 - 2n\epsilon} \frac{1}{q_0}. \end{aligned}$$

Proof. Let $\hat{x} \in \hat{\Omega}_\epsilon^n$ and let

$$E_n := f_{\hat{x}_n}^{-n} \left(B_{x_n} \left(\frac{1}{4m_0} \right) \right).$$

The inverse branch $f_{\hat{x}_n}^{-n}$ is well defined on $B_{x_n}(\frac{1}{4m_0})$ since $\hat{x}_n \in \hat{\Omega}_\epsilon$. Let g_n be the restriction of f^n on E_n . By using $f_{\hat{x}_n}^{-n} \circ g_n = \text{Id}_{E_n}$ and $T = \frac{1}{d^n} g_n^* T$ on E_n , we obtain

$$\begin{aligned} T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon &= \frac{1}{d^n} g_n^* T \wedge g_n^* (f_{\hat{x}_n}^{-n})^* \left(\frac{i}{2} dZ_{\hat{x}_n}^\epsilon \wedge d\overline{Z}_{\hat{x}_n}^\epsilon \right) \\ &= \frac{1}{d^n} g_n^* \left[T \wedge \frac{i}{2} (dZ_{\hat{x}_n}^\epsilon \circ (f_{\hat{x}_n}^{-n})) \wedge d(\overline{Z}_{\hat{x}_n}^\epsilon \circ (f_{\hat{x}_n}^{-n})) \right] \end{aligned}$$

on the open subset E_n . Now we use (2.1) to write $Z_{\hat{x}_n}^\epsilon \circ (f_{\hat{x}_n}^{-n}) = \alpha_{n, \hat{x}_n} Z_{\hat{x}_n}$. Since $|\alpha_{n, \hat{x}_n}|^2 \geq e^{-2n\lambda_1 - 2n\epsilon}$, we get on E_n :

$$T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon \geq \frac{1}{d^n} e^{-2n\lambda_1 - 2n\epsilon} g_n^* \left[T \wedge \frac{i}{2} dZ_{\hat{x}_n}^\epsilon \wedge d\overline{Z}_{\hat{x}_n}^\epsilon \right]. \quad (3.2)$$

Now we bound from above the left hand side and we bound from below the right hand side (applied to E_n). Using Lemma 2.3 with $r = 1/m_0$ and $n \geq n_1(L_0)$ we obtain $E_n \subset B_x \left(\frac{1}{m_0} e^{-n\lambda_2 + 3n\epsilon} \right)$, hence

$$T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon \left(B_x \left(\frac{1}{m_0} e^{-n\lambda_2 + 3n\epsilon} \right) \right) \geq T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon (E_n). \quad (3.3)$$

For the right hand side, since g_n is injective on E_n and $g_n(E_n) = B_{x_n} \left(\frac{1}{4m_0} \right)$, we get

$$g_n^* \left[T \wedge \frac{i}{2} (dZ_{\hat{x}_n}^\epsilon \wedge d\overline{Z}_{\hat{x}_n}^\epsilon) \right] (E_n) = \left[T \wedge \frac{i}{2} (dZ_{\hat{x}_n}^\epsilon \wedge d\overline{Z}_{\hat{x}_n}^\epsilon) \right] \left(B_{x_n} \left(\frac{1}{4m_0} \right) \right) \geq \frac{1}{q_0}, \quad (3.4)$$

where the inequality comes from $\hat{x}_n \in \hat{\Omega}_\epsilon$. By combining (3.2), (3.3) and (3.4) we obtain

$$\left[T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon \right] \left(B_x \left(\frac{1}{m_0} e^{-n\lambda_2 + 3n\epsilon} \right) \right) \geq \frac{1}{d^n} e^{-2n\lambda_1 - 2n\epsilon} \frac{1}{q_0}.$$

We use $W_{\hat{x}}^\epsilon \circ (f_{\hat{x}_n}^{-n}) = \beta_{n, \hat{x}} W_{\hat{x}_n}$ and $|\beta_{n, \hat{x}}|^2 \geq e^{-2n\lambda_2 - 2n\epsilon}$ to prove the other estimate. \square

3.2 Proof of Theorem 1.2

We take the notations of Section 3.1. Let

$$\hat{\Theta}_\epsilon := \limsup_{n \in \mathbb{N}} \hat{\Omega}_\epsilon \cap \hat{f}^{-n}(\hat{\Omega}_\epsilon) = \limsup_{n \in \mathbb{N}} \hat{\Omega}_\epsilon^n.$$

We have $\hat{\nu}(\hat{\Theta}_\epsilon) \geq 1 - 2\delta$ according to (3.1). Let $\hat{x} \in \hat{\Theta}_\epsilon$, there exists an increasing sequence of integers $(l_p)_p$ such that

$$\hat{x} \in \hat{\Omega}_\epsilon \cap \hat{f}^{-l_p}(\hat{\Omega}_\epsilon) = \hat{\Omega}_\epsilon^{l_p}.$$

Proposition 3.3 then asserts for p large enough:

$$\left[T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\bar{Z}_{\hat{x}}^\epsilon \right] \left(B_x \left(\frac{1}{m_0} e^{-l_p \lambda_2 + 3l_p \epsilon} \right) \right) \geq \frac{1}{d^{l_p}} e^{-2l_p \lambda_1 - 2l_p \epsilon} \frac{1}{q_0}.$$

If p is also large enough so that $e^{l_p \epsilon} \geq \frac{1}{m_0}$ and $\frac{1}{q_0} \geq e^{-l_p \epsilon}$, we obtain with $r_p := e^{-l_p(\lambda_2 - 4\epsilon)}$:

$$\left[T \wedge \frac{i}{2} dZ_{\hat{x}}^\epsilon \wedge d\bar{Z}_{\hat{x}}^\epsilon \right] (B_x(r_p)) \geq e^{-l_p(\log d + 2\lambda_1 + 3\epsilon)} = r_p^{(\log d + 2\lambda_1 + 3\epsilon)/(\lambda_2 - 4\epsilon)}.$$

Since $(r_p)_p$ tends to 0 and $\hat{\nu}(\hat{\Theta}_\epsilon) > 0$, we get

$$\underline{d}_{T,Z}(\nu) \leq \frac{\log d + 2\lambda_1 + 3\epsilon}{\lambda_2 - 4\epsilon} =: 2 \frac{\lambda_1}{\lambda_2} + \frac{\log d}{\lambda_2} + O(\epsilon).$$

We prove

$$\underline{d}_{T,W}(\nu) \leq \frac{\log d + 2\lambda_2 + 3\epsilon}{\lambda_2 - 4\epsilon} =: 2 + \frac{\log d}{\lambda_2} + O(\epsilon)$$

in a similar way.

3.3 Monge-Ampère mass

We prove Proposition 3.1. Let $x \in \text{Supp } \mu$, let V be a neighbourhood of x and let $Z : V \rightarrow \mathbb{C}$ be a holomorphic coordinate (submersion) on V . We want to prove that the positive measure $T \wedge \frac{i}{2} dZ \wedge d\bar{Z}$ is not the zero measure on V . With no loss of generality, we can assume that $x = (0, 0)$, $V = \mathbb{D}(2) \times \mathbb{D}(2)$ and $Z(z, w) = z$. Let also $T = 2i\partial\bar{\partial}G$ on V , where G is a continuous psh function, see Section 1.2. We denote $\sigma_z(u) := (z, u)$.

Lemma 3.4. *If $(T \wedge \frac{i}{2} dZ \wedge d\bar{Z})(\mathbb{D}(2) \times \mathbb{D}(2)) = 0$, then $G \circ \sigma_z$ is harmonic on \mathbb{D} for every $z \in \mathbb{D}$.*

Proof. Let $z_0 \in \mathbb{D}$ and let $\varphi \in C_0^\infty(\mathbb{D})$ be a non negative test function. Let $\psi \in C_0^\infty(\mathbb{D}^2)$ non negative such that $\psi \circ \sigma_{z_0} = \varphi$ on \mathbb{D} . According to Proposition A.4, we have

$$\left(T \wedge \frac{i}{2} dZ \wedge d\bar{Z} \right) (\psi) = \int_{z \in \mathbb{D}} \left(\int_{w \in \mathbb{D}} (G \circ \sigma_z)(w) \times \Delta(\psi \circ \sigma_z)(w) \, d\text{Leb}(w) \right) \, d\text{Leb}(z),$$

which is equal to zero by our assumption. Since G is psh, the measurable function

$$z \mapsto \int_{w \in \mathbb{D}} (G \circ \sigma_z)(w) \times \Delta(\psi \circ \sigma_z)(w) \, d\text{Leb}(w)$$

is non negative, hence there exists $A \subset \mathbb{D}$ such that $\text{Leb}(A) = \text{Leb}(\mathbb{D})$ and

$$\forall z \in A, \quad \int_{w \in \mathbb{D}} (G \circ \sigma_z)(w) \times \Delta(\psi \circ \sigma_z)(w) \, d\text{Leb}(w) = 0. \quad (3.5)$$

Let us extend this property to every $z \in \mathbb{D}$. Since A is a dense subset of \mathbb{D} , there exists a sequence $(z_n)_{n \geq 1}$ of points in A which converges to z . Using (3.5), we get

$$\forall n \geq 1, \quad \int_{w \in \mathbb{D}} (G \circ \sigma_{z_n})(w) \times \Delta(\psi \circ \sigma_{z_n})(w) \, d\text{Leb}(w) = 0. \quad (3.6)$$

Since G is continuous on $\overline{\mathbb{D}^2}$ and ψ is smooth on $\overline{\mathbb{D}^2}$, G and $2i\partial\bar{\partial}\psi$ are uniformly continuous on $\overline{\mathbb{D}^2}$. This implies that $G \circ \sigma_{z_n}$ uniformly converges to $G \circ \sigma_z$ on \mathbb{D} and that $\Delta(\psi \circ \sigma_{z_n})$ uniformly converges to $\Delta(\psi \circ \sigma_z)$ on \mathbb{D} . Taking the limits in (3.6), we get

$$\forall z \in \mathbb{D}, \quad \int_{w \in \mathbb{D}} (G \circ \sigma_z)(w) \times \Delta(\psi \circ \sigma_z)(w) \, d\text{Leb}(w) = 0.$$

In particular, we obtain using $\psi \circ \sigma_{z_0} = \varphi$:

$$\int_{w \in \mathbb{D}} (G \circ \sigma_{z_0})(w) \times \Delta\varphi(w) \, d\text{Leb}(w) = 0.$$

This holds for every non negative $\varphi \in C_0^\infty(\mathbb{D})$, hence $G \circ \sigma_{z_0}$ is harmonic on \mathbb{D} . \square

Now we use the following result, see [6, Lemme IV.1.1] and [27, Section A.10].

Theorem 3.5 (Briend). *Let G be a continuous psh function on $\mathbb{D}(2) \times \mathbb{D}(2)$. Let E be the set of points $p \in \mathbb{D}(\frac{1}{4}) \times \mathbb{D}(\frac{1}{4})$ such that there exists a holomorphic disc $\sigma_p : \mathbb{D} \rightarrow \mathbb{D}(2) \times \mathbb{D}(2)$ satisfying*

1. *the boundary of σ_p is outside $\mathbb{D}(\frac{1}{2}) \times \mathbb{D}(\frac{1}{2})$,*
2. *$G \circ \sigma_p$ is harmonic on \mathbb{D} .*

Then $(2i\partial\bar{\partial}G \wedge 2i\partial\bar{\partial}G)(E) = 0$.

In our situation, we have $\mathbb{D}(\frac{1}{4}) \times \mathbb{D}(\frac{1}{4}) = E$ by taking for σ_p the discs $\sigma_z : \mathbb{D} \rightarrow \mathbb{D} \times \mathbb{D}$, $u \mapsto (z, u)$. Indeed, the boundary of σ_z is contained in $\{z\} \times \partial\mathbb{D}$ and $G \circ \sigma_z$ is harmonic on \mathbb{D} according to Lemma 3.4. Theorem 3.5 then gives:

$$(2i\partial\bar{\partial}G \wedge 2i\partial\bar{\partial}G)(\mathbb{D}(\frac{1}{4}) \times \mathbb{D}(\frac{1}{4})) = 0,$$

which contradicts $x = 0 \in \text{Supp } \mu = \text{Supp}(2i\partial\bar{\partial}G \wedge 2i\partial\bar{\partial}G)$.

4 Lower bounds for the directional dimensions of T and S

We prove Theorem 1.3 and Corollary 1.5. Let ν be a dilating measure with exponents $\lambda_1 > \lambda_2$, and let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . Theorems 4.6 and 4.7 provide lower bounds for directional dimensions depending on d_ν , Theorem 1.3 follows from the lower bound (1.4) on d_ν . We shall use arguments of [13]. Precisely, we shall substitute the lower bound $(f^n)^*\omega \geq e^{2n\lambda_2}e^{-n\epsilon}\omega$ obtained in [13] by slicing arguments, by the two lower bounds

$$(f^n)^*\omega \geq e^{2n\lambda_1}e^{-6n\epsilon}\frac{i}{2}dZ \wedge d\bar{Z} \quad \text{and} \quad (f^n)^*\omega \geq e^{2n\lambda_2}e^{-6n\epsilon}\frac{i}{2}dW \wedge d\bar{W}$$

coming from Proposition 2.6. In Sections 4.1 and 4.2 we construct a set $\hat{\Lambda}_\epsilon \subset \hat{X}$ of uniformizations satisfying $\hat{\nu}(\hat{\Lambda}_\epsilon) \geq 1 - \delta/2$. Section 4.3 deals with separated sets.

4.1 Dynamical balls

The dynamical distance is defined by $d_n(x, y) := \max_{0 \leq k \leq n} d(f^k(x), f^k(y))$. We denote by $B_n(x, r)$ the ball centered at x and of radius r for the distance d_n .

Lemma 4.1. *Let $\delta > 0$ and $R > 0$. There exist $\eta_1 < R$, $n_3 \geq 1$ and $C \subset \mathbb{P}^2$ such that $\nu(C) \geq 1 - \delta/8$ and satisfying the following properties. For every $x \in C$ and $n \geq n_3$:*

$$\nu(B_n(x, \eta_1/8)) \geq e^{-nh_\nu - \epsilon n}.$$

$$\forall r \leq \eta_1, \nu(B_n(x, 5r)) \leq \nu(B_n(x, 5\eta_1)) \leq e^{-nh_\nu + \epsilon n}.$$

Proof. Brin-Katok Theorem [9] ensures that there exists $C_1 \subset \mathbb{P}^2$ of full ν -measure such that for every $x \in C_1$:

$$\lim_{r \rightarrow 0} \left(\liminf_{n \rightarrow +\infty} \frac{-1}{n} \log \nu(B_n(x, r)) \right) = \lim_{r \rightarrow 0} \left(\limsup_{n \rightarrow +\infty} \frac{-1}{n} \log \nu(B_n(x, r)) \right) = h_\nu.$$

Hence, for every $x \in C_1$ there exists $\eta_1(x) \leq R$ such that $r \leq \eta_1(x)$ implies

$$\liminf_{n \rightarrow +\infty} \frac{-1}{n} \log(\nu(B_n(x, 5r))) \geq h_\nu - \epsilon/2 \quad \text{and} \quad \limsup_{n \rightarrow +\infty} \frac{-1}{n} \log(\nu(B_n(x, r/8))) \leq h_\nu + \epsilon/2.$$

Let $\eta_1 > 0$ small enough such that $C_2 := \{x \in C_1, \eta_1 \leq \eta_1(x) \leq R\}$ satisfies $\nu(C_2) \geq 1 - \delta/16$. For every $x \in C_2$, there exists $n_3(x)$ such that $n \geq n_3(x)$ implies

$$\nu(B_n(x, \eta_1/8)) \geq e^{-nh_\nu - \epsilon n}.$$

$$\forall r \leq \eta_1, \nu(B_n(x, 5r)) \leq \nu(B_n(x, 5\eta_1)) \leq e^{-nh_\nu + \epsilon n}.$$

Let $n_3 \geq 1$ such that $C := C_2 \cap \{x \in C_1, n_3(x) \leq n_3\}$ satisfies $\nu(C) \geq 1 - \delta/8$. \square

For every $L > 0$, let m_L be the smallest integer m such that $Le^{-m(\lambda_2 + \epsilon)} \leq 1$ and let $n_4(L)$ be the smallest integer n larger than m_L such that $e^{-n\epsilon} \leq M^{-m_L}$, where $M := \max\{\|Df\|_{\infty, \mathbb{P}^2}, 1\}$. The set \hat{F} is defined in Theorem 2.1.

Lemma 4.2. *Let $\hat{x} \in \hat{F}$ such that $\eta_\epsilon(\hat{x}) \geq \eta$ and $L_\epsilon(\hat{x}) \leq L$. For every $n \geq n_4(L)$ and every $r \leq \eta$,*

$$f_{\hat{x}_n}^{-n}(B_{x_n}(re^{-2n\epsilon})) \subset B_n(x_0, r).$$

Proof. Let us observe that for every $0 \leq k \leq n$, f^k is injective on $f_{\hat{x}_n}^{-n}(B_{x_n}(re^{-2n\epsilon}))$ and that $f^k f_{\hat{x}_n}^{-n} = f_{\hat{x}_n}^{-n+k}$. By setting $p = n - k$, it suffices to show that

$$\forall p \in \llbracket 0, n \rrbracket \quad f_{\hat{x}_n}^{-p}(B_{x_n}(re^{-2n\epsilon})) \subset B_{x_{n-p}}(r). \quad (4.1)$$

To simplify let us set $m := m_L$ and $n_4 := n_4(L)$. We immediately have $n_4 \geq m$ and also

$$\forall n \geq n_4, \forall p \in \llbracket 0, n \rrbracket, \quad f_{\hat{x}_n}^{-p}(B_{x_n}(re^{-2n\epsilon})) \subset f_{\hat{x}_n}^{-p}(B_{x_n}(\frac{r}{M^m}e^{-n\epsilon})). \quad (4.2)$$

To verify (4.1), we shall consider separately the cases $p \leq m$ and $p > m$. We know that for every p , $\text{Lip } f_{\hat{x}_n}^{-p} \leq L(\hat{x}_n)e^{-p\lambda_2+p\epsilon} \leq Le^{n\epsilon}e^{-p\lambda_2+p\epsilon}$ on $B_{x_n}(\eta_\epsilon(\hat{x}_n))$, which contains $B_{x_n}(\eta e^{-n\epsilon})$. Hence for every $n \geq n_4 \geq m$, $p \in \llbracket m, n \rrbracket$ and $r \leq \eta$:

$$f_{\hat{x}_n}^{-p}(B_{x_n}(re^{-n\epsilon})) \subset B_{x_{n-p}}(re^{-n\epsilon}Le^{n\epsilon}e^{-p\lambda_2+p\epsilon}) = B_{x_{n-p}}(rLe^{-p\lambda_2+p\epsilon}) \subset B_{x_{n-p}}(r).$$

Since $M^m \geq 1$ this implies for every $n \geq n_4 \geq m$, $p \in \llbracket m, n \rrbracket$ and $r \leq \eta$:

$$f_{\hat{x}_n}^{-p}(B_{x_n}(\frac{r}{M^m}e^{-n\epsilon})) \subset B_{x_{n-p}}(\frac{r}{M^m}). \quad (4.3)$$

Thus, by using (4.2) and $M^m \geq 1$:

$$\forall p \in \llbracket m, n \rrbracket, \quad f_{\hat{x}_n}^{-p}(B_{x_n}(re^{-2n\epsilon})) \subset B_{x_{n-p}}(r). \quad (4.4)$$

We have proved (4.1) for $p \in \llbracket m, n \rrbracket$. Let us show this inclusion for $p \in \llbracket 0, m \rrbracket$. For every $p \in \llbracket 0, m \rrbracket$, let us set $p = m - p'$ where $p' \in \llbracket 0, m \rrbracket$. Then

$$f_{\hat{x}_n}^{-p}(B_{x_n}(\frac{r}{M^m}e^{-n\epsilon})) = f^{p'}(f_{\hat{x}_n}^{-m}(B_{x_n}(\frac{r}{M^m}e^{-n\epsilon}))) \subset f^{p'}(B_{x_{n-m}}(\frac{r}{M^m})),$$

where the inclusion comes from (4.3) with $p = m$. Using $e^{-n\epsilon} \leq 1/M^m$ for the left hand side, and $\|Df^{p'}\|_{\infty, \mathbb{P}^2} \leq M^{p'}$ for the right hand side, we get:

$$\forall p \in \llbracket 0, m \rrbracket, \quad f_{\hat{x}_n}^{-p}(B_{x_n}(re^{-2n\epsilon})) \subset B_{x_{n-m+p'}}(\frac{r}{M^m}M^{p'}) \subset B_{x_{n-p}}(r). \quad (4.5)$$

We finally obtain (4.1) by combining (4.4) and (4.5). \square

4.2 Uniformizations

In this Section we introduce uniformizations for the functions of Theorem 2.1 and for radius of balls in relation with their ν -measure or their directional measures. This will allow us to consider those quantities as constant on Borel sets of $\hat{\nu}$ -measure close to 1.

1 - Control of the functions $n_\epsilon, \rho_\epsilon, L_\epsilon, \eta_\epsilon, \beta_\epsilon$ of Theorem 2.1

We recall that \hat{F} provided by Theorem 2.1 satisfies $\hat{\nu}(\hat{F}) = 1$. Let $n_0 \geq 1, \rho_0 > 0, L_0 \geq 1, \eta_0 > 0, \beta_0 \geq 1$ such that

$$\Lambda^{(1)} := \left\{ \hat{x} \in \hat{F}, n_\epsilon(\hat{x}) \leq n_0, \rho_\epsilon(\hat{x}) \geq \rho_0, L_\epsilon(\hat{x}) \leq L_0, \eta_\epsilon(\hat{x}) \geq \eta_0, \beta_\epsilon(\hat{x}) \leq \beta_0 \right\}$$

satisfies $\hat{\nu}(\Lambda^{(1)}) \geq 1 - \delta/8$.

2 - Uniformization of the directional dimensions

Let S be a positive closed current on \mathbb{P}^2 whose support contains the support of ν . Let $r_1 > 0$ such that

$$\Lambda^{(2)} := \left\{ \hat{x} \in \hat{F}, \forall r \leq r_1, \begin{array}{l} r^{\overline{d_{S,Z}(\hat{x})+\epsilon}} \leq (S \wedge (\frac{i}{2}dZ_{\hat{x}}^\epsilon \wedge d\overline{Z}_{\hat{x}}^\epsilon))(B_{x_0}(r)) \leq r^{\underline{d_{S,Z}(\hat{x})-\epsilon}} \\ (S \wedge (\frac{i}{2}dW_{\hat{x}}^\epsilon \wedge d\overline{W}_{\hat{x}}^\epsilon))(B_{x_0}(r)) \leq r^{\underline{d_{S,W}(\hat{x})-\epsilon}} \end{array} \right\}$$

satisfies $\hat{\nu}(\Lambda^{(2)}) \geq 1 - \delta/8$.

3 - Uniformization of the dimension of the measure

Let $r_2 > 0$ such that

$$D := \left\{ x \in \mathbb{P}^2, \forall r \leq r_2, \nu(B_x(r)) \leq r^{\underline{d_\nu}-\epsilon} \right\}$$

satisfies $\nu(D) \geq 1 - \delta/8$. We set $\Lambda^{(3)} := \pi_0^{-1}(D) \cap \hat{F}$.

4 - Measure of dynamical balls, definition of η_1

We apply Lemma 4.1 with $R := \min\{\eta_0, r_1, r_2\}$. There exist $\eta_1 \leq R, n_3 \geq 1$ and $C \subset \mathbb{P}^2$ such that $\nu(C) \geq 1 - \delta/8$ and for every $x \in C$ and $n \geq n_3$:

$$\nu(B_n(x, \eta_1/8)) \geq e^{-nh_\nu - \epsilon n},$$

$$\forall r \leq \eta_1, \nu(B_n(x, 5r)) \leq \nu(B_n(x, 5\eta_1)) \leq e^{-nh_\nu + \epsilon n}.$$

We denote $\Lambda^{(4)} := \pi_0^{-1}(C) \cap \hat{F}$.

5 - Definitions of $\hat{\Lambda}_\epsilon$ and N_ϵ

We set

$$\hat{\Lambda}_\epsilon := \Lambda^{(1)} \cap \Lambda^{(2)} \cap \Lambda^{(3)} \cap \Lambda^{(4)}.$$

We have $\hat{\nu}(\hat{\Lambda}_\epsilon) \geq 1 - \delta/2$. We recall that $n_1(L), n_2(\beta)$ and $n_4(L)$ are defined before Lemma 2.3, Proposition 2.6 and Lemma 4.2. Let n_5 be the smallest integer n such that $e^{-n\epsilon} \leq 1/2$ and $2e^{-n(\lambda_1+\epsilon)} < 1$. We set

$$N_\epsilon := \max\{n_0, n_1(L_0), n_2(\beta_0), n_3, n_4(L_0), n_5\}.$$

6 - Definition of $\hat{\Delta}_\epsilon^n$

For all $n \geq N_\epsilon$, we define

$$\hat{\Delta}_\epsilon^n := \hat{F} \cap \hat{f}^{-n} \{n_\epsilon(\hat{x}) \leq n\} = \{\hat{x} \in \hat{F}, n_\epsilon(\hat{x}_n) \leq n\}.$$

Since $\hat{\nu}$ is \hat{f} -invariant and $\Lambda^{(1)} \subset \{n_\epsilon(\hat{x}) \leq n\}$ for every $n \geq N_\epsilon$, we have $\hat{\nu}(\hat{\Delta}_\epsilon^n) \geq \hat{\nu}(\Lambda^{(1)}) \geq 1 - \delta/8$. Hence

$$\forall n \geq N_\epsilon, \quad \hat{\nu}(\hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n) \geq 1 - \delta.$$

4.3 Separated sets

A subset $\{x_1, \dots, x_N\} \subset \mathbb{P}^2$ is r -separated if $d(x_i, x_j) \geq r$ for every $i \neq j$. For $A \subset \mathbb{P}^2$, a subset $\{x_1, \dots, x_N\} \subset A$ is maximal r -separated with respect to A if it is r -separated and if for every $y \in A$, there exists $i \in \{1, \dots, N\}$ such that $d(y, x_i) < r$. We use similar definitions for the distance d_n , in which case the subsets are called (n, r) separated.

Lemma 4.3. *Let $A \subset \pi_0(\hat{\Lambda}_\epsilon)$ such that $\nu(A) > 0$ and let $c \in]0, 1]$. Let $n \geq N_\epsilon$ and let $\{x_1, \dots, x_{N_n}\} \subset A$ be a maximal $(n, c\eta_1)$ -separated with respect to A . Then*

1. $A \subset \cup_{i=1}^{N_n} B_n(x_i, c\eta_1)$ and $B_n(x_i, c\eta_1/2) \cap B_n(x_j, c\eta_1/2) = \emptyset$ for every $i \neq j$.
2. $\nu(B_n(x_i, c\eta_1)) \leq e^{-nh_\nu + n\epsilon}$ and $e^{-nh_\nu - n\epsilon} \leq \nu(B_n(x_i, c\eta_1))$ if $c \geq 1/8$.
3. $N_n \geq \nu(A)e^{nh_\nu - n\epsilon}$.

Proof. Item 1 comes from the definitions, and Item 2 from §4 of Section 4.2. They imply $\nu(A) \leq \sum_{i=1}^{N_n} \nu(B_n(x_i, c\eta_1)) \leq N_n e^{-nh_\nu + n\epsilon}$, which gives Item 3. \square

The next lemmas give an amount of concentrated separation. We take the arguments of de Thélin-Vigny [13, Section 7]. By applying Lemma 4.3 with $c = 1/4$ we obtain $\nu(B_n(x_i, \eta_1/4)) \geq e^{-nh_\nu - n\epsilon}$ for every x_i in a maximal $(n, \eta_1/4)$ -separated subset of A . Lemma 4.4 allows to select a large number of x_i such that $\nu(B_n(x_i, \eta_1/4) \cap A) \geq e^{-nh_\nu - 2n\epsilon}$ and which are (n, η_1) -separated, then Lemma 4.5 deals with these x_i . Let n_δ be the smallest n such that $e^{-n\epsilon} \leq \delta/2$.

Lemma 4.4. *Let $A \subset \pi_0(\hat{\Lambda}_\epsilon)$ such that $\nu(A) \geq \delta$. For every $n \geq \max\{N_\epsilon, n_\delta\}$, there exist a $(n, \eta_1/4)$ -separated subset $\{x_1, \dots, x_{N_{n,2}}\}$ of A and $N_{n,3} \in \llbracket 1, N_{n,2} \rrbracket$ such that*

1. for $i \neq j$ in $\llbracket 1, N_{n,2} \rrbracket$, $B_n(x_i, \eta_1/8) \cap B_n(x_j, \eta_1/8) = \emptyset$.
2. for $i \in \llbracket 1, N_{n,2} \rrbracket$, $\nu(B_n(x_i, \eta_1/4) \cap A) \geq e^{-nh_\nu - 2\epsilon n}$, where $N_{n,2} \geq \nu(A)e^{nh_\nu - 2n\epsilon}$.
3. for $i \in \llbracket 1, N_{n,2} \rrbracket$ and $c \geq 1/8$, $e^{-nh_\nu - n\epsilon} \leq \nu(B_n(x_i, c\eta_1))$.
4. for $i \neq j$ in $\llbracket 1, N_{n,3} \rrbracket$, $B_n(x_i, \eta_1/2) \cap B_n(x_j, \eta_1/2) = \emptyset$, where $N_{n,3} \geq \nu(A)e^{nh_\nu - 4n\epsilon}$.

Proof. Let us apply Lemma 4.3 with $c = 1/4$ and $n \geq \max\{N_\epsilon, n_\delta\}$. There exists a maximal $(n, \eta_1/4)$ -separated subset $\{x_1, \dots, x_{N_{n,1}}\}$ of A satisfying:

- $A \subset \cup_{i=1}^{N_{n,1}} B_n(x_i, \eta_1/4)$ and $B_n(x_i, \eta_1/8) \cap B_n(x_j, \eta_1/8) = \emptyset$ for every $i \neq j$,
- $e^{-nh_\nu - n\epsilon} \leq \nu(B_n(x_i, \eta_1/8))$,
- $N_{n,1} \geq \nu(A)e^{nh_\nu - n\epsilon}$.

Let us set $I := \{1 \leq i \leq N_{n,1}, \nu(B_n(x_i, \eta_1/4) \cap A) \geq e^{-nh_\nu - 2\epsilon n}\}$. Let $N_{n,2}$ be the cardinality of I , and assume that I consists of the first $N_{n,2}$ elements of $\llbracket 1, N_{n,1} \rrbracket$. We want to bound $N_{n,2}$ from below. We know that $A \subset \cup_{i=1}^{N_{n,1}} B_n(x_i, \eta_1/4)$, hence

$$\nu(A) \leq \sum_{i=1}^{N_{n,2}} \nu(B_n(x_i, \eta_1/4) \cap A) + \sum_{i=N_{n,2}+1}^{N_{n,1}} \nu(B_n(x_i, \eta_1/4) \cap A).$$

If $i \notin \llbracket 1, N_{n,2} \rrbracket$, we have $\nu(B_n(x_i, \eta_1/4) \cap A) < e^{-nh_\nu - 2\epsilon n}$ by definition of I . Otherwise, $\nu(B_n(x_i, \eta_1/4)) \leq e^{-nh_\nu + \epsilon n}$ since $x_i \in C$. This implies

$$\nu(A) \leq N_{n,2}e^{-nh_\nu + \epsilon n} + (N_{n,1} - N_{n,2})e^{-nh_\nu - 2\epsilon n}. \quad (4.6)$$

Let us give an upper bound for $N_{n,1}$. Since the balls $B_n(x_i, \eta_1/8)$ are pairwise disjoint and since $\nu(B_n(x_i, \eta_1/8)) \geq e^{-nh_\nu - \epsilon n}$, we get $e^{nh_\nu + \epsilon n} \geq N_{n,1} \geq N_{n,1} - N_{n,2}$. Combining this and (4.6), we obtain

$$\nu(A) \leq N_{n,2}e^{-nh_\nu + \epsilon n} + e^{-\epsilon n}.$$

Since $n \geq n_\delta$, we have $e^{-n\epsilon} \leq \delta/2 \leq \nu(A)/2$, and hence $N_{n,2} \geq \nu(A)e^{nh_\nu - \epsilon n}/2$. We deduce $N_{n,2} \geq \nu(A)e^{nh_\nu - 2\epsilon n}$ because $n \geq N_\epsilon \geq n_5$. Let us verify the last item. Let E_n be a maximal (n, η_1) -separated subset of $\{x_1, \dots, x_{N_{n,2}}\}$. By reordering we can assume that $E_n = \{x_1, \dots, x_{N_{n,3}}\}$. For every $i \in \llbracket 1, N_{n,3} \rrbracket$, let m_i be the cardinality of $\{j \in \llbracket 1, N_{n,2} \rrbracket, x_j \in B_n(x_i, \eta_1)\}$, and let $m_{i_0} := \max m_i$, simply denoted m in the sequel. We have $N_{n,2} \leq mN_{n,3}$. Let us verify that $m \leq e^{2n\epsilon}$, which implies $N_{n,3} \geq N_{n,2}e^{-2n\epsilon}$ as desired. For that purpose, let $\{x'_1, \dots, x'_m\} := B_n(x_{i_0}, \eta_1) \cap \{x_1, \dots, x_{N_{n,2}}\}$. We obviously have $\cup_{i=1}^m B_n(x'_i, \eta_1/8) \subset B_n(x_{i_0}, 2\eta_1)$. Since the dynamical balls $B_n(x'_i, \eta_1/8)$ are pairwise disjoint, we finally get $me^{-nh_\nu - n\epsilon} \leq e^{-nh_\nu + n\epsilon}$ by considering the ν -measures and using $x'_i, x_{i_0} \in C$ (see §4 of Section 4.2). \square

Now for $1 \leq i \leq N_{n,3}$ we put in $B_n(x_i, \eta_1/2)$ a lot of balls centered in $B_n(x_i, \eta_1/4) \cap A$.

Lemma 4.5. *Let $A \subset \pi_0(\hat{\Lambda}_\epsilon)$ such that $\nu(A) > 0$. Let $x \in A$ and let $n \geq N_\epsilon$ such that*

$$\nu(B_n(x, \eta_1/4) \cap A) \geq e^{-nh_\nu - 2n\epsilon}.$$

Let $\{y_1, \dots, y_{M_n}\}$ be a maximal $2\eta_1 e^{-n\lambda_1 - 4n\epsilon}$ -separated subset in $B_n(x, \eta_1/4) \cap A$.

1. *for every $i \neq j$, $B(y_i, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) \cap B(y_j, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) = \emptyset$.*
2. *$B_n(x, \eta_1/4) \cap A \subset \cup_{i=1}^{M_n} B(y_i, 2\eta_1 e^{-n\lambda_1 - 4n\epsilon})$.*

$$3. B(y_i, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) \subset B_n(x, \eta_1/2).$$

$$4. M_n \geq e^{-nh_\nu - 2n\epsilon} \left(\frac{1}{2\eta_1} e^{n\lambda_1 + 4n\epsilon} \right)^{\underline{d}_\nu - \epsilon}.$$

Proof. Items 1 and 2 come from the definitions. Lemmas 2.3 and 4.2 successively give

$$B(y_i, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) \subset f_{\hat{y}_{i,n}}^{-n}(B_{y_{i,n}}(\eta_1 e^{-2n\epsilon}/4) \subset B_n(y_i, \eta_1/4).$$

Since $y_i \in B_n(x, \eta_1/4)$, we get $B_n(y_i, \eta_1/4) \subset B_n(x, \eta_1/2)$, and Item 3 follows. Item 2 implies

$$\nu(B_n(x, \eta_1/4) \cap A) \leq \sum_{i=1}^{M_n} \nu(B(y_i, 2\eta_1 e^{-n\lambda_1 - 4n\epsilon})). \quad (4.7)$$

By our assumption, the left hand side of (4.7) is larger than $e^{-nh_\nu - 2n\epsilon}$. For the right hand side, since $n \geq N_\epsilon \geq n_5$, we have $2\eta_1 e^{-n\lambda_1 - 4n\epsilon} < \eta_1 \leq r_2$ by the definition of n_5 and η_1 (given in §5 and §4 of Section 4.2). Then, by using $y_i \in A \subset \pi_0(\hat{\Lambda}_\epsilon) \subset D$, we get by the definition of D (given in §3 of Section 4.2):

$$\nu(B(y_i, 2\eta_1 e^{-n\lambda_1 - 4n\epsilon})) \leq (2\eta_1 e^{-n\lambda_1 - 4n\epsilon})^{\underline{d}_\nu - \epsilon}.$$

We finally deduce from (4.7) that $e^{-nh_\nu - 2n\epsilon} \leq M_n (2\eta_1 e^{-n\lambda_1 - 4n\epsilon})^{\underline{d}_\nu - \epsilon}$, as desired. \square

4.4 Lower bounds for the upper dimensions $\overline{d_{T,Z}}$ and $\overline{d_{T,W}}$

Theorem 4.6. *Let ν be a dilating measure whose exponents $\lambda_1 > \lambda_2$ do not resonate. Let $\epsilon > 0$ and let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . Then*

$$\overline{d_{T,Z}}(\nu) \geq 2 + \underline{d}_\nu - \frac{\log d}{\lambda_1} - O(\epsilon),$$

$$\overline{d_{T,W}}(\nu) \geq 2 \frac{\lambda_2}{\lambda_1} + \underline{d}_\nu - \frac{\log d}{\lambda_1} - O(\epsilon).$$

Proof. We recall that $\overline{d_{T,Z}}(\nu) := \overline{d_{T,Z}}(\hat{x})$ and $\overline{d_{T,W}}(\nu) = \overline{d_{T,W}}(\hat{x})$ for $\hat{\nu}$ -almost every \hat{x} (see Proposition 2.4). Let us set $\overline{d_{T,Z}} := \overline{d_{T,Z}}(\nu)$. We are going to show

$$(\lambda_1 + 4\epsilon)(\overline{d_{T,Z}} - \underline{d}_\nu + 2\epsilon) + 12\epsilon \geq 2\lambda_1 - \log d. \quad (4.8)$$

This yields as desired

$$\overline{d_{T,Z}} - \underline{d}_\nu \geq \frac{2\lambda_1}{\lambda_1 + 4\epsilon} - \frac{\log d + 12\epsilon}{\lambda_1 + 4\epsilon} - 2\epsilon = 2 - \frac{\log d}{\lambda_1} - O(\epsilon). \quad (4.9)$$

Let $\delta > 0$. Let $\hat{\Lambda}_\epsilon$, N_ϵ and $\hat{\Delta}_\epsilon^n$ be given by Section 4.2. For every $n \geq N_\epsilon$ we set $A_n := \pi_0(\hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n)$, it satisfies $\nu(A_n) \geq 1 - \delta$. Lemma 4.4 provides $\{x_1, \dots, x_{N_{n,3}}\} \subset A_n$ satisfying $N_{n,3} \geq \nu(A_n) e^{nh_\nu - 4n\epsilon}$, $\nu(B_n(x_i, \eta_1/4) \cap A_n) \geq e^{-nh_\nu - 2n\epsilon}$ and $B_n(x_i, \eta_1/2) \cap B_n(x_j, \eta_1/2) = \emptyset$ for $i \neq j$. Then, for every x_i , we set a maximal $2\eta_1 e^{-n\lambda_1 - 4n\epsilon}$ -separated subset $\{y_1^i, \dots, y_{M_n}^i\}$ in $B_n(x_i, \eta_1/4) \cap A_n$ given by Lemma 4.5. Let us note

that $B(y_j^i, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) \subset B_n(x_i, \eta_1/2)$. Finally, for every i we choose $\hat{x}_i \in \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n$ such that $\pi_0(\hat{x}_i) = x_i$, and for every j we choose $\hat{y}_j^i \in \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n$ such that $\pi_0(\hat{y}_j^i) = y_j^i$.

Now we use $d^n = \int_{\mathbb{P}^2} (f^n)_* T \wedge \omega$ provided by Proposition A.3 to get

$$d^n \geq \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} \int_{\mathbb{P}^2} (f^n)_* (1_{B(y_j^i, \eta_1 e^{-n\lambda_1 - 4n\epsilon})} T) \wedge \omega = \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} \int_{B(y_j^i, \eta_1 e^{-n\lambda_1 - 4n\epsilon})} T \wedge (f^n)^* \omega.$$

Lemma 2.3 with $\hat{y}_j^i \in \hat{\Lambda}_\epsilon$ implies

$$B(y_j^i, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) \subset f_{\hat{y}_j^i, n}^{-n} (B(y_{j,n}^i, \frac{\eta_1}{4} e^{-2n\epsilon})). \quad (4.10)$$

Since $\hat{y}_j^i \in \hat{\Delta}_\epsilon^n$, we can apply Proposition 2.6 to bound $(f^n)^* \omega$ from below:

$$d^n \geq \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} e^{2n\lambda_1 - 6n\epsilon} \left(T \wedge \frac{i}{2} dZ_{\hat{y}_j^i}^\epsilon \wedge d\overline{Z}_{\hat{y}_j^i}^\epsilon \right) (B_{y_j^i}(\eta_1 e^{-n\lambda_1 - 4n\epsilon})). \quad (4.11)$$

Now $\hat{y}_j^i \in \hat{\Lambda}_\epsilon \subset \Lambda^{(2)}$ and $n \geq N_\epsilon$, hence

$$d^n \geq \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} e^{2n\lambda_1 - 6n\epsilon} (\eta_1 e^{-n\lambda_1 - 4n\epsilon})^{\overline{d_{T,Z} + \epsilon}}.$$

Finally, we use the lower bounds for $N_{n,3}$ (Lemma 4.4) and for M_n (Lemma 4.5), and $\nu(A_n) \geq 1 - \delta$. We obtain for every $n \geq \max\{N_\epsilon, n_{1-\delta}\}$:

$$d^n \geq (1 - \delta) e^{nh_\nu - 4n\epsilon} \cdot e^{-nh_\nu - 2n\epsilon} \left(\frac{1}{2\eta_1} e^{n\lambda_1 + 4n\epsilon} \right)^{d_\nu - \epsilon} \cdot e^{2n\lambda_1 - 6n\epsilon} (\eta_1 e^{-n\lambda_1 - 4n\epsilon})^{\overline{d_{T,Z} + \epsilon}}.$$

Let us note that the entropy h_ν disappear for the benefit of d_ν , and we get

$$d^n \geq c e^{-12n\epsilon} \left(e^{n\lambda_1 + 4n\epsilon} \right)^{d_\nu - \overline{d_{T,Z} - 2\epsilon}} e^{2n\lambda_1}, \quad (4.12)$$

where $c := (1 - \delta) \eta_1^{\overline{d_{T,Z} + \epsilon}} / (2\eta_1)^{d_\nu - \epsilon}$. We obtain (4.8) by looking at the exponential growth rates. Similarly, we can prove

$$(\lambda_1 + 4\epsilon)(\overline{d_{T,W}} - \underline{d}_\nu + 2\epsilon) + 12\epsilon \geq 2\lambda_2 - \log d$$

by using again Proposition 2.6 to bound $(f^n)^* \omega$ from below. \square

4.5 Lower bounds for the upper dimensions $\overline{d_{S,Z}}$ and $\overline{d_{S,W}}$

Let S be a $(1,1)$ closed positive current of \mathbb{P}^2 . If S does not satisfy $f^*S = dS$, the directional dimensions may be not $\hat{\nu}$ -almost everywhere constant (see Proposition 2.4). In this case, in the manner of de Thélin-Vigny [13], we take on an adapted definition and obtain the following result. It implies Theorem 1.3 by using the lower bound (1.4).

Theorem 4.7. *Let S be a $(1, 1)$ closed positive current of \mathbb{P}^2 . Let ν be a dilating measure whose exponents $\lambda_1 > \lambda_2$ do not resonate. We assume that $\text{Supp}(\nu) \subset \text{Supp} S$. Let $\epsilon > 0$ and let $(Z, W)_\epsilon$ be Oseledec-Poincaré coordinates for (f, ν) . For every $\hat{\Lambda} \subset \hat{F}$ such that $\hat{\nu}(\hat{\Lambda}) > 0$, we set*

$$\overline{d_{S,Z}}(\hat{\Lambda}) := \sup_{\hat{x} \in \hat{\Lambda}} \overline{d_{S,Z}}(\hat{x}) \quad , \quad \overline{d_{S,W}}(\hat{\Lambda}) := \sup_{\hat{x} \in \hat{\Lambda}} \overline{d_{S,W}}(\hat{x}).$$

Then

$$\begin{aligned} \overline{d_{S,Z}}(\hat{\Lambda}) &\geq 2 + \underline{d}_\nu - \frac{\log d}{\lambda_1} - O(\epsilon), \\ \overline{d_{S,W}}(\hat{\Lambda}) &\geq 2 \frac{\lambda_2}{\lambda_1} + \underline{d}_\nu - \frac{\log d}{\lambda_1} - O(\epsilon). \end{aligned}$$

Proof. Let $2\delta := \hat{\nu}(\hat{\Lambda})$. We construct $\hat{\Lambda}_\epsilon$, N_ϵ and $\hat{\Delta}_\epsilon^n$ for the current S as in Section 4.2. We have $\hat{\nu}(\hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n) \geq 1 - \delta$ for every $n \geq N_\epsilon$, thus $\hat{\nu}(\hat{\Lambda} \cap \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n) \geq \delta > 0$. We follow the arguments of Theorem 4.6. Let $n \geq \max\{N_\epsilon, n_\delta\}$ and $A_n := \pi_0(\hat{\Lambda} \cap \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n)$. Lemma 4.4 provides $\{x_1, \dots, x_{N_{n,3}}\} \subset A_n$ satisfying $N_{n,3} \geq \nu(A_n) e^{nh_\nu - 4n\epsilon}$, $\nu(B_n(x_i, \eta_1/4) \cap A_n) \geq e^{-nh_\nu - 2\epsilon n}$ and $B_n(x_i, \eta_1/2) \cap B_n(x_j, \eta_1/2) = \emptyset$ for $i \neq j$. Then, for every x_i , let $\{y_1^i, \dots, y_{M_n}^i\}$ be a maximal $2\eta_1 e^{-n\lambda_1 - 4n\epsilon}$ -separated subset in $B_n(x_i, \eta_1/4) \cap A_n$ given by Lemma 4.5. We have the inclusions $B(y_j^i, \eta_1 e^{-n\lambda_1 - 4n\epsilon}) \subset B_n(x_i, \eta_1/2)$. Finally, for every i, j let $\hat{x}_i \in \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n$ such that $\pi_0(\hat{x}_i) = x_i$, and let $\hat{y}_j^i \in \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n$ such that $\pi_0(\hat{y}_j^i) = y_j^i$.

Using $d^n = \int_{\mathbb{P}^2} (f^n)_* S \wedge \omega$, we deduce as in (4.11):

$$d^n \geq \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} e^{2n\lambda_1 - 6n\epsilon} \left(S \wedge \frac{i}{2} dZ_{\hat{y}_j^i}^\epsilon \wedge d\overline{Z}_{\hat{y}_j^i}^\epsilon \right) \left(B_{y_j^i}(\eta_1 e^{-n\lambda_1 - 4n\epsilon}) \right).$$

Since $\hat{y}_j^i \in \hat{\Lambda}_\epsilon$ and $n \geq N_\epsilon$, we have

$$d^n \geq \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} e^{2n\lambda_1 - 6n\epsilon} (\eta_1 e^{-n\lambda_1 - 4n\epsilon}) \overline{d_{S,Z}}(\hat{y}_j^i) + \epsilon.$$

Then we use the inequality $\overline{d_{S,Z}}(\hat{y}_j^i) \leq \overline{d_{S,Z}}(\hat{\Lambda})$ and conclude as in the proof of Theorem 4.6. The lower bound concerning W is proved in a similar way. \square

4.6 Proof of Corollary 1.5

We recall that $\overline{d_T}(x)$ is μ -almost everywhere constant and denoted $\overline{d_T}(\mu)$, see Proposition 2.5. According to Proposition A.1, $\mu \ll T \wedge \omega$ implies

$$\overline{d_T}(\mu) \leq \overline{d_\mu}. \quad (4.13)$$

Let us analyse these quantities. On the one hand, Proposition A.2 yields $\overline{d_T}(\mu) = \min \left\{ \overline{d_{T,Z}}(x), \overline{d_{T,W}}(x) \right\}$ for μ -almost every $x \in \mathbb{P}^2$ and for every holomorphic coordinates

(Z, W) near x . On the other hand, since $\underline{d}_\mu = \overline{d}_\mu$, we have $\underline{d}_\mu = \overline{d}_\mu = \dim_H(\mu)$. Observe also that $\lambda_2 = \frac{1}{2} \log d$, since $\mu \ll T \wedge \omega$, see [19]. Hence $\dim_H(\mu) = 2 + \frac{\log d}{\lambda_1}$ by (1.6). Then one deduces from (4.13) that

$$\min \left\{ \overline{d_{T,Z}}(x), \overline{d_{T,W}}(x) \right\} \leq 2 + \frac{\log d}{\lambda_1}. \quad (4.14)$$

Now we use Theorem 4.6. If $(Z, W)_\epsilon$ are Oseledec-Poincaré coordinates for (f, μ) ,

$$\begin{aligned} \overline{d_{T,Z}}(\mu) &\geq 2 + \underline{d}_\mu - \frac{\log d}{\lambda_1} - O(\epsilon), \\ \overline{d_{T,W}}(\mu) &\geq 2 \frac{\lambda_2}{\lambda_1} + \underline{d}_\mu - \frac{\log d}{\lambda_1} - O(\epsilon). \end{aligned}$$

Let us note that $\lambda_2 = \frac{1}{2} \log d$ implies $\underline{d}_\mu - \frac{\log d}{\lambda_1} \geq 2$ by (1.4). We deduce that $\overline{d_{T,Z}}(\mu) \geq 4 - O(\epsilon)$ and that $\overline{d_{T,W}}(\mu) \geq 2 + \frac{\log d}{\lambda_1} - O(\epsilon)$. It remains to show that $\overline{d_{T,W}}(\mu) \leq 2 + \frac{\log d}{\lambda_1}$. Let ϵ small enough such that $4 - O(\epsilon) > 2 + \frac{\log d}{\lambda_1}$. This implies $\overline{d_{T,Z}}(\mu) > 2 + \frac{\log d}{\lambda_1}$, thus the minimum in (4.14) is attained for W .

5 Proof of Theorem 1.1

Let ν be a dilating measure with exponents $\lambda_1 \geq \lambda_2$ whose support is contained in the support of μ . If $\lambda_1 = \lambda_2$, then $\overline{d}_\nu = \underline{d}_\nu = 2 \frac{\log d}{\lambda_1}$ by (1.1), and Theorem 1.1 is true. So we can assume that $\lambda_1 > \lambda_2$. The proof of Theorem 1.1 will rely on the estimates of Proposition 3.3, on the arguments of Theorem 4.6 and on a rescaling of the time n with respect to the exponents. A crucial point is that Proposition 3.3 provides estimates for the directional measures of T in terms of λ_1, λ_2, d , we shall use them in (5.6). The dimension \underline{d}_ν will appear in (5.3).

Let $\epsilon > 0$. The set $\hat{\Omega}_\epsilon^n$ has been defined in Section 3.1, it satisfies $\hat{\nu}(\hat{\Omega}_\epsilon^n) \geq 1 - 2\delta$ for every $n \geq 1$. Let $\hat{\Lambda}_\epsilon, \hat{\Delta}_\epsilon^n$ be the sets defined in Section 4.2. We have $\hat{\nu}(\hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n \cap \hat{\Omega}_\epsilon^n) \geq 1 - 3\delta$ for every $n \geq N_\epsilon$. Now let K_n be the unique integer in $]\alpha_n, \alpha_n + 1]$, where $\alpha_n := \frac{1}{\lambda_2 - 3\epsilon}(-\log(\eta_1 m_0) + n(\lambda_1 + 4\epsilon))$. It satisfies

$$\eta_1 e^{-n\lambda_1 - 4n\epsilon} e^{-\lambda_2 + 3\epsilon} \leq \frac{1}{m_0} e^{-K_n \lambda_2 + 3K_n \epsilon} < \eta_1 e^{-n\lambda_1 - 4n\epsilon}, \quad (5.1)$$

these inequalities will be useful for (5.5). Let us have in mind that $K_n \simeq n\lambda_1/\lambda_2 \geq n \geq N_\epsilon$. Let $\{x_1, \dots, x_{N_{n,3}}\}$ be a $(n, \eta_1/4)$ -separated subset of $A_n := \pi_0(\hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n \cap \hat{\Omega}_\epsilon^{K_n})$ provided by Lemma 4.4. We have for every $n \geq \max\{N_\epsilon, n_{1-3\delta}\}$:

$$N_{n,3} \geq \nu(A_n) e^{nh_\nu - 4n\epsilon} \geq (1 - 3\delta) e^{nh_\nu - 4n\epsilon}. \quad (5.2)$$

Then, for every x_i , let $\{y_1^i, \dots, y_{M_n}^i\}$ be a maximal $2\eta_1 e^{-n\lambda_1 - 4n\epsilon}$ -separated subset of $B_n(x_i, \eta_1/4) \cap A_n$ provided by Lemma 4.5. The cardinality of this set satisfies:

$$M_n \geq e^{-nh_\nu - 2n\epsilon} \left(\frac{1}{2\eta_1} e^{n\lambda_1 + 4n\epsilon} \right)^{\underline{d}_\nu - \epsilon}. \quad (5.3)$$

For every $j \in \{1, \dots, M_n\}$, we set $\hat{y}_j^i \in \hat{\Lambda}_\epsilon \cap \hat{\Delta}_\epsilon^n \cap \hat{\Omega}_\epsilon^{K_n}$ such that $y_j^i = \pi_0(\hat{y}_j^i)$. By following the proof of Theorem 4.6 until Inequality (4.11), we get:

$$d^n \geq \sum_{i=1}^{N_{n,3}} \sum_{j=1}^{M_n} e^{2n\lambda_1 - 6n\epsilon} \left[T \wedge \left(\frac{i}{2} dZ_{\hat{y}_j^i}^\epsilon \wedge d\overline{Z_{\hat{y}_j^i}^\epsilon} \right) \right] (B_{y_j^i}(\eta_1 e^{-n\lambda_1 - 4n\epsilon})). \quad (5.4)$$

Now, according to (5.1),

$$B_{y_j^i}(\eta_1 e^{-n\lambda_1 - 4n\epsilon}) \supset B_{y_j^i} \left(\frac{1}{m_0} e^{-K_n \lambda_2 + 3K_n \epsilon} \right). \quad (5.5)$$

Since $\hat{y}_j^i \in \hat{\Omega}_\epsilon^{K_n}$, Proposition 3.3 implies for every n satisfying $n \geq N_\epsilon$ and $K_n \geq N_\epsilon$:

$$\left[T \wedge \left(\frac{i}{2} dZ_{\hat{y}_{i,j}^\epsilon} \wedge d\overline{Z_{\hat{y}_{i,j}^\epsilon}} \right) \right] (B_{y_j^i}(\eta_1 e^{-n\lambda_1 - 4n\epsilon})) \geq \frac{1}{d^{K_n}} e^{-2K_n \lambda_1 - 2K_n \epsilon} \frac{1}{q_0}. \quad (5.6)$$

We infer from (5.4) that for every n satisfying $n \geq N_\epsilon$ and $K_n \geq N_\epsilon$,

$$d^n \geq N_{n,3} \cdot M_n \cdot e^{2n\lambda_1 - 6n\epsilon} \frac{1}{d^{K_n}} e^{-2K_n \lambda_1 - 2K_n \epsilon} \frac{1}{q_0}. \quad (5.7)$$

By using the upper bounds for $N_{n,3}$ and M_n given by (5.2) and (5.3):

$$d^{n+K_n} \geq (1 - 3\delta) e^{nh_\nu - 4n\epsilon} \cdot e^{-nh_\nu - 2n\epsilon} \left(\frac{1}{2\eta_1} e^{n\lambda_1 + 4n\epsilon} \right)^{\underline{d}_\nu - \epsilon} \cdot e^{2n\lambda_1 - 6n\epsilon} e^{-2K_n \lambda_1 - 2K_n \epsilon} \frac{1}{q_0}.$$

If $C_1(\epsilon) := (1 - 3\delta)/q_0(2\eta_1)^{\underline{d}_\nu - \epsilon}$, we get:

$$\log d + \frac{K_n}{n} \log d \geq \frac{1}{n} \log C_1(\epsilon) - 12\epsilon + (\lambda_1 + 4\epsilon)(\underline{d}_\nu - \epsilon) + 2\lambda_1 - 2\frac{K_n}{n}(\lambda_1 + \epsilon).$$

By using the lower and upper bounds defining K_n just before (5.1), we obtain

$$\log d + \frac{\lambda_1 + 4\epsilon}{\lambda_2 - 3\epsilon} \log d \geq \frac{1}{n} \log C_2(\epsilon) - 12\epsilon + (\lambda_1 + 4\epsilon)(\underline{d}_\nu - \epsilon) + 2\lambda_1 - 2\frac{\lambda_1 + 4\epsilon}{\lambda_2 - 3\epsilon}(\lambda_1 + \epsilon),$$

where $C_2(\epsilon)$ is another constant. Letting n tend to $+\infty$ and then ϵ to 0, we get

$$\underline{d}_\nu \leq \frac{\log d}{\lambda_1} + \frac{\log d}{\lambda_2} + 2 \left(\frac{\lambda_1}{\lambda_2} - 1 \right).$$

To obtain the other upper bound, we use the analogue of (5.4) for W . Applying Proposition 3.3 with respect to W , we obtain instead of (5.7):

$$d^n \geq N_{n,3} \cdot M_n \cdot e^{2n\lambda_2 - 6n\epsilon} \frac{1}{d^{K_n}} e^{-2K_n \lambda_2 - 2K_n \epsilon} \frac{1}{q_0}.$$

Then we get

$$\underline{d}_\nu \leq \frac{\log d}{\lambda_1} + \frac{\log d}{\lambda_2} + 2 \left(1 - \frac{\lambda_2}{\lambda_1} \right),$$

which completes the proof of Theorem 1.1.

A Appendix

A.1 Dimension of measures

Proposition A.1. *Let ν_1 and ν_2 be two probability measures on \mathbb{P}^2 such that $\nu_1 \ll \nu_2$. Then for ν_1 -almost every $x \in \mathbb{P}^2$, we have:*

$$\underline{d}_{\nu_1}(x) \geq \underline{d}_{\nu_2}(x) \quad \text{and} \quad \overline{d}_{\nu_1}(x) \geq \overline{d}_{\nu_2}(x).$$

Proof. Let $\varphi \in L^1(\nu_2)$ such that $\nu_1(A) = \int_A \varphi d\nu_2$ for every Borel set A of \mathbb{P}^2 . Using the dominated convergence Theorem,

$$\lim_{M \rightarrow +\infty} \int_{\mathbb{P}^2} 1_{\{\varphi \leq M\}} \varphi d\nu_2 = \int_{\mathbb{P}^2} \varphi d\nu_2 = 1.$$

For every $n \geq 1$, we let M_n satisfy $\int_{\mathbb{P}^2} 1_{\{\varphi \leq M_n\}} \varphi d\nu_2 \geq 1 - \frac{1}{n}$. By the Lebesgue density Theorem, for ν_1 -almost every x in $\{\varphi \leq M_n\}$, we have

$$\lim_{r \rightarrow 0} \frac{\nu_1(B_x(r) \cap \{\varphi \leq M_n\})}{\nu_1(B_x(r))} = 1.$$

Then for every r small enough, we have

$$\frac{1}{2} \nu_1(B_x(r)) \leq \nu_1(B_x(r) \cap \{\varphi \leq M_n\}) \leq M_n \int_{B_x(r)} d\nu_2.$$

And thus $\nu_1(B_x(r)) \leq 2M_n \nu_2(B_x(r))$. Taking limits when r tends to 0, we deduce that

$$\underline{d}_{\nu_1}(x) \geq \underline{d}_{\nu_2}(x) \quad \text{and} \quad \overline{d}_{\nu_1}(x) \geq \overline{d}_{\nu_2}(x)$$

for ν_1 -almost every $x \in \{\varphi \leq M_n\}$. We end with $\nu_1(\cup_{n \in \mathbb{N}} \{\varphi \leq M_n\}) = 1$. \square

Proposition A.2. *Let S be a $(1, 1)$ -closed positive current on \mathbb{P}^2 . Let $x \in \mathbb{P}^2$ and let (Z, W) be holomorphic coordinates near x . Then*

$$\underline{d}_S(x) = \min \{ \underline{d}_{S,Z}(x), \underline{d}_{S,W}(x) \}, \quad \overline{d}_S(x) = \min \{ \overline{d}_{S,Z}(x), \overline{d}_{S,W}(x) \}.$$

Proof. Let us set $\sigma_{S,Z} = S \wedge (\frac{i}{2} dZ \wedge d\bar{Z})$ and $\sigma_{S,W} = S \wedge (\frac{i}{2} dW \wedge d\bar{W})$. There exists $c > 0$ such that $\frac{1}{c}(\sigma_{S,Z} + \sigma_{S,W}) \leq \sigma_S \leq c(\sigma_{S,Z} + \sigma_{S,W})$ on a neighbourhood of x , see [14] Chapter III, §3. We deduce for every r small enough

$$\frac{1}{c} \max [\sigma_{S,Z}(B_x(r)), \sigma_{S,W}(B_x(r))] \leq \sigma_S(B_x(r)) \leq 2c \max [\sigma_{S,Z}(B_x(r)), \sigma_{S,W}(B_x(r))].$$

We finish by observing that the local dimension of the maximum of two measures is equal to the minimum of the local dimensions, since one divides by $\log r$ which is negative. \square

A.2 Cohomology and slices

We refer to Sections 1.2 and A.3 of Dinh-Sibony's book [18].

Proposition A.3. *Let S be a $(1, 1)$ -closed positive current of \mathbb{P}^2 of mass 1. Let ω be the Fubini-Study form on \mathbb{P}^2 and let $f : \mathbb{P}^2 \rightarrow \mathbb{P}^2$ be an endomorphism of degree d . Then,*

$$\int_{\mathbb{P}^2} (f^n)_* S \wedge \omega = \int_{\mathbb{P}^2} S \wedge (f^n)^* \omega = d^n.$$

Proof. The first equality comes from the definition of duality. We verify the second one. By using $f^* \omega = d \cdot \omega + 2i\partial\bar{\partial}u$, where u is a smooth function on \mathbb{P}^2 , we obtain by induction

$$(f^n)^* \omega = d^n \omega + 2i\partial\bar{\partial}v_n,$$

where $v_n := (d^{n-1}u + \dots + du \circ f^{n-2} + u \circ f^{n-1})$. Since S is a closed current of mass 1, we have $\int_{\mathbb{P}^2} S \wedge 2i\partial\bar{\partial}v_n = 0$ and $\int_{\mathbb{P}^2} S \wedge d^n \omega = d^n$. \square

Proposition A.4. *Let G be a continuous psh function on \mathbb{D}^2 and let $S = 2i\partial\bar{\partial}G$. Let (Z, W) be the coordinates on $\mathbb{D}^2 = \mathbb{D} \times \mathbb{D}$ and let $\phi \in C_0^\infty(\mathbb{D}^2)$. Then*

$$\begin{aligned} S \wedge \frac{i}{2} dZ \wedge d\bar{Z}(\phi) &= \int_{z \in \mathbb{D}} \left(\int_{w \in \mathbb{D}} G_z(w) \times \Delta \phi_z(w) \, d\text{Leb}(w) \right) d\text{Leb}(z) \\ &= \int_{z \in \mathbb{D}} (\sigma_z^* S)(\phi_z) d\text{Leb}(z). \end{aligned}$$

where $\sigma_z : u \mapsto (z, u)$, $G_z := G \circ \sigma_z$ and $\phi_z := \phi \circ \sigma_z$.

Proof. By definition,

$$S \wedge \frac{i}{2} dZ \wedge d\bar{Z}(\phi) = 2i\partial\bar{\partial}G(\phi \frac{i}{2} dZ \wedge d\bar{Z}) = \int_{\mathbb{D}^2} G \cdot 2i\partial\bar{\partial}(\phi \frac{i}{2} dZ \wedge d\bar{Z}).$$

The computation

$$2i\partial\bar{\partial}(\phi \frac{i}{2} dZ \wedge d\bar{Z}) = 4 \left(\frac{\partial^2 \phi}{\partial w \partial \bar{w}} \right) \frac{i}{2} dW \wedge d\bar{W} \wedge \frac{i}{2} dZ \wedge d\bar{Z} = 4 \left(\frac{\partial^2 \phi}{\partial w \partial \bar{w}} \right) d\text{Leb}(z, w)$$

allows to write

$$\begin{aligned} S \wedge \frac{i}{2} dZ \wedge d\bar{Z}(\phi) &= \int_{(z, w) \in \mathbb{D}^2} G(z, w) \times 4 \frac{\partial^2 \phi}{\partial w \partial \bar{w}}(z, w) \, d\text{Leb}(z, w) \\ &= \int_{z \in \mathbb{D}} \left(\int_{w \in \mathbb{D}} G_z(w) \times \Delta \phi_z(w) \, d\text{Leb}(w) \right) d\text{Leb}(z), \end{aligned}$$

where the second equality comes from Fubini's Theorem. Finally, the quantity in brackets is equal to $(\Delta G_z)(\phi_z) = (\sigma_z^* S)(\phi_z)$. \square

References

- [1] F. Berteloot and C. Dupont. Une caractérisation des endomorphismes de Lattès par leur mesure de Green. *Comment. Math. Helv.*, 80(2):433–454, 2005.
- [2] F. Berteloot, C. Dupont, and L. Molino. Normalization of bundle holomorphic contractions and applications to dynamics. *Ann. Inst. Fourier (Grenoble)*, 58(6):2137–2168, 2008.
- [3] F. Berteloot and J.-J. Loeb. Spherical hypersurfaces and Lattès rational maps. *J. Math. Pures Appl. (9)*, 77(7):655–666, 1998.
- [4] F. Berteloot and J.-J. Loeb. Une caractérisation géométrique des exemples de Lattès de \mathbb{P}^k . *Bull. Soc. Math. France*, 129(2):175–188, 2001.
- [5] I. Binder and L. DeMarco. Dimension of pluriharmonic measure and polynomial endomorphisms of \mathbb{C}^n . *Int. Math. Res. Not.*, 2003(11):613–625, 2003.
- [6] J.-Y. Briend. *Exposants de Liapounoff et points périodiques d’endomorphismes holomorphes de $CP(k)$* . PhD thesis, Université Paul Sabatier de Toulouse, 1997.
- [7] J.-Y. Briend and J. Duval. Exposants de Liapounoff et distribution des points périodiques d’un endomorphisme de $\mathbb{C}P^k$. *Acta Math.*, 182(2):143–157, 1999.
- [8] J.-Y. Briend and J. Duval. Deux caractérisations de la mesure d’équilibre d’un endomorphisme de $\mathbb{P}^k(\mathbb{C})$. *Publ. Math. Inst. Hautes Études Sci.*, 2001(93):145–159, 2001.
- [9] M. Brin and A. Katok. On local entropy. In *Geometric dynamics (Rio de Janeiro, 1981)*, volume 1007 of *Lecture Notes in Math.*, pages 30–38. Springer, Berlin, 1983.
- [10] H. de Thélin. Minoration de la dimension de hausdorff du courant de Green. <https://arxiv.org/abs/1709.01356>.
- [11] H. de Thélin. Un phénomène de concentration de genre. *Math. Ann.*, 332(3):483–498, 2005.
- [12] H. de Thélin. Sur les exposants de Lyapounov des applications méromorphes. *Invent. Math.*, 172(1):89–116, 2008.
- [13] H. de Thélin and G. Vigny. On the measures of large entropy on a positive closed current. *Math. Z.*, 280(3-4):919–944, 2015.
- [14] J.-P. Demailly. *Complex Analytic and Differential Geometry*. 2012. available at <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [15] T.-C. Dinh. Sur les applications de Lattès de \mathbb{P}^k . *J. Math. Pures Appl. (9)*, 80(6):577–592, 2001.

- [16] T.-C. Dinh. Attracting current and equilibrium measure for attractors on \mathbb{P}^k . *J. Geom. Anal.*, 17(2):227–244, 2007.
- [17] T.-C. Dinh and C. Dupont. Dimension de la mesure d'équilibre d'applications méromorphes. *J. Geom. Anal.*, 14(4):613–627, 2004.
- [18] T.-C. Dinh and N. Sibony. *Dynamics in several complex variables: endomorphisms of projective spaces and polynomial-like mappings*, volume 1998 of *Lecture Notes in Math.*, pages 165–294. Springer, Berlin, 2010.
- [19] R. Dujardin. Fatou directions along the Julia set for endomorphisms of $\mathbb{C}\mathbb{P}^k$. *J. Math. Pures Appl. (9)*, 98(6):591–615, 2012.
- [20] C. Dupont. On the dimension of invariant measures of endomorphisms of $\mathbb{C}\mathbb{P}(k)$. *Math. Ann.*, 349(3):509–528, 2011.
- [21] C. Dupont. Large entropy measures for endomorphisms of $\mathbb{C}\mathbb{P}^k$. *Israel J. Math.*, 192(2):505–533, 2012.
- [22] C. Dupont and J. Taflin. Dynamics of fibered endomorphisms of \mathbb{P}^k . Article in preparation.
- [23] J.E. Fornæss and N. Sibony. Some open problems in higher dimensional complex analysis and complex dynamics. *Publ. Mat.*, 45(2):529–547, 2001.
- [24] M. Jonsson and D. Varolin. Stable manifolds of holomorphic diffeomorphisms. *Invent. Math.*, 149(2):409–430, 2002.
- [25] R. Mañé. The Hausdorff dimension of invariant probabilities of rational maps. In *Dynamical systems, Valparaiso 1986*, volume 1331 of *Lecture Notes in Math.*, pages 86–117. Springer, Berlin, 1988.
- [26] Y. B. Pesin. *Dimension theory in dynamical systems*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1997. Contemporary views and applications.
- [27] N. Sibony. Dynamique des applications rationnelles de \mathbf{P}^k . In *Dynamique et géométrie complexes (Lyon, 1997)*, volume 8 of *Panor. Synthèses*, pages ix–x, xi–xii, 97–185. Soc. Math. France, Paris, 1999.
- [28] P. Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [29] L.-S. Young. Dimension, entropy and Lyapunov exponents. *Ergodic Theory Dynamical Systems*, 2(1):109–124, 1982.

C. Dupont and A. Rogue
Univ Rennes
CNRS, IRMAR - UMR 6625
F-35000 Rennes, France
christophe.dupont@univ-rennes1.fr
axel.rogue@univ-rennes1.fr