

Diffusion dynamics of classical systems driven by an oscillatory force

F. Castella⁽¹⁾, P. Degond⁽²⁾, Th. Goudon⁽³⁾

(1) IRMAR, Université de Rennes 1
Campus de Beaulieu, 35042 Rennes Cedex, France
francois.castella@univ-rennes1.fr

(2) MIP, UMR 5640 (CNRS-UPS-INSA)
Université Paul Sabatier
118, route de Narbonne, 31062 Toulouse Cedex, France
degond@mip.ups-tlse.fr

(3) Team SIMPAF–INRIA Futurs & Labo. Paul Painlevé, UMR 8524
Université des Sciences et Technologies Lille 1
F-59655 Villeneuve d’Ascq Cedex, France
thierry.goudon@math.univ-lille1.fr

J. Stat. Phys., Vol. 124, N. 2-4, pp. 913-950 (2006).

Abstract

We investigate the asymptotic behavior of solutions to a kinetic equation describing the evolution of particles subject to the sum of a fixed, confining, Hamiltonian, and a small time-oscillating perturbation. Additionally, the equation involves an interaction operator which projects the distribution function onto functions of the fixed Hamiltonian. The paper aims at providing a classical counterpart to the derivation of rate equations from the atomic Bloch equations. Here, the homogenization procedure leads to a diffusion equation in the energy variable. The presence of the interaction operator regularizes the limit process and leads to finite diffusion coefficients.

Key words: Kinetic equation, Homogenization, Diffusion limit.

AMS Subject classification: 74Q10, 35Q99, 35B25, 82C70

1 Setting of the problem

We consider a particle system described by its phase-space density, or distribution function, $f(t, x, p)$: $x \in \mathbb{R}^d$ is the position variable, $p \in \mathbb{R}^d$ is the momentum, and t is the time. In practice, $d = 1, 2$ or 3 . It is convenient to also introduce the phase space variable $X = (x, p) \in \mathbb{R}^{2d}$. The evolution of the density f is governed by a kinetic equation of the form

$$\partial_t f + \{H, f\} = \frac{1}{\tau} Q(f). \quad (1.1)$$

Given the Hamiltonian of the system $H = H(t, X) = H(t, x, p)$, the Poisson bracket $\{H, f\}$ denotes the operator

$$\{H, f\} = \nabla_p H \cdot \nabla_x f - \nabla_x H \cdot \nabla_p f.$$

The left-hand side of (1.1) describes the total time derivative of f along the trajectories of the particles, i.e.

$$\frac{d}{dt} f(t, \bar{x}(t), \bar{p}(t)) = \left(\partial_t f + \{H, f\} \right) (t, \bar{x}(t), \bar{p}(t)),$$

where $\bar{X}(t) = (\bar{x}(t), \bar{p}(t))$ is any solution of the characteristic system

$$\frac{d}{dt} \bar{x}(t) = \nabla_p H(t, \bar{x}(t), \bar{p}(t)), \quad \frac{d}{dt} \bar{p}(t) = -\nabla_x H(t, \bar{x}(t), \bar{p}(t)).$$

Then, (1.1) translates the fact that the time variations of f produced by transport along the Hamiltonian flow of H balances the rate of change of f . The latter is due to complex interaction phenomena, the description of which is embodied into the operator Q (see below). The parameter $\tau > 0$ in (1.1) then appears as a relaxation time.

We are interested in a situation in which the Hamiltonian H splits into an unperturbed time-independent Hamiltonian $H_0(x, p)$, and a time dependent potential perturbation $\mathcal{V}(t, x)$, i.e.

$$H(t, x, p) = H_0(x, p) + \mathcal{V}(t, x).$$

The technical requirements on H_0 and \mathcal{V} will be specified later on. A typical example is that of a classical particle in an unperturbed potential $V_0(x)$ which leads to

$$H_0(X) = \frac{p^2}{2} + V_0(x).$$

The prototype situation is the case where H_0 is the harmonic oscillator

$$H_0(X) = \frac{p^2 + x^2}{2} = H_{\text{harm}}(X).$$

This situation is presented in detail in Appendix E.

Besides, we assume that the potential \mathcal{V} is small but has very fast time variations. Precisely, let us denote by ε the ratio between the order of magnitude of the perturbation to that of

the free Hamiltonian. We also have to define the observation time scale T , in comparison to both the typical time scale of the perturbation θ and the relaxation time τ . It turns out that the perturbation is still negligible when looking at too short time scales (say of order $\mathcal{O}(1/\varepsilon)$). This is reminiscent of the well established fact that perturbations of size ε in an integrable Hamiltonian dynamics enter at second order only: they induce an effect of typical size $\mathcal{O}(\varepsilon^2)$. In this paper, the ‘‘integrability’’ assumption is played by Hypothesis 1.2 below. For that reason, we define the time scale so that $T/\theta = 1/\varepsilon^2$, $T/\tau = \gamma/\varepsilon^2$, with $\gamma > 0$ a fixed dimensionless parameter. Accordingly, the Hamiltonian can be recast in dimensionless form as

$$H(t, x, p) = H_0(x, p) + \varepsilon V(t/\varepsilon^2, x)$$

and we wish to perform the asymptotic analysis $\varepsilon \rightarrow 0$ in the following scaled version of (1.1)

$$\varepsilon^2 \partial_t f^\varepsilon + \{H_0, f^\varepsilon\} + \varepsilon \{V(t/\varepsilon^2, x), f^\varepsilon\} = \gamma Q(f^\varepsilon). \quad (1.2)$$

The derivation of (1.2) from (1.1) is detailed in Appendix B. Such a scaling is known under the name of weak-coupling regime, and is a well-identified regime both in quantum mechanics and in classical Hamiltonian systems, see [Sp].

The present situation is the standard setting for the description of an atom which interacts with a light field. In that case, the unperturbed Hamiltonian H_0 is the atomic Hamiltonian, and the perturbation $\mathcal{V}(t, x) = \varepsilon V(t/\varepsilon^2, x)$ is the potential energy induced by the light wave in the vicinity of the atom. If a quantum mechanical setting is retained instead of a classical one, the kinetic equation (1.1) must be replaced by the quantum Liouville equation, which, for atoms, is often referred to as the atomic Bloch equation. It reads

$$i\varepsilon^2 \partial_t \rho^\varepsilon(t) = [H_0, \rho^\varepsilon(t)] + \varepsilon [V(t/\varepsilon^2), \rho^\varepsilon(t)] + \gamma Q(\rho^\varepsilon(t)), \quad (1.3)$$

where the unknown now is a time dependent trace class operator $\rho^\varepsilon(t)$, the so-called density matrix of the quantum mechanical system, and all Poisson brackets $\{\cdot, \cdot\}$ are formally replaced by commutators $[\cdot, \cdot]$ between operators, in the passage from the kinetic equation (1.2) to the quantum equation (1.3). Also, in (1.3), $Q(\rho^\varepsilon)$ is a relaxation operator that describes, at a heuristic level, the observed trend of various atomic systems to relax towards equilibrium states of the unperturbed Hamiltonian H_0 . We do not give the precise expression of $Q(\rho^\varepsilon)$ here, and refer e.g. to [Lo] for a physical discussion.

Let us now turn to the definition of the operator Q that is relevant in our context. Our basic approach follows the analogy between the quantum mechanical situation (1.3) and the associated classical setting (1.2). For quantum mechanical systems, the large time behavior of the system can be described by a time-differential system of rate equations, which describes the evolution of the populations of the atomic energy levels (see e.g. [Lo] and references therein). The rate constants depend on the frequency of the light field and the differences between the atomic energy levels (transition energies). They are large when a resonance occurs i.e. when the frequency of the light field matches one (or more) of the transition energies. These facts have been recently proved on a rigorous basis in [BCD, BCDG], starting from equation (1.3)

and performing both a density matrix analysis in the spirit of [CD, Ca1, Ca2], and an averaging procedure for Ordinary Differential Equations in the spirit of [SV]. In the present work, we would like to explore a similar situation with a classical system. The classical counterpart of the level population is the number of particles on a given energy surface. Hence, we shall assume that this number is well defined and finite for almost all energies. For that purpose, let us introduce the following requirements on the free Hamiltonian H_0 .

Hypothesis 1.1 *We assume that*

$$H_0(X) \in C^\infty(\mathbb{R}^{2d}), \quad H_0(X) \geq -C_0 \quad \text{for some } C_0 \geq 0, \quad \lim_{|X| \rightarrow \infty} H_0(X) = +\infty.$$

Hypothesis 1.2 (Well defined energy levels, having finite measure) *We assume that*

(i) *For almost all $E \in \mathbb{R}$, the set¹*

$$S_E = \{X = (x, p) \in \mathbb{R}^{2d} \mid H_0(X) = E\},$$

is a smooth orientable $2d - 1$ submanifold of \mathbb{R}^{2d} . For any such E , we let $d\sigma_E(X)$ denote the induced euclidean surface measure, and we define the measure $\delta(H_0(X) - E)$ as

$$\delta(H_0(X) - E) := \frac{d\sigma_E(X)}{|\nabla_X H_0(X)|}.$$

(ii) *For any E as in (i), S_E also has **finite measure** with respect to $\delta(H_0(X) - E)$. In other words*

$$h_0(E) := \int_{S_E} \delta(H_0(X) - E) < +\infty, \quad \text{a.e. } E \in \mathbb{R}.$$

This serves as a definition for $h_0(E)$.

Hypothesis 1.3 *Let $\bar{X} : s \in \mathbb{R} \mapsto \bar{X}(s) \in \mathbb{R}^{2d}$ stand for the solution of the ODE system*

$$\frac{d}{ds} \bar{X}(s) = (\nabla_p H_0, -\nabla_x H_0)(\bar{X}(s)), \quad \bar{X}(0) = (x, p).$$

Then we assume that the matrix of the derivatives with respect to the initial data is such that for any $0 < R < \infty$, there exist $C_R, q_R \geq 0$ verifying

$$\sup_{|(x,p)| \leq R} |\nabla_{x,p} \bar{X}(s)| \leq C_R (1 + |s|)^{q_R}$$

for any $s \in \mathbb{R}$.

¹We should write here $E \in H_0(\mathbb{R}^{2d})$ instead of $E \in \mathbb{R}$ to be rigorous. Since the distinction between $H_0(\mathbb{R}^{2d})$ and \mathbb{R} is anyhow obvious – there is nothing to assume for energies $E \notin H_0(\mathbb{R}^{2d})$ – we shall systematically consider energies $E \in \mathbb{R}$ in this article, meaning implicitly that energies should actually satisfy the rigorous condition $E \in H_0(\mathbb{R}^{2d})$.

Remark 1.1 *Of course, these assumptions are fulfilled by the harmonic potential H_{harm} . Then, the energy shells reduce to spheres $\{X \in \mathbb{R}^{2d}, X^2 = 2E\}$ and Hypothesis 1.3 simply holds with $C_R = 1, q_R = 0$. Moreover, one may take any smooth diffeomorphism of phase-space $\Phi : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$. Clearly, the new Hamiltonian $H_0(X) = H_{\text{harm}}(\Phi(X))$ also satisfies these Hypotheses. Then, energy shells are deformed spheres.*

Remark 1.2 *Hypothesis 1.1 is essentially a confining condition. As discussed in Appendix A, once H_0 is assumed C^∞ , Sard's Theorem together with the coarea formula imply that S_E is indeed a smooth codimension one submanifold, for almost every $E \in \mathbb{R}$. Hence part (i) of Hypothesis 1.2 is indeed a consequence of Hypothesis 1.1. The important point in Hypothesis 1.2 is part (ii). It can be seen as an additional growth condition on H_0 with respect to the space variable. It allows us to normalize the measure $\delta(H_0(x) - E)$. This is a key assumption in the present paper, both from the point of view of the model (it allows us to define the operator Q), and of the techniques: through Jensen's inequality, it gives us the desired "entropy estimates" suited for our asymptotic analysis. Note that the measure $\delta(H_0(x) - E)$ is a standard object in statistical physics: it is known as the microcanonical measure on the energy shell $S_E = \{H_0(X) = E\}$. It is also referred to as the Liouville measure, which is the unique invariant measure under the Hamiltonian flow generated by H_0 .*

Also, Hypothesis 1.3 is a strong stability assumption on the unperturbed potential V_0 . Its role will appear clear in Section 4.2, and is related to the regularity of the solutions of certain profile equations. Note that this Hypothesis can be relaxed, but at the price of restricting the relaxation parameter γ to large enough values.

Associated with $\delta(H_0(X) - E)$, the following mean-value operator is defined:

$$\Pi f(t, E) := \frac{1}{h_0(E)} \int_{S_E} f(t, X) \delta(H_0(X) - E) = \frac{\int_{S_E} f(t, X) \delta(H_0(X) - E)}{\int_{S_E} \delta(H_0(X) - E)}. \quad (1.4)$$

For each energy level E , Πf defines the average of f over the energy shell $\{X \mid H_0(X) = E\}$. In Appendix A, we check that $\Pi f(t, E)$ is well-defined for functions f belonging to the spaces $L^p(\mathbb{R}^{2d})$. Physically, $\Pi f(t, E)$ denotes the mean number of particles which belong to the energy shell S_E at time t . Now, the classical counterpart of the level populations being the number of particles on a given energy surface, it is natural to define the following operator

$$P : f \longmapsto Pf(t, X) := \Pi f(t, H_0(X)). \quad (1.5)$$

We shall see that P enjoys the natural self-adjointness and contraction properties of a projection: it is the projection onto functions depending only on the energy. Going on with the analogy between classical and quantum mechanics, we also observe that the classical counterpart of the density-matrix correlations is the projection of the distribution function onto the space

orthogonal to functions of the energy only. This leads us to the following definition of the relaxation operator to be used in (1.2):

$$Q(f) := Pf - f. \tag{1.6}$$

This operator models the relaxation of the distribution function towards a function of the total energy of the system only. Physically, it describes a redistribution of the particles which makes the distribution uniform on any energy shell. To motivate this interaction, we can think of some resonant interaction process: two particles with different energies do not spend enough time in a coherent motion one with respect to each other to interact significantly. Only particles which have the same energy do interact, and if this interaction is repulsive, it eventually produces a uniform distribution on the energy shell. Further considerations on how such a relaxation operator can be derived are beyond the scope of this work.

Let us give some intuition of the phenomena involved in (1.2), endowed with the operator (1.6). First, as $\varepsilon \rightarrow 0$, we can expect that f^ε relaxes towards an equidistributed repartition i.e. towards a solution to $Pf = f$. However, the fluctuations $f^\varepsilon - Pf^\varepsilon$, which are small but definitely non zero, are transported by the Hamiltonian flow. Then, resonant interactions are possible with the motion induced by the perturbation εV which oscillates with frequency $1/\varepsilon^2$. These intricate interactions will eventually give rise to diffusion in the energy variable. Of course, the asymptotics is highly governed by the precise time dependence of V . It turns out that the relaxation operator Q somewhat regularizes the situation in this respect, in that it prevents the possibility of too strong resonances (small denominators), through the introduction of some damping in the model. Let us comment further the introduction of this operator:

- On the one hand, as explained above, the situation has to be compared with the quantum Bloch equation (1.3), which has been analyzed in [BCD] and further in [BCDG]. There, the term $Q(\rho^\varepsilon)$ gives damping terms for the off-diagonal elements of the density matrix (the correlations, analogous to $f^\varepsilon - Pf^\varepsilon$ here). These damping terms make the large-time dynamics dominated by the diagonal elements (the populations, analogous to Pf^ε here). They also contribute to making the rate constants finite even at resonances (the “width” of the resonance being related to the damping rates). These damping terms can be physically motivated in a number of ways (for instance they can model the decoherence effects of atomic collisions in a gaseous medium, see the discussion in [Lo]). Under more restrictive assumptions on the data, smaller damping rates of order $\mathcal{O}(\varepsilon^\mu)$ with $\mu < 1/2$ could be considered and the usual (undamped) formulae for the Einstein rate equations [Lo] could be recovered, see [BCD, BCDG].

- On the other hand, the operator Q introduces non reversibility in the system through dissipation mechanisms. Without damping rates, the Bloch equation is time-reversible while the rate equations are time-irreversible. The damping terms in the quantum Liouville equation make it an irreversible equation from the beginning and simplifies the mathematical theory. A similar idea was used in [CD, Ca1, Ca2] for the derivation of the Pauli master equation from the quantum Liouville equation in a deterministic framework. Indeed, it is a well-known fact, since the work of Lanford [La] about the derivation of the Boltzmann equation, that rigorously passing from a reversible to an irreversible dynamics is extremely difficult. A second, probably more standard, approach to overcome this problem is the introduction of stochastic averaging

in the model, as in [EY1], [EY2], [PV], [LV] (see also [KPR] in a different context). There are several other examples of such an alternative: homogenization of convection(-diffusion) equations (see [GP2, GP3] and references therein), Lorentz gas evolving in a billiard (see [BDG] and [BCS]), quantum scattering limit of the Schrödinger equation [BPR], [EY2], [PR], [PV]... For the (space-)homogenization of the kinetic equation without dissipative term, we refer e.g. to [Ale2], [FH]. Here, as well as in [BCD] and [BCDG], we wish to treat the problem in a fully deterministic framework. To some extent, in this framework, the damping term plays the same role as the stochastic averaging process (see remark 3.2 below).

We wish to add a last comment. In the quantum context, it has been proved (see [BCD] and [BCDG] for extensions) that the asymptotic behavior of the Bloch equations (1.3) leads to an Ordinary Differential System (the system of rate equations) describing the occupation numbers of the various energy levels. This system describes the jump process of the electrons between the energy levels. However, in contrast with the quantum case where the energy levels are naturally discrete (like the lowest energy levels of an atom), a classical system possesses a continuum of allowed energies and the corresponding transition energies are infinitesimally small. Therefore, the large time evolution of a classical system (or equivalently, in our framework, the $\varepsilon \rightarrow 0$ limit of eqs. (1.2), (1.6)) is expected to take place through infinitesimal energy jumps, i.e. through a diffusion process in energy, rather than through a finite jump process. For this reason, the limit model will be in the form of a diffusion equation in the energy variable, or in other words, of a Fokker-Planck type equation. The goal of the paper is to rigorously show this fact and to obtain the classical mechanics counterparts of the results proved in [BCD]. The main result of this work can be summarized as follows.

Formal statement. *We suppose that V oscillates quasi-periodically: $V(\tau, x) = V_q(\omega\tau, x)$, where $\omega \in \mathbb{R}^r$ has rationally independent components and $\theta \mapsto V_q(\theta, x)$ is $(0, 1)^r$ -periodic. Then, up to some “reasonable” assumptions on V_q , $f^\varepsilon(t, X)$ converges to some $F(t, H_0(X))$, where $F(t, E)$ satisfies a diffusion equation, which can be written in the following conservative form*

$$\partial_t(h_0F) - \partial_E(h_0b \partial_E F) = 0, \tag{1.7}$$

with h_0 defined in Hypothesis 1.2. The coefficient $b(E) \geq 0$ is defined by an expression involving some average of V_q .

The expression of the effective coefficient b , as well as the precise notion of convergence will be stated later on (see Section 3). In (1.7), $h_0F(E) dE$ can be interpreted as the number of particles having their energies in the interval $(E, E + dE)$ while $h_0b \partial_E F(E)$ gives the particle flux through the energy surface S_E .

The remainder of this paper is organized as follows. Section 2 is devoted to the basic properties of both the relaxation and transport operators, which will be crucial for our analysis. In Section 3, we provide a formal derivation of the asymptotic model. To this aim, we restrict ourselves to the framework of quasi-periodic perturbation potentials V . In this framework, we are able to give the precise and complete statement of our convergence result. This discussion allows us to

point out the mathematical difficulties related to the resolution of adequate profile equations. These difficulties are analyzed in Section 4. Next, details of the convergence proof are presented in Section 5. We postpone the proofs of several technical facts – which could be interesting in themselves – to the Appendix.

2 Preliminary considerations: properties of the relaxation operator

Since equations (1.2), (1.6) describe a relaxation phenomenon, we are naturally led to investigate the dissipation properties of the operator Q . This will give a particular form of the “entropy dissipation estimates” that are suited to our problem. Also, the commutator between both operators $f \mapsto Pf$ and $f \mapsto \{H_0, f\}$ is an important object in the asymptotic analysis of (1.2). Hence, the following statement will be useful.

Lemma 2.1 *The operator P satisfies the following properties:*

(i) P is a continuous projection operator on L^p spaces:

$$P(Pf) = Pf, \quad \|Pf\|_{L^p(\mathbb{R}^{2d})} \leq \|f\|_{L^p(\mathbb{R}^{2d})} \quad 1 \leq p \leq \infty.$$

(ii) P is conservative in the sense that for any integrable function, we get

$$\int_{\mathbb{R}^{2d}} Pf \, dX = \int_{\mathbb{R}^{2d}} f \, dX.$$

(iii) P is self-adjoint with respect to the inner product in $L^2(\mathbb{R}^{2d})$ (denoted by $\langle \cdot, \cdot \rangle$ throughout the paper). Consequently, the following orthogonality property holds: for any function $f \in L^2(\mathbb{R}^{2d})$ and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ such that $X \mapsto \varphi(H_0(X))$ lies in $L^2(\mathbb{R}^{2d})$, we have

$$\langle \varphi(H_0(X)), (\text{Id} - P)f \rangle = 0.$$

(iv) P is a non negative operator: if $f \geq 0$ almost everywhere (a.e.), then $Pf \geq 0$ a.e. as well. Moreover, the stronger property holds:

$$\text{If } f \geq 0 \text{ a.e., and } Pf = 0 \text{ a.e., then } f = 0 \text{ a.e.}$$

(v) The operators $f \mapsto Pf$ and $f \mapsto \{H_0, f\}$ are orthogonal, in the sense that

$$P\{H_0, f\} = 0,$$

holds for any $f \in L^2(\mathbb{R}^{2d})$ such that $\{H_0, f\} \in L^2(\mathbb{R}^{2d})$. Consequently, for any $f, g \in L^2(\mathbb{R}^{2d})$ such that $\{H_0, f\}$ and $\{H_0, g\}$ in $L^2(\mathbb{R}^{2d})$, we have

$$P(\{H_0, f\}g) = -P(f\{H_0, g\}).$$

Property (iii) implies that

$$\int_{\mathbb{R}^{2d}} (Pf - f)Pf \, dX = 0.$$

Therefore, we deduce the following key property of the relaxation operator.

Corollary 2.2 *The operator Q is a bounded operator on $L^2(\mathbb{R}^{2d})$ and the relation*

$$-\int_{\mathbb{R}^{2d}} Q(f)f \, dX = \int_{\mathbb{R}^{2d}} |Pf - f|^2 \, dX \geq 0$$

holds for any $f \in L^2(\mathbb{R}^{2d})$.

Proof of Lemma 2.1. We split the proof as follows.

Proof of (i)-(ii)-(iii)

The continuity of P on L^p spaces is an immediate consequence of the coarea formula recalled in Appendix A, together with the assumption that S_E has finite measure for $E \in \mathbb{R}$ a.e. Indeed,

$$\begin{aligned} \|Pf\|_{L^p(dX)}^p &= \int_{\mathbb{R}^{2d}} |\Pi f(H_0(X))|^p \, dX \\ &= \int_{\mathbb{R}} |\Pi f(E)|^p h_0(E) \, dE \\ &\leq \int_{\mathbb{R}} \left(\int_{S_E} |f(X)|^p \frac{\delta(H_0(X) - E)}{h_0(E)} \right) h_0(E) \, dE \\ &\leq \int_{\mathbb{R}^{2d}} |f(X)|^p \, dX \end{aligned}$$

where the coarea formula (A.5) is used for the second equality, Jensen's inequality for the first inequality and the coarea formula again for the second inequality. Note that equality holds for $p = 1$. The relation $P(Pf) = Pf$ is obvious since P leaves any function depending only on $H_0(X)$ invariant. Finally, the self-adjointness of P simply comes from the identity $P = \Pi^*\Pi$, where Π^* is the adjoint of Π (with the notations of the Appendix – see Lemma A.1).

Proof of (iv)

It is obvious that P preserves non negativity. Let $f \geq 0$ such that $Pf = 0$ a.e.. Since $\int_{\mathbb{R}^{2d}} f \, dX = \int_{\mathbb{R}^{2d}} Pf \, dX = 0$, then, f is a nonnegative function with vanishing integral, which implies that $f(X) = 0$ for $X \in \mathbb{R}^{2d}$ a.e.

Proof of (v)

We deduce that $P\{H_0, f\} = 0$ from $\Pi\{H_0, f\} = 0$. To prove the latter, we take any test function $\psi(E) \in L^2(\mathbb{R}, h_0(E) \, dE)$. We write

$$\begin{aligned} \langle \Pi\{H_0, f\}, \psi \rangle_{L^2(\mathbb{R}; h_0(E) \, dE)} &= \int_{\mathbb{R}} \Pi\{H_0, f\}(E) \psi(E) h_0(E) \, dE \\ &= \langle \{H_0, f\}, \Pi^*\psi \rangle_{L^2(\mathbb{R}^{2d})} \\ &= \langle \{H_0, f\}, \psi(H_0(X)) \rangle_{L^2(\mathbb{R}^{2d})} \\ &= -\langle f, \{H_0, \psi(H_0(X))\} \rangle_{L^2(\mathbb{R}^{2d})} \\ &= 0. \end{aligned}$$

where the definition of Π^* can be found in Lemma A.1 of the Appendix and where we have used an integration by parts to obtain the fourth equality. Then, combining this property together with the Leibniz rule $\{H_0, fg\} = \{H_0, f\}g + f\{H_0, g\}$ allows to conclude the proof. \blacksquare

3 Formal derivation; quasi-periodicity

We consider a perturbation V which oscillates in a quasi-periodic way. To be more precise, let \mathbb{Y} be the unit cube in \mathbb{R}^r , for some integer $r \geq 1$. We assume the following

Quasi-periodicity Hypothesis: *There exists a vector $\omega \in \mathbb{R}^r \setminus \{0\}$ and a smooth and bounded function $V_q : \mathbb{R}^r \times \mathbb{R}^d \rightarrow \mathbb{R}$, which is \mathbb{Y} -periodic with respect to its first variable, such that*

$$V(\tau, x) = V_q(\omega\tau, x), \quad \text{for any } \tau \in \mathbb{R}, x \in \mathbb{R}^d.$$

The periodicity condition means that $V_q(\theta + j, x) = V_q(\theta, x)$ holds for any $\theta \in \mathbb{Y}$, $x \in \mathbb{R}^d$, $j \in \mathbb{N}^r$. The vector ω is called the frequency vector. It collects the r frequencies of V . We assume that the r components of ω are rationally independent, which means that $k \cdot \omega = 0$, for $k \in \mathbb{Q}^r$ iff $k = 0$. When $r = 1$, V is simply said to be periodic, and one can take $\omega = 1$ without loss of generality. It will be convenient later to make use of the Fourier series associated to V_q

$$V_q(\theta, x) = \sum_{k \in \mathbb{Z}^r} \widehat{V}_q(k, x) \exp(2i\pi k \cdot \theta), \quad \widehat{V}_q(k, x) = \int_{\mathbb{Y}} V_q(\theta, x) \exp(-2i\pi k \cdot \theta) d\theta.$$

Provided V_q has the smoothness $V_q(\theta, x) \in L^2(\mathbb{Y} \times \mathbb{R}^d)$, the above Fourier series is convergent in the topology $\ell^2(\mathbb{Z}^r; L^2(\mathbb{R}^d))$ (note that we shall need the stronger regularity $V_q \in C_b^2$, see assumption 3.1 below).

With the help of this assumption, we can now guess the behavior of f^ε by inserting into eq. (1.2) a double scale ansatz in the spirit of [BLP], [SP]:

$$f^\varepsilon(t, X) = f_q^{(0)}(t, \omega t/\varepsilon^2, X) + \varepsilon f_q^{(1)}(t, \omega t/\varepsilon^2, X) + \varepsilon^2 f_q^{(2)}(t, \omega t/\varepsilon^2, X) + \dots$$

where all functions $f_q^{(i)}$ are supposed \mathbb{Y} -periodic with respect to the second variable. Then, we formally identify all terms which appear with the same power of ε . Remarking that

$$\partial_t \left(f_q^{(i)}(t, \omega t/\varepsilon^2, X) \right) = \left(\partial_t f_q^{(i)} + \frac{1}{\varepsilon^2} \omega \cdot \nabla_\theta f_q^{(i)} \right) (t, \omega t/\varepsilon^2, X),$$

it becomes convenient to introduce the operator

$$\mathcal{T} f_q = \omega \cdot \nabla_\theta f_q + \{H_0, f_q\} - \gamma Q(f_q),$$

and its formal adjoint $\mathcal{T}^*\varphi = -\omega \cdot \nabla_{\theta}\varphi - \{H_0, \varphi\} - \gamma Q(\varphi)$. We obtain the following profile equations

$$\varepsilon^0 \text{ term:} \quad \mathcal{T}f_q^{(0)} = 0, \quad (3.1)$$

$$\varepsilon^1 \text{ term:} \quad \mathcal{T}f_q^{(1)} = \nabla_x V_q(\theta, x) \cdot \nabla_p f_q^{(0)}, \quad (3.2)$$

$$\varepsilon^2 \text{ term:} \quad \mathcal{T}f_q^{(2)} = -\partial_t f_q^{(0)} + \nabla_x V_q(\theta, x) \cdot \nabla_p f_q^{(1)} \quad (3.3)$$

and so on. The general form of these equation reads $\mathcal{T}f_q = h_q$, and the time variable t appears only as a parameter. As a matter of fact, we readily check that any function depending only on the energy variable, but not on θ , belongs to the kernel of \mathcal{T} . Therefore, it is tempting to infer from (3.1) that

$$f_q^{(0)}(t, \theta, X) = F(t, H_0(X)).$$

Since such a function also lies in the kernel of the adjoint operator \mathcal{T}^* , we might imagine that the orthogonality relation

$$\int_{\mathbb{Y}} P h_q d\theta = 0$$

can serve as a compatibility condition. Assuming that these considerations hold true, and forgetting for the time being any functional difficulties, we rewrite (3.2) as

$$\mathcal{T}f_q^{(1)} = \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \partial_E F(t, H_0(X)).$$

Note that $\nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) = -\{V_q, H_0\}$ fulfils the compatibility condition, thanks to Lemma 2.1-(v). Thus, we can define $\chi_q(\theta, X)$, a solution of the auxiliary equation

$$\mathcal{T}\chi_q = \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X),$$

and we set $f_q^{(1)}(t, \theta, X) = \chi_q(\theta, X) \partial_E F(t, H_0(X))$. Inserting this expression into the ε^2 order equation (3.3), and using the compatibility condition, we are led to

$$\begin{aligned} 0 &= \partial_t P\left(F(t, H_0(X))\right) - \int_{\mathbb{Y}} P\left(\nabla_x V_q(\theta, x) \cdot \nabla_p f_q^{(1)}(t, \theta, X)\right) d\theta \\ &= \partial_t F(t, H_0(X)) - \left(\int_{\mathbb{Y}} P\left(\nabla_x V_q(\theta, x) \cdot \nabla_p \chi_q(\theta, X)\right) d\theta\right) \partial_E F(t, H_0(X)) \\ &\quad - \left(\int_{\mathbb{Y}} P\left(\nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \chi_q(\theta, X)\right) d\theta\right) \partial_{EE}^2 F(t, H_0(X)). \end{aligned}$$

Thanks to the coarea formula (A.5), we deduce that $F(t, E)$ verifies the following drift-diffusion equation

$$\partial_t (h_0(E)F(t, E)) = h_0(E)a(E)\partial_E F(t, E) + h_0(E)b(E)\partial_{EE}^2 F(t, E), \quad (3.4)$$

the coefficients of which are defined by

$$\begin{cases} a(E) = \Pi\left(\int_{\mathbb{Y}} \nabla_x V_q(\theta, x) \cdot \nabla_p \chi_q(\theta, X) d\theta\right)(E), \\ b(E) = \Pi\left(\int_{\mathbb{Y}} \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \chi_q(\theta, X) d\theta\right)(E). \end{cases}$$

For further purposes, it is also convenient to introduce χ_q^* , a solution of the adjoint profile equation

$$\mathcal{T}^* \chi_q^* = \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X).$$

This function is precisely defined in Corollary 4.4 below. Let us set

$$\begin{cases} a^*(E) = \Pi \left(\int_{\mathbb{Y}} \nabla_x V_q(\theta, x) \cdot \nabla_p \chi_q^*(\theta, X) d\theta \right) (E) \\ b^*(E) = \Pi \left(\int_{\mathbb{Y}} \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \chi_q^*(\theta, X) d\theta \right) (E). \end{cases} \quad (3.5)$$

The following claim will make the connection with (1.7) clear.

Lemma 3.1 *The following relations hold true:*

$$h_0(E)b^*(E) = h_0(E)b(E), \quad h_0(E)a^*(E) = h_0(E)a(E) = \frac{d}{dE} \left(h_0(E)b^*(E) \right).$$

These relations are consequences of the coarea formula; detailed computations are presented in Appendix C. Therefore, from (3.4), we are led to (1.7):

$$\partial_t(h_0 F) = \partial_E(h_0 b) \partial_E F(t, E) + h_0(E)b(E) \partial_{EE}^2 F(t, E) = \partial_E(h_0 b \partial_E F).$$

We are now left with the task of making this formal guess rigorous. To this end, we need some technical assumptions on the perturbation V .

Hypothesis 3.1 *We assume that*

(i) *the quasiperiodic potential $V(t, x) = V_q(\omega t, x)$ possesses the regularity $V_q \in C_b^2(\mathbb{Y} \times \mathbb{R}^d)$, where V_q is \mathbb{Y} -periodic with respect to the first variable.*

(ii) *There exists some $\beta \geq 0$ such that*

$$\sup_{\theta \in \mathbb{Y}} \int_{\mathbb{R}^{2d}} \frac{|\nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X)|^2}{w(X)^\beta} dX < \infty,$$

where $w(X) = (1 + H_0(X)^2)^{1/2}$.

Remark 3.1 *Considering the harmonic Hamiltonian, we get $\nabla_x V \cdot \nabla_p H_{\text{harm}}(X) p \cdot \nabla_x V$ which clearly does not belong to $L^2(\mathbb{R}^{2d})$. However, Hypothesis 3.1-(ii) holds for any $\beta > d + 1$. Thus, the ε order equation (3.2) makes sense in a reasonable functional space since the right-hand side belongs to the weighted space $L^2(\mathbb{R}^{2d}, w(X)^{-\beta} dX)$.*

We are now ready to give the statement of our main result.

Theorem 3.2 *Let $f_0^\varepsilon \geq 0$ be the initial data for (1.2). We suppose that f_0^ε is bounded in $L^2(\mathbb{R}^{2d})$. We suppose that Hypothesis 1.1, 1.2, 1.3 and 3.1 are satisfied. Then, $f^\varepsilon = P f^\varepsilon + \varepsilon g_\varepsilon$ where g_ε is bounded in $L^2((0, T) \times \mathbb{R}^{2d})$ and, up to a subsequence, $P f^\varepsilon(t, X)$ converges in $C^0([0, T]; L^2(\mathbb{R}^{2d}) - \text{weak})$ to $F(t, H_0(X))$, where $F : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}^+$ satisfies the diffusion equation (1.7) weakly in $L^2(\mathbb{R})$, with the initial data $F(t = 0, E)$ given by the weak limit of $\Pi f_0^\varepsilon(E)$ in $L^2(\mathbb{R}, h_0(E) dE)$.*

Remark 3.2 We point out that assuming $\gamma > 0$ is crucial in our analysis since the operator Q plays the role of a dissipation which allows to avoid all resonance phenomena. The explicit computations presented in Appendix E may shed some light on this aspect. Without such a relaxation, the mathematical analysis becomes very delicate and certainly does not lead to a diffusion process. We refer in particular to [Ale2], [FH] where it is shown that the homogenization of a kinetic equation with highly oscillatory force fields leads to an effective equation involving memory effects. These results are in the spirit of those concerning the homogenization of transport equations with transverse oscillations [Ale1], [AHZ], [Pe] as initiated by [T]. In the present approach, we avoid these effects thanks to the presence of a dissipation operator.

4 Profile Equations

This section is devoted to the analysis of the profile equation $\mathcal{T}f_q = h_q$. We denote by $L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$ the class of functions $f_q : \mathbb{R}^r \times \mathbb{R}^{2d} \rightarrow \mathbb{R}$ which are \mathbb{Y} -periodic with respect to the first variable and such that

$$\int_{\mathbb{Y} \times \mathbb{R}^{2d}} |f_q(\theta, X)|^2 d\theta dX < \infty.$$

We also introduce

$$H_{\#} = \left\{ f_q \in L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d}), \mathcal{T}f_q \in L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d}) \right\}.$$

4.1 General setting

Proposition 4.1 Let $h_q \in L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$. We suppose that h_q is either purely periodic or has finitely many harmonics, which means that either $r = 1$, or, when $r \geq 2$,

$$h_q(\theta, x) = \sum_{k \in \mathbb{Z}^r, |k| \leq K} \widehat{h}_q(k, x) \exp(ik \cdot \theta), \quad (4.1)$$

for some finite integer K . Then, the problem $\mathcal{T}f_q = h_q$ has a solution $f_q \in H_{\#}$ iff h_q satisfies the compatibility condition

$$\int_{\mathbb{Y}} Ph_q(\theta, X) d\theta = 0. \quad (4.2)$$

The solution is unique when imposing the additional constraint $\int_{\mathbb{Y}} Pf_q(\theta, X) d\theta = 0$. This uniquely defined solution depends continuously on h_q : there exists $C > 0$ such that

$$\|f_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \leq C \|h_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}.$$

Other solutions differ from f_q by a function $\varphi(H_0(X))$.

Proof. The arguments are inspired from [GP1], but specific difficulties appear, since in particular the operators $\omega \cdot \nabla_{\theta}$ and $\{H_0, \cdot\} - Q$ act on independent variables. As it will become clear in the proof, the restriction contained in (4.1) is related to small denominator difficulties

when solving the profile equations. These difficulties disappear in the purely periodic case. The proof splits as follows.

Uniqueness

For any $f_q \in H_\#$, we observe that

$$\int_{\mathbb{Y} \times \mathbb{R}^{2d}} \omega \cdot \nabla_\theta f_q f_q d\theta dX = 0, \quad \int_{\mathbb{Y} \times \mathbb{R}^{2d}} \{H_0, f_q\} f_q d\theta dX = 0.$$

Let $f_q \in H_\#$ be a solution of $\mathcal{T}f_q = 0$. Multiplying by f_q and integrating yields

$$-\gamma \int_{\mathbb{Y} \times \mathbb{R}^{2d}} Q(f_q) f_q d\theta dX = 0 = \gamma \|f_q - Pf_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2$$

thanks to Corollary 2.2. We deduce that $f_q(\theta, X) = Pf_q(\theta, X)$ depends on X only through the energy. Then, we apply the operator P to the equation. We get

$$\omega \cdot \nabla_\theta Pf_q = 0$$

thanks to Lemma 2.1-(ii) and (v). Accordingly, the Fourier coefficients of Pf_q verify

$$\omega \cdot k \widehat{Pf_q}(k, X) = 0.$$

Since the components of the frequency vector ω are assumed rationally independent, we deduce that $\widehat{Pf_q}(k, X) = 0$ for any $k \neq 0$, and thus this implies that $Pf_q(\theta, X)$ does not depend on the variable $\theta \in \mathbb{Y}$. We proved that $f_q \in L^2(\mathbb{Y} \times \mathbb{R}^{2d})$ verifies $\mathcal{T}f_q = 0$ iff $f_q(\theta, X) = F(H_0(X))$, for some F such that $\int_{\mathbb{R}^{2d}} |F(H_0(X))|^2 dX < \infty$. In particular, if we impose that $\int_{\mathbb{Y}} Pf_q d\theta = 0$, this implies that $f_q = 0$, proving the uniqueness result.

Existence

Applying the projector P to the equation $\mathcal{T}f_q = h_q$ and integrating over \mathbb{Y} , we realize that (4.2) is a necessary condition for having a solution. From now on, we thus assume that (4.2) holds true and we prove that it is also a sufficient condition. Let us temporarily assume that, for any $\lambda > 0$, there exists $f_q^{(\lambda)} \in H_\#$ verifying

$$\lambda f_q^{(\lambda)} + \mathcal{T}f_q^{(\lambda)} = h_q. \tag{4.3}$$

We wish to prove the existence part of Proposition 4.1 by passing to the limit $\lambda \rightarrow 0$. This is completely obvious once we know that the sequence $(f_q^{(\lambda)})_{\lambda > 0}$ remains bounded in $L^2(\mathbb{Y} \times \mathbb{R}^{2d})$.

Suppose that there exists a subsequence, say $\{\lambda^{(n)}, n \in \mathbb{N}\}$ such that $\lim_{n \rightarrow \infty} \lambda^{(n)} = 0$ and

$$N^{(n)} = \|f_q^{(\lambda^{(n)})}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \xrightarrow{n \rightarrow \infty} +\infty.$$

We set $F_q^{(n)} = f_q^{(\lambda^{(n)})}/N^{(n)}$. Without loss of generality, we can assume that $F_q^{(n)} \rightharpoonup F_q$ weakly in $L^2(\mathbb{Y} \times \mathbb{R}^{2d})$ as $n \rightarrow \infty$. We have

$$\lambda^{(n)} F_q^{(n)} + \mathcal{T}F_q^{(n)} = \frac{h_q}{N^{(n)}}.$$

Hence, multiplying by $F_q^{(n)}$ leads to

$$\gamma \|F_q^{(n)} - PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 \leq \int_{\mathbb{Y} \times \mathbb{R}^{2d}} \frac{h_q}{N^{(n)}} F_q^{(n)} d\theta dX \leq \frac{\|h_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}}{N^{(n)}}.$$

We deduce that

$$\|F_q^{(n)} - PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 \xrightarrow{n \rightarrow \infty} 0. \quad (4.4)$$

Accordingly, $F_q^{(n)} = PF_q^{(n)} + (F_q^{(n)} - PF_q^{(n)}) \rightarrow F_q = PF_q$ as $n \rightarrow \infty$. Now, we apply the projection operator and we get

$$\lambda^{(n)} PF_q^{(n)} + \omega \cdot \nabla_\theta PF_q^{(n)} = \frac{Ph_q}{N^{(n)}}. \quad (4.5)$$

Integrating with respect to θ , we obtain for any $n \in \mathbb{N}$

$$\int_{\mathbb{Y}} PF_q^{(n)}(\theta, X) d\theta = 0,$$

as a consequence of (4.2). Besides, passing to the limit in (4.5) yields

$$\omega \cdot \nabla_\theta PF_q^{(n)} \xrightarrow{n \rightarrow \infty} \omega \cdot \nabla_\theta PF_q = 0 \quad \text{strongly in } L^2(\mathbb{Y} \times \mathbb{R}^{2d}).$$

Hence the limit is nothing but $F_q = 0$. We will obtain a contradiction by showing that $F_q^{(n)}$ converges strongly.

Let us consider the Fourier series associated with $PF_q^{(n)}$

$$PF_q^{(n)}(\theta, X) = \sum_{k \in \mathbb{Z}^r} \widehat{PF_q^{(n)}}(k, X) e^{2i\pi k \cdot \theta}.$$

We have already remarked that the first Fourier coefficient vanishes

$$\widehat{PF_q^{(n)}}(0, X) = \int_{\mathbb{Y}} PF_q^{(n)}(\theta, X) d\theta = 0.$$

Therefore, the Plancherel theorem gives

$$\begin{aligned} \|PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 &= \sum_{k \in \mathbb{Z}^r \setminus \{0\}} |\widehat{PF_q^{(n)}}(k, X)|^2 \\ &= \sum_{k \in \mathbb{Z}^r \setminus \{0\}} \frac{1}{|\omega \cdot k|^2} |\omega \cdot k|^2 |\widehat{PF_q^{(n)}}(k, X)|^2 \\ &= \sum_{k \in \mathbb{Z}^r \setminus \{0\}} \frac{1}{4\pi^2 |\omega \cdot k|^2} |\omega \cdot \nabla_\theta \widehat{PF_q^{(n)}}(k, X)|^2. \end{aligned}$$

When $r \geq 2$, we use the assumption that the data h_q has finitely many harmonics. By (4.5), $PF_q^{(n)}$ shares the same property, with the same truncation index K and we are thus led to

$$\|PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 \leq \sup_{k \in \mathbb{Z}^r \setminus \{0\}, |k| \leq K} \left(\frac{1}{4\pi^2 |\omega \cdot k|^2} \right) \|\omega \cdot \nabla_\theta PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 \xrightarrow{n \rightarrow \infty} 0.$$

When $r = 1$ the conclusion is immediate since we get $\|PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 \leq \|\partial_\theta PF_q^{(n)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2$.

It remains to justify the existence of $F^{(\lambda)}$. This is obtained by a Banach fixed point argument. Indeed, consider the operator $\Phi^{(\lambda)}$, which to a function $\phi \in L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$ associates the solution $\psi^{(\lambda)} = \Phi^{(\lambda)}(\phi) \in L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$ to the transport equation

$$\lambda\psi^{(\lambda)}(\theta, X) + \omega \cdot \nabla_\theta \psi^{(\lambda)} + \{H_0, \psi^{(\lambda)}\} + \gamma\psi^{(\lambda)} = \gamma P\phi + h_q(\theta, X).$$

We prove that $\Phi^{(\lambda)}$ is a contraction over $L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$. Since (4.3) also reads $f_q^{(\lambda)} = \Phi^{(\lambda)}(f_q^{(\lambda)})$, this clearly implies the existence and uniqueness of $f_q^{(\lambda)}$, the solution to (4.3). Now, to prove the contraction property of $\Phi^{(\lambda)}$, we take two functions ϕ and $\tilde{\phi}$, with the associated $\psi^{(\lambda)} = \Phi^{(\lambda)}(\phi)$ and $\tilde{\psi}^{(\lambda)} = \Phi^{(\lambda)}(\tilde{\phi})$. We readily obtain the following energy estimate

$$\begin{aligned} (\lambda + \gamma) \|\psi^{(\lambda)} - \tilde{\psi}^{(\lambda)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}^2 &\leq \gamma \left| \langle P\phi - P\tilde{\phi}, \psi^{(\lambda)} - \tilde{\psi}^{(\lambda)} \rangle_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \right| \\ &\leq \gamma \|\phi - \tilde{\phi}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \|\psi^{(\lambda)} - \tilde{\psi}^{(\lambda)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}. \end{aligned}$$

The second estimate uses the Cauchy-Schwarz inequality together with the continuity of P over $L^2(\mathbb{Y} \times \mathbb{R}^{2d})$ (see Lemma 2.1). As a consequence, we have

$$\|\psi^{(\lambda)} - \tilde{\psi}^{(\lambda)}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \leq \frac{\gamma}{\gamma + \lambda} \|\phi - \tilde{\phi}\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}.$$

This is the claimed contraction property.

This ends the proof of Proposition 4.1. The continuity estimate follows from the closed graph theorem, once we have remarked that the set of functions verifying the compatibility condition is a closed subspace of $L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$. \blacksquare

The distinction between the purely periodic case and the genuinely quasiperiodic case is due to small denominator difficulties: while the transport operator ∂_θ is (essentially) invertible over $L^2(d\theta)$ in one dimension, the inverse of the transport operator $\omega \cdot \nabla_\theta$ ceases to be bounded in reasonable spaces when the angular variable θ belongs to the $r > 1$ dimensional torus. This appears clearly when we try to deduce the behavior of $PF^{(n)}$ from informations on $\omega \cdot \nabla_\theta PF^{(n)}$. In the periodic case the required estimate is actually nothing but the classical Poincaré-Wirtinger estimate for periodic functions on $(0, 1)$. When $r \geq 2$, the quantity $|\omega \cdot k|^2$ is never zero when $k \neq 0$, due to the rational independence of the components of ω . Nevertheless, small denominators might appear, corresponding to cases where $\omega \cdot k$ is small but nonzero. This typically happens for large values of $|k|$. This is the reason why we assume, in the case $r \geq 2$, that h_q has finitely many harmonics. Another (classical) way to analyze this difficulty consists in saying that the Fredholm alternative does not apply to the transport operator $\omega \cdot \nabla_\theta$; its range is not closed in general. The difficulty can also be illustrated by imposing some diophantine condition on ω (which is therefore satisfied for almost all ω). Some slight adaptations of the previous proof then lead to the following claim

Proposition 4.2 *Let ω satisfy the following diophantine condition: for any $k \in \mathbb{Z}^r$,*

$$|\omega \cdot k| \geq \frac{C_\gamma}{|k|^\gamma},$$

holds for some $\gamma > 0$ and $C_\gamma > 0$. Let $h_q \in L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$ satisfy

$$\|Ph_q\|_{H^{\gamma}_{\#}(\mathbb{Y}; L^2(\mathbb{R}^{2d}))}^2 \sum_{k \in \mathbb{Z}^r} |k|^{2\gamma} \|\widehat{Ph_q}(k, \cdot)\|_{L^2(\mathbb{R}^{2d})}^2 < \infty.$$

Then, the problem $\mathcal{T}f_q = h_q$ has a solution $f_q \in H_{\#}$ iff h_q satisfies the compatibility condition (4.2). The solution is unique when imposing the additional constraint $\int_{\mathbb{Y}} Pf_q(\theta, X) d\theta = 0$. This uniquely defined solution depends continuously on h_q in the sense that

$$\|(I - P)f_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \leq C \|h_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})}, \quad \|Pf_q\|_{L^2(\mathbb{Y} \times \mathbb{R}^{2d})} \leq C \|h_q\|_{H^{\gamma}(\mathbb{Y}; L^2(\mathbb{R}^{2d}))}.$$

Other solutions differ from f_q by a function $\varphi(H_0(X))$.

In the course of the formal derivation, we have seen that we actually have to consider data belonging to some weighted space:

$$h_q : \mathbb{Y} \times \mathbb{R}^{2d} \rightarrow \mathbb{R}, \quad \mathbb{Y} - \text{periodic}, \quad \int_{\mathbb{Y} \times \mathbb{R}^{2d}} |h_q(\theta, X)|^2 w(X)^{\alpha} dX d\theta < \infty$$

for some real α . The profile equation in such a weighted space is easily reduced to the simpler L^2 framework. Indeed, define $\tilde{h}_q(\theta, X) = h_q(\theta, X)w(X)^{\alpha/2}$. Then, \tilde{h}_q belongs to $L^2_{\#}(\mathbb{Y} \times \mathbb{R}^{2d})$. Hence, we solve $\mathcal{T}\tilde{f}_q = \tilde{h}_q$ with $\tilde{f}_q \in H_{\#}$, $\int_{\mathbb{Y}} P\tilde{f}_q d\theta = 0$ and we set $f_q(\theta, X) = \tilde{f}_q(\theta, X)w(X)^{-\alpha/2}$. f_q satisfies

$$\int_{\mathbb{Y} \times \mathbb{R}^{2d}} |f_q(\theta, X)|^2 w(X)^{\alpha} dX d\theta < \infty, \quad \mathcal{T}f_q = h_q, \quad \int_{\mathbb{Y}} Pf_q d\theta = 0$$

since multiplication by a (smooth enough) function of $H_0(X)$ commutes with the operator \mathcal{T} . Clearly, similar conclusions hold for the adjoint operator \mathcal{T}^* , which shows that the results can easily be extended to the weighted space framework.

Let us now turn to the very particular case we are interested in.

4.2 Solution of the profile equation (3.2)

The computation of the effective coefficients relies on the resolution of the profile equation with data $\nabla_x V_q(\theta, x) \cdot \nabla_p H_0(x, p)$. The compatibility condition (4.2) is satisfied in a strong way since we actually have

$$P(\nabla_x V_q \cdot \nabla_p H_0) = P\{V_q, H_0\} = 0.$$

This allows us to derive a more explicit expression for the solution χ_q (resp. χ_q^*) of the profile equation $\mathcal{T}\chi_q = \nabla_x V_q \cdot \nabla_p H_0$ (resp. $\mathcal{T}^*\chi_q^* = \nabla_x V_q \cdot \nabla_p H_0$).

Indeed, let us consider the profile equation $\mathcal{T}f_q = h_q$ under the condition $Ph_q = 0$. (Similar computations can be performed for the adjoint equation.) Then, applying the operator P to the equation yields $\omega \cdot \nabla_\theta Pf_q = Ph_q = 0$ which implies that Pf_q does not depend on θ . Requiring $\int_{\mathbb{Y}} Pf_q d\theta = 0$ gives $Pf_q = 0$. Therefore, we are led to solve

$$\begin{cases} \omega \cdot \nabla_\theta f_q + \{H_0, f_q\} + \gamma f_q = h_q, \\ Pf_q = 0. \end{cases}$$

Let us introduce the characteristics $\Theta \in \mathbb{R}^r$, $\bar{X} \in \mathbb{R}^{2d}$, the solutions of the ODEs system

$$\begin{cases} \frac{d}{ds}\Theta(s) = \omega, & \frac{d}{ds}\bar{X}(s) = (\nabla_p H_0(\bar{X}(s)), -\nabla_x H_0(\bar{X}(s))), \\ \Theta(0) = \theta, & \bar{X}(0) = (x, p). \end{cases}$$

Note in particular that $\Theta(s) = \theta + s\omega$. Hence, we get

$$\frac{d}{ds} \left(e^{\gamma s} f_q(\Theta(s), \bar{X}(s)) \right) = e^{\gamma s} h_q(\Theta(s), \bar{X}(s)).$$

Integration with respect to s yields the following statement:

Lemma 4.3 *Let $h_q \in L^2(\mathbb{Y} \times \mathbb{R}^{2d})$ be such that $Ph_q = 0$. Then the solution $f_q \in H_\#$ of $\mathcal{T}h_q = h_q$ with $Pf_q = 0$ is given by*

$$f_q(\theta, x, p) = \int_{-\infty}^0 e^{\gamma s} h_q(\Theta(s), \bar{X}(s)) ds. \quad (4.6)$$

Accordingly, if h_q lies in $C^0(\mathbb{Y}; L^2(\mathbb{R}^{2d}))$, then, f_q lies in the same space. If, furthermore $\nabla_X h_q$ lies in $C^0(\mathbb{Y}; L^2_{\text{loc}}(\mathbb{R}^{2d}))$, then, f_q also satisfies this property.

There only remains to discuss the regularity statement, which follows from a direct application of Lebesgue's dominated convergence theorem. Similarly, we can differentiate (4.6) with respect to X and conclude thanks to Hypothesis 1.3. Let us now state the precise result which will be used in the sequel:

Corollary 4.4 *Assume Hypothesis 1.1, 1.2, 1.3, 3.1. Then, there exists a unique function $\chi_q^* : \mathbb{Y} \times \mathbb{R}^{2d} \rightarrow \mathbb{R}$ such that*

$$\int_{\mathbb{Y} \times \mathbb{R}^{2d}} |\chi_q^*(\theta, X)|^2 \frac{dX d\theta}{w(X)^\beta} < \infty, \quad \mathcal{T}^* \chi_q^* = \nabla_x V_q \cdot \nabla_p H_0, \quad \int_{\mathbb{Y}} P \chi_q^* d\theta = 0.$$

It is defined by the formula

$$\chi_q^*(\theta, x, p) = \int_0^\infty e^{-\gamma s} \nabla_x V_q \cdot \nabla_p H_0(\theta + s\omega, \bar{X}(s; x, p)) ds.$$

Furthermore, for any $0 < R < \infty$, χ_q^* and $\nabla_X \chi_q^*$ belong to $C^0(\mathbb{Y}; L^2(B(0, R)))$, where $B(0, R) = \{X \in \mathbb{R}^{2d}, |X| \leq R\}$, and $P \chi_q^* = 0$.

Remark 4.1 *The role of Hypothesis 1.3 is to guarantee that χ_q^* possesses enough regularity to justify some algebraic manipulations below. If, instead of Hypothesis 1.3, we assume the weaker hypothesis $H_0 \in W^{2,\infty}(\mathbb{R}^{2d})$, we readily obtain the following estimate: $|\nabla_{x,p}\bar{X}(s)| \leq e^{Cs} (1 + |(x,p)|)$ for some $C > 0$. Then, all our results will remain true provided that we consider large enough values of the parameter γ (which should be $> C$). However, this looks too strong a restriction from a physical viewpoint because usually, relaxation rates are rather weak.*

5 Proof of Theorem 3.2

5.1 A priori estimates

We obtain the basic uniform estimate by multiplying (1.2) by f^ε and performing some integration by parts. Since the transport terms are antisymmetric, we get

$$\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^{2d}} |f^\varepsilon|^2 dX = \frac{\gamma}{\varepsilon^2} \int_{\mathbb{R}^{2d}} Q(f^\varepsilon) f^\varepsilon dX = -\frac{\gamma}{\varepsilon^2} \int_{\mathbb{R}^{2d}} |Pf^\varepsilon - f^\varepsilon|^2 dX \leq 0,$$

thanks to Corollary 2.2. Hence, we deduce the following claim.

Proposition 5.1 *Suppose that the initial data f_0^ε is bounded in $L^2(\mathbb{R}^{2d})$. Then,*

- (i) $(f^\varepsilon)_{\varepsilon>0}$ is bounded in $L^\infty(\mathbb{R}^+; L^2(\mathbb{R}^{2d}))$,
- (ii) $(g^\varepsilon = \frac{1}{\varepsilon}(f^\varepsilon - Pf^\varepsilon))_{\varepsilon>0}$ is bounded in $L^2(\mathbb{R}^+ \times \mathbb{R}^{2d})$.

Remark 5.1 *For any convex function $\Psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, we have*

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^{2d}} \Psi(f^\varepsilon) dX &= \gamma \int_{\mathbb{R}^{2d}} Q(f^\varepsilon) \Psi'(f^\varepsilon) dX \\ &= -\frac{\gamma}{\varepsilon^2} \int_{\mathbb{R}^{2d}} (Pf^\varepsilon - f^\varepsilon) \left(\Psi'(Pf^\varepsilon) - \Psi'(f^\varepsilon) \right) dX \leq 0 \end{aligned}$$

In particular, this provides uniform estimates of f^ε in any $L^p(\mathbb{R}^{2d})$ space, $1 \leq p \leq \infty$. However, these estimates will not be needed in the sequel.

5.2 Convergence proof

A possible proof would consist in solving the successive profile equations (3.1)-(3.3), constructing an approximate solution $f_{\text{app}}^\varepsilon = f^{(0)} + \varepsilon f^{(1)} + \varepsilon^2 f^{(2)}$, evaluating the difference $f^\varepsilon - f_{\text{app}}^\varepsilon$ and showing that it is $\mathcal{O}(\varepsilon)$. Such an approach is usually very demanding in terms of regularity of the solution and would lead to tedious technicalities. Moreover, the resolution of the profile equation (3.3) can impose more restrictions on the potential V_q than those detailed in Proposition 4.1. Here, we adopt another viewpoint, trying to pass to the limit in the equation. To this end, we follow the general homogenization strategy developed e.g. in [GP2]. It combines double scale convergence tools, as introduced in [N] and [A], combined with a suitable choice of test functions, the so-called ‘‘oscillating test functions method’’, see [Ev1, Ev2, T0, T]. First of all, let us give the following double scale convergence statement, which is adapted to the quasi-periodic framework.

Proposition 5.2 *Let f_ε be a bounded sequence in $L^2(\mathbb{R})$. Let $\omega \in \mathbb{R}^r$ the components of which are rationally independent. Then, there exists a subsequence, still labelled by ε , and a function $F_q \in L^2_{\#}(\mathbb{R} \times \mathbb{Y})$ such that for any test function $\psi_q \in L^2(\mathbb{R}; C^0_{\#}(\mathbb{Y}))$,² we have*

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} f_\varepsilon(t) \psi_q(t, \omega t / \varepsilon^2) dt = \int_{\mathbb{R}} \int_{\mathbb{Y}} F_q(t, \theta) \psi_q(t, \theta) d\theta dt.$$

The proof follows the arguments of [A], which are combined to the ergodic condition “ ω has rationally independent components”, through the use of a variant of the Birkhoff theorem (see [DS]). This is detailed in Appendix D. Further adaptations to sequences of functions with values in a Hilbert space can be readily obtained as in [GP1]. Therefore, coming back to Proposition 5.1, we have the following compactness property, where $C^0_{c,\#}(\mathbb{R} \times \mathbb{Y}; L^2(\mathbb{R}^{2d}))$ denotes the space of functions $\psi_q : \mathbb{R} \times \mathbb{R}^r \times \mathbb{R}^{2d} \rightarrow \mathbb{R}$ which are continuous with respect to $(t, \theta) \in \mathbb{R} \times \mathbb{R}^r$, \mathbb{Y} -periodic with respect to the second variable, with values in $L^2(\mathbb{R}^{2d})$, and such that $\psi_q(t, \theta, X) = 0$ when $t \notin K$, for some compact set $K \subset \mathbb{R}$.

Lemma 5.3 *We can suppose, up to the extraction of a subsequence, that f^ε converges to $F_q(t, \theta, X) \in L^2_{\#}((0, T) \times \mathbb{Y} \times \mathbb{R}^{2d})$ in the sense that*

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \psi_q(t, \omega t / \varepsilon^2, X) dX dt = \int_{\mathbb{R}} \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \psi_q(t, \theta, X) d\theta dX dt,$$

holds for any trial function $\psi_q \in C^0_{c,\#}(\mathbb{R} \times \mathbb{Y}; L^2(\mathbb{R}^{2d}))$. Furthermore, f^ε converges weakly in $L^2((0, T) \times \mathbb{R}^{2d})$ to $f(t, X) = \int_{\mathbb{Y}} F_q(t, \theta, X) d\theta$.

Let us multiply (1.2) by $\psi_q(t, \omega t / \varepsilon^2, X)$, where ψ_q is a C^∞ function of its arguments and is \mathbb{Y} -periodic with respect to the second variable. Integrations by parts yield

$$\begin{aligned} & \varepsilon \frac{d}{dt} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \psi_q(t, \omega t / \varepsilon^2, X) dX - \varepsilon \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \partial_t \psi_q(t, \omega t / \varepsilon^2, X) dX \\ & - \frac{1}{\varepsilon} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \omega \cdot \nabla_\theta \psi_q(t, \omega t / \varepsilon^2, X) dX - \frac{1}{\varepsilon} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \{H_0, \psi_q\}(t, \omega t / \varepsilon^2, X) dX \\ & + \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \nabla_x V_q(\omega t / \varepsilon^2, x) \cdot \nabla_p \psi_q(t, \omega t / \varepsilon^2, X) dX \\ & - \frac{1}{\varepsilon} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \gamma Q(\psi_q)(t, \omega t / \varepsilon^2, X) dX = 0 \end{aligned} \quad (5.1)$$

since $Q^* = Q$.

Hence, we first conclude that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \left[\omega \cdot \nabla_\theta \psi_q + \{H_0, \psi_q\} + \gamma Q(\psi_q) \right] (t, \omega t / \varepsilon^2, X) dX dt = 0 \\ & = \int_0^\infty \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \left[\omega \cdot \nabla_\theta \psi_q + \{H_0, \psi_q\} + \gamma Q(\psi_q) \right] (t, \theta, X) d\theta dX dt. \end{aligned}$$

²Referring to [A], section 5, $L^2(\mathbb{R}; C^0_{\#}(\mathbb{Y}))$ is the class of functions $\psi_q : \mathbb{R} \times \mathbb{R}^r \rightarrow \mathbb{R}$ which are measurable and square integrable with respect to the variable $t \in \mathbb{R}$, with values in the Banach space of continuous and \mathbb{Y} -periodic functions.

It implies that the double scale limit F_q does not depend on θ and is only a function of the energy; we denote $F_q(t, \theta, X) = F(t, H_0(X)) = f(t, X)$.

Next, we remark that for any function only depending on the energy, the most singular term in (5.1) vanishes. Accordingly, let us choose $\psi_q(t, \theta, X) = \varphi(H_0(X)) + \varepsilon\phi_q(t, \theta, X)$, with $\varphi \in C_c^\infty(\mathbb{R})$, as a test function. We get

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \left\{ \int_0^\infty \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \left[\omega \cdot \nabla_\theta \phi_q + \{H_0, \phi_q\} + \gamma Q(\phi_q) \right] (t, \omega t/\varepsilon^2, X) dX dt \right. \\ & \quad \left. - \int_0^\infty \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \nabla_x V_q(\omega t/\varepsilon^2, X) \cdot \nabla_p (\varphi(H_0(X))) dX dt \right\} = 0 \\ & = \int_0^\infty \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \left[\omega \cdot \nabla_\theta \phi_q + \{H_0, \phi_q\} + \gamma Q(\phi_q) \right] (t, \theta, X) d\theta dX dt \\ & \quad - \int_0^\infty \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \nabla_x V_q(\theta, X) \cdot \nabla_p H_0(X) \partial_E \varphi(H_0(X)) d\theta dX dt \\ & = - \int_0^\infty \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \nabla_x V_q(\theta, X) \cdot \nabla_p H_0(X) \partial_E \varphi(H_0(X)) d\theta dX dt = 0. \end{aligned}$$

Eventually, we choose ϕ_q depending on φ in such a way that the order $\mathcal{O}(1)$ term in (5.1) also vanishes. This is indeed possible by choosing ϕ_q a solution of the (adjoint) profile equation

$$\omega \cdot \nabla_\theta \phi_q + \{H_0, \phi_q\} + \gamma Q(\phi_q) = -\mathcal{T}^* \phi_q = \nabla_x V_q(\theta, X) \cdot \nabla_p H_0(X) \partial_E \varphi(H_0(X)).$$

Precisely, we set

$$\phi_q(\theta, X) = -\chi_q^*(\theta, X) \partial_E \varphi(H_0(X)).$$

with χ_q^* defined in Corollary 4.4. Note that by the regularity properties in Corollary 4.4, $\phi_q(\theta, X)$ and $\nabla_p \phi_q(\theta, X) = -\nabla_p \chi_q^* \partial_E \varphi(H_0(X)) - \chi_q^* \nabla_p H_0(X) \partial_{EE}^2 \varphi(H_0(X))$ can indeed be used as “admissible” test functions. It follows that

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \left(\varphi(H_0(X)) + \varepsilon \phi_q(\omega t/\varepsilon^2, X) \right) dX \\ & \quad + \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \nabla_x V_q(\omega t/\varepsilon^2, X) \cdot \nabla_p \phi_q(\omega t/\varepsilon^2, X) dX = 0, \end{aligned} \tag{5.2}$$

holds in $\mathcal{D}'((0, +\infty))$.

Equation (5.2) indicates that

$$\begin{aligned} & \left| \frac{d}{dt} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \left(\varphi(H_0(X)) + \varepsilon \phi_q(\omega t/\varepsilon^2, X) \right) dX \right| \\ & \leq \|f^\varepsilon\|_{L^\infty(\mathbb{R}^+; L^2(\mathbb{R}^{2d}))} \|\nabla_x V_q\|_{L^\infty(\mathbb{Y} \times \mathbb{R}^{2d})} \|\nabla_p \phi_q\|_{L^\infty(\mathbb{Y}; L^2(\mathbb{R}^{2d}))}, \end{aligned}$$

is uniformly bounded with respect to $\varepsilon > 0$, $0 \leq t \leq T < \infty$, thanks to Proposition 5.1, Hypothesis 3.1, Corollary 4.4 and the fact that φ has a compact support. Hence, for any φ fixed in $C_c^\infty(\mathbb{R})$, the family

$$\left\{ \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) (\varphi(H_0(X)) + \varepsilon \phi_q(\omega t/\varepsilon^2, X)) dX, \varepsilon > 0 \right\}$$

is relatively compact in $C^0([0, T])$, by virtue of the Arzela-Ascoli theorem. But, we also have

$$\begin{aligned} \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \varphi(H_0(X)) dX &= \int_{\mathbb{R}^{2d}} P f^\varepsilon(t, X) \varphi(H_0(X)) dX \\ &= \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) (\varphi(H_0(X)) + \varepsilon \phi_q(\omega t/\varepsilon^2, X)) dX - \varepsilon \int_{\mathbb{R}^{2d}} f^\varepsilon(t, X) \phi_q(\omega t/\varepsilon^2, X) dX \end{aligned}$$

where the last integral is dominated by

$$\|f^\varepsilon\|_{L^\infty(\mathbb{R}^+; L^2(\mathbb{R}^{2d}))} \|\chi_q^*\|_{L^\infty(\mathbb{Y}; L^2(\{X \in \mathbb{R}^{2d}, H_0(X) \in \text{supp}\varphi\}))} \|\varphi\|_{W^{1,\infty}(\mathbb{R})}.$$

Thus, the family

$$\left\{ \int_{\mathbb{R}^{2d}} P f^\varepsilon(t, X) \varphi(H_0(X)) dX, \varepsilon > 0 \right\}$$

is relatively compact in $C^0([0, T])$. Combining a separability and a diagonal extraction argument, we conclude that we can consider a subsequence, still labelled by ε , such that

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^{2d}} P f^\varepsilon(t, X) \varphi(H_0(X)) dX = \int_{\mathbb{R}^{2d}} F(t, H_0(X)) \varphi(H_0(X)) dX$$

uniformly on $[0, T]$, for any φ verifying $\int_{\mathbb{R}^{2d}} |\varphi(H_0(X))|^2 dX < \infty$.

Furthermore, the limit of the second integral in (5.2) as $\varepsilon \rightarrow 0$ reads

$$\begin{aligned} & \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \nabla_x V_q(\theta, x) \cdot \nabla_p \chi_q^*(\theta, X) \partial_E \varphi(H_0(X)) d\theta dX \\ & + \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} F_q(t, \theta, X) \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \chi_q^*(\theta, X) \partial_{EE}^2 \varphi(H_0(X)) d\theta dX \\ & = \int_{\mathbb{R}^{2d}} F(t, H_0(X)) \left(\int_{\mathbb{Y}} \nabla_x V_q(\theta, x) \cdot \nabla_p \chi_q^*(\theta, X) d\theta \right) \partial_E \varphi(H_0(X)) dX \\ & + \int_{\mathbb{R}^{2d}} F(t, H_0(X)) \left(\int_{\mathbb{Y}} \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \chi_q^*(\theta, X) d\theta \right) \partial_{EE}^2 \varphi(H_0(X)) dX \\ & = \int_{\mathbb{R}^{2d}} F(t, H_0(X)) \left(a^* \partial_E \varphi(H_0(X)) + b^* \partial_{EE}^2 \varphi(H_0(X)) \right) dX, \end{aligned}$$

since we have seen that $F_q(t, \theta, X) = F(t, H_0(X))$. Hence, letting ε tend to 0 in (5.2) yields

$$\frac{d}{dt} \int_{\mathbb{R}^{2d}} F(t, H_0(X)) \varphi(H_0(X)) dX = \int_{\mathbb{R}^{2d}} F(t, H_0(X)) \left(a^* \partial_E \varphi(H_0(X)) + b^* \partial_{EE}^2 \varphi(H_0(X)) \right) dX. \quad (5.3)$$

■

Let us detail some properties of the effective coefficients.

Lemma 5.4 *The coefficients a^* and b^* belong to $L_{\text{loc}}^2(\mathbb{R}, h_0(E) dE)$, and we have $b^*(E) \geq 0$ for almost all $E \in \mathbb{R}$. If furthermore, for any measurable set $A \subset \mathbb{R}$, and $\theta \in \mathbb{Y}$, we have*

$$(I - P)(\nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X)) \neq 0 \quad \text{on } \{X \in \mathbb{R}^{2d}, H_0(X) \in A\}$$

then, $b^(E) > 0$ almost everywhere.*

Proof. Regularity is a consequence of Corollary 4.4. Next, let $\varphi \in C_c^\infty(\mathbb{R})$. Thanks to Lemma 2.1-(iii), we get

$$\begin{aligned}
& \int_{\mathbb{R}^{2d}} b^*(H_0(X)) \varphi^2(H_0(X)) dX \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} (\nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X) \varphi(H_0(X))) (\chi_q^*(\theta, X) \varphi(H_0(X))) d\theta dX \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \mathcal{T}^*(\chi_q^*(\theta, X) \varphi(H_0(X))) \chi_q^*(\theta, X) \varphi(H_0(X)) d\theta dX \\
&= \gamma \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} |P \chi_q^*(\theta, X) \varphi(H_0(X)) - \chi_q^*(\theta, X) \varphi(H_0(X))|^2 d\theta dX \geq 0.
\end{aligned}$$

Next, suppose that $b^*(E) = 0$ for E in some measurable set $A \subset \mathbb{R}$. Let us set

$$\chi_{q,A}^*(\theta, X) = \chi_q^*(\theta, X) \mathbb{1}_{\{X \in \mathbb{R}^{2d}, H_0(X) \in A\}}(X).$$

Reasoning as above we obtain

$$\int_{\{X \in \mathbb{R}^{2d}, H_0(X) \in A\}} b^*(H_0(X)) dX = 0 = \gamma \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} |P \chi_{q,A}^* - \chi_{q,A}^*|^2 d\theta dX.$$

Therefore, $P \chi_{q,A}^* = \chi_{q,A}^*$, which implies that

$$\begin{aligned}
\mathbb{1}_{\{X \in \mathbb{R}^{2d}, H_0(X) \in A\}}(X) \mathcal{T}^* \chi_q^* &= \mathcal{T}^* \chi_{q,A}^* \\
&= \omega \cdot \nabla_\theta \chi_{q,A}^* \\
&= P(\omega \cdot \nabla_\theta \chi_{q,A}^*) \\
&= \mathbb{1}_{\{X \in \mathbb{R}^{2d}, H_0(X) \in A\}}(X) \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X)
\end{aligned}$$

holds. This would contradict the assumption $(I-P)(\mathbb{1}_{\{X \in \mathbb{R}^{2d}, H_0(X) \in A\}}(X) \nabla_x V_q(\theta, x) \cdot \nabla_p H_0(X)) \neq 0$ and proves that $b^*(E) > 0$ for $E \in \mathbb{R}$ a.e. . \blacksquare

We end the proof by showing that (5.3) is a weak formulation of the conservative equation (1.7). The coarea formula yields

$$\frac{d}{dt} \int_{\mathbb{R}} F(t, E) \varphi(E) h_0(E) dE = \int_{\mathbb{R}} F(t, E) \left(a^*(E) \partial_E \varphi(E) + b^*(E) \partial_{EE}^2 \varphi(E) \right) h_0(E) dE,$$

with $F \in L^\infty(\mathbb{R}^+; L^2(\mathbb{R}, h_0(E) dE))$, and the right hand side makes sense by Lemma 5.4. Then, Lemma 3.1 allows us to write:

$$\begin{aligned}
h_0(E) b^*(E) &= h_0(E) b(E) \in L_{\text{loc}}^2(\mathbb{R}, h_0(E)^{-1} dE), \\
h_0(E) a^*(E) &= \partial_E (h_0(E) b(E)) \in L_{\text{loc}}^2(\mathbb{R}, h_0(E)^{-1} dE).
\end{aligned}$$

Therefore, the right hand side in (5.3) becomes

$$\int_{\mathbb{R}} F(t, E) \partial_E \left(h_0(E) b(E) \partial_E \varphi(E) \right) dE,$$

which proves the expected result. \blacksquare

A The coarea formula and its consequences

Let $H_0 : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ be a C^∞ function. The Sard Theorem (see e.g. [M]) asserts that, for almost every real number³ $E \in \mathbb{R}$, and for any X such that $H_0(X) = E$, one has $\nabla_X H_0(X) \neq 0$. As a consequence, for almost every $E \in \mathbb{R}$, the level set $S_E := \{X \in \mathbb{R}^{2d}, H_0(X) = E\}$ is a smooth, codimension one, submanifold of \mathbb{R}^{2d} . Now, the coarea formula asserts that the following equality holds

$$\int_{\mathbb{R}^{2d}} f(X) dX = \int_{\mathbb{R}} \left(\int_{S_E} f(X) \delta(H_0(X) - E) \right) dE, \quad (\text{A.1})$$

for any function $f \in L^1(\mathbb{R}^{2d})$. We recall that the measure $\delta(H_0(X) - E)$ is defined by

$$\int_{S_E} f(X) \delta(H_0(X) - E) := \int_{S_E} f(X) \frac{d\sigma_E(X)}{|\nabla_X H_0(X)|}, \quad (\text{A.2})$$

using again the fact that the gradient $\nabla_X H_0(X)$ never vanishes on S_E , $d\sigma_E(X)$ being the euclidian surface measure on the level set S_E . We recall that a crucial hypothesis in our work is

$$h_0(E) := \int_{S_E} \delta(H_0(X) - E) < \infty \quad (\text{A.3})$$

for almost every $E \in \mathbb{R}$. Having defined the normalized average

$$\Pi f(E) = \frac{1}{h_0(E)} \int_{S_E} f(X) \delta(H_0(X) - E), \quad (\text{A.4})$$

for $f \in L^1(\mathbb{R}^{2d})$, the coarea formula then takes the form

$$\int_{\mathbb{R}^{2d}} f(X) dX = \int_{\mathbb{R}} \Pi f(E) h_0(E) dE. \quad (\text{A.5})$$

In particular, Π is an isometry from $L^1(\mathbb{R}^{2d})$ to $L^1(\mathbb{R}; h_0(E) dE)$. Since the analysis developed in the present paper needs an L^2 framework, we next turn to the L^2 properties of the operator Π .

Lemma A.1 *Let $f(X) : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ be in $L^2(\mathbb{R}^{2d})$. Then, we have*

$$\|\Pi f\|_{L^2(\mathbb{R}; h_0(E) dE)} \leq \|f\|_{L^2(\mathbb{R}^{2d})}.$$

Furthermore, let $g : \mathbb{R} \rightarrow \mathbb{R}$ satisfy $g \in L^2(\mathbb{R}; h_0(E) dE)$. The adjoint Π^ of the operator Π with respect to the scalar product in $L^2(\mathbb{R}; h_0(E) dE)$ is*

$$\Pi^* g(X) = g(H_0(X)).$$

It satisfies

$$\|\Pi^* g\|_{L^2(\mathbb{R}^{2d})} = \|g\|_{L^2(h_0(E) dE)}.$$

³Note that here, we make the same abuse of notation as in the main part of the present paper: instead of writing the correct condition $E \in H_0(\mathbb{R}^{2d})$, we simply write $E \in \mathbb{R}$.

Proof. First we use the Cauchy-Schwarz inequality together with the coarea formula and we get

$$\begin{aligned} \int_{\mathbb{R}} |\Pi f(E)|^2 h_0(E) dE &= \int_{\mathbb{R}} h_0(E) \left(\int_{S_E} f(X) \frac{\delta(H_0(X) - E)}{h_0(E)} \right)^2 dE \\ &\leq \int_{\mathbb{R}} \frac{h_0(E)}{h_0(E)^2} \left(\int_{S_E} |f(X)|^2 \delta(H_0(X) - E) \right) \left(\int_{S_E} \delta(H_0(X) - E) \right) dE \\ &\leq \int_{\mathbb{R}} \int_{S_E} |f(X)|^2 \delta(H_0(X) - E) dE = \int_{\mathbb{R}^{2d}} |f(X)|^2 dX. \end{aligned}$$

Next, we observe that

$$\begin{aligned} \langle \Pi f, g \rangle_{L^2(h_0(E)dE)} &= \int_{\mathbb{R}} g(E) \left(\int_{S_E} f(X) \delta(H_0(X) - E) \right) dE \\ &= \int_{\mathbb{R}} \int_{S_E} f(X) g(H_0(X)) \delta(H_0(X) - E) dE = \int_{\mathbb{R}^{2d}} f(X) g(H_0(X)) dX. \end{aligned}$$

Eventually, the coarea formula yields

$$\|\Pi^* g\|_{L^2(\mathbb{R}^{2d})}^2 = \int_{\mathbb{R}^{2d}} |g(H_0(X))|^2 dX = \int_{\mathbb{R}} \int_{S_E} |g(H_0(X))|^2 \delta(H_0(X) - E) dE = \int_{\mathbb{R}} |g(E)|^2 h_0(E) dE. \quad \blacksquare$$

B Dimensionless Equations

Let us detail the passage from (1.1) to its dimensionless version (1.2). The coefficients of the operator Q being dimensionless, $Q(f)$ has the same dimension as f itself, while $\tau > 0$ is a relaxation time. Let us introduce time and length scales, denoted by T and L respectively, and let P stand for a momentum unit. Then, we set

$$\begin{cases} t_* = t/T, & x_* = x/L, & p_* = p/P, \\ f_*(t_*, x_*, p_*) = L^d P^d f(t_* T, x_* L, p_* P), & H_{0,*}(x_*, p_*) = \frac{1}{H} H_0(x_* L, p_* P), \end{cases}$$

where the energy scale $H > 0$ characterizes the amplitude of the hamiltonian H_0 . It remains to discuss the perturbation \mathcal{V} . To this end, we introduce additional parameters:

- $\varepsilon > 0$, which is a dimensionless quantity measuring the strength of the perturbation compared with the free hamiltonian,

- $\theta > 0$, which is a characteristic time scale of the evolution of \mathcal{V} .

Hence, we have

$$\mathcal{V}(t, x) = \varepsilon H V_* \left(\frac{t}{\theta}, \frac{x}{L} \right).$$

Finally, (1.1) can be recast in the following dimensionless form

$$\partial_{t_*} f_* + \frac{TH}{LP} \{H_{0,*}, f_*\} + \varepsilon \frac{TH}{LP} \{V_*(t_* T/\theta), f_*\} = \frac{T}{\tau} Q_*(f_*).$$

Then, our analysis is based on the following scaling assumptions. First, we suppose that

$$\frac{\text{TH}}{\text{LP}} = \frac{1}{\varepsilon^2} \gg 1.$$

Roughly speaking it means that the time unit we adopt is large compared with the characteristic time scale of the free hamiltonian H_0 (e.g. for the harmonic oscillator the period of the characteristic curves). Next, we are interested in the behavior of the system as $\varepsilon \ll 1$ when the time scales involved in the problem satisfy the following ordering:

$$\frac{\text{T}}{\theta} = \frac{1}{\varepsilon^2}, \quad \frac{\text{T}}{\tau} = \frac{\gamma}{\varepsilon^2}, \quad \gamma = \mathcal{O}(1).$$

Here, $\gamma > 0$ is a fixed dimensionless quantity. This sets up the asymptotic regime we are dealing with.

C Effective coefficients: Proof of Lemma 3.1

Let $\psi \in C_c^\infty(\mathbb{R})$. The coarea formula (A.5) yields

$$\begin{aligned} \int_{\mathbb{R}} h_0 b^* \psi dE &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \{V_q, H_0\} \chi_q^*(\theta, X) \psi(H_0(X)) d\theta dX \\ &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \mathcal{T} \chi_q \chi_q^*(\theta, X) \psi(H_0(X)) d\theta dX \\ &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q \mathcal{T}^* \left(\chi_q^*(\theta, X) \psi(H_0(X)) \right) d\theta dX \\ &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q \mathcal{T}^* \chi_q^*(\theta, X) \psi(H_0(X)) d\theta dX \\ &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q \{V_q, H_0\} \psi(H_0(X)) d\theta dX = \int_{\mathbb{R}} h_0 b \psi dE. \end{aligned}$$

Similarly, combining the coarea formula and integration by parts, we get

$$\begin{aligned} \int_{\mathbb{R}} h_0 a^* \psi dE &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \{V_q, \chi_q^*\}(\theta, X) \psi(H_0(X)) d\theta dX \\ &= - \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q^* \{V_q, \psi(H_0(X))\} d\theta dX \\ &= - \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q^* \{V_q, H_0(X)\} (\partial_E \psi)(H_0(X)) d\theta dX \\ &= - \int_{\mathbb{R}} h_0 b^* \partial_E \psi dE, \end{aligned}$$

which proves $h_0 a^* = \partial_E(h_0 b^*)$. We obtain the equality $h_0 a^* = h_0 a$ by remarking that

$$\begin{aligned}
& \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q^* \{V_q, H_0(X)\} (\partial_E \psi)(H_0(X)) d\theta dX \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \chi_q^* \mathcal{T} \chi_q (\partial_E \psi)(H_0(X)) d\theta dX \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \mathcal{T}^* \chi_q^* \chi_q (\partial_E \psi)(H_0(X)) d\theta dX \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \{V_q, H_0(X)\} \chi_q (\partial_E \psi)(H_0(X)) d\theta dX \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{Y}} \{V_q, \psi(H_0(X))\} \chi_q d\theta dX
\end{aligned}$$

holds. An integration by parts allows to conclude the proof. \blacksquare

D Double scale convergence: Proof of Proposition 5.2

The double scale convergence framework has been extended to very complicated and general oscillating coefficients, which leads to tedious technicalities; we refer on these aspects to [CG] and [N2]. The case of quasi-periodic coefficients we are dealing with can be treated by following closely the arguments of [A]. Indeed, consider a bounded sequence in $L^2(\mathbb{R})$

$$\sup_{\varepsilon > 0} \int_{\mathbb{R}} |f_\varepsilon(t)|^2 dt \leq C < \infty.$$

Let \mathcal{A} stand for the space $L^2(\mathbb{R}; C_{\#}^0(\mathbb{Y}))$, which is a separable Banach space. Let $\phi \in \mathcal{A}$ and remark that

$$\begin{aligned}
\left| \int_{\mathbb{R}} f_\varepsilon(t) \phi(t, \omega t/\varepsilon) dt \right| &\leq \|f_\varepsilon\|_{L^2(\mathbb{R})} \left(\int_{\mathbb{R}} \left(\sup_{z \in \mathbb{Y}} |\phi(t, z)| \right)^2 dt \right)^{1/2} \\
&\leq C \|\phi\|_{\mathcal{A}}.
\end{aligned}$$

Hence, if we denote by Θ_ε the linear form defined by

$$\langle \Theta_\varepsilon, \phi \rangle = \int_{\mathbb{R}} f_\varepsilon(t) \phi(t, \omega t/\varepsilon) dt,$$

we conclude that $(\Theta_\varepsilon)_{\varepsilon > 0}$ is bounded in the dual set \mathcal{A}' . Hence, by the Banach-Alaoglu theorem, we can suppose that Θ_ε converges to some ν weakly- \star in \mathcal{A}' . However, we also have:

$$\left| \int_{\mathbb{R}} f_\varepsilon(t) \phi(t, \omega t/\varepsilon) dt \right| \leq C \left(\int_{\mathbb{R}} |\phi(t, \omega t/\varepsilon)|^2 dt \right)^{1/2},$$

so that letting ε tend to 0 yields:

$$|\langle \nu, \phi \rangle| \leq C \lim_{\varepsilon \rightarrow 0} \left(\int_{\mathbb{R}} |\phi(t, \omega t/\varepsilon)|^2 dt \right)^{1/2}.$$

Therefore, we can identify ν with a function $F \in L^2_{\#}(\mathbb{R} \times \mathbb{Y})$ by the Riesz theorem once we are able to justify that

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} |\phi(t, \omega t/\varepsilon)|^2 dt = \int_{\mathbb{Y}} \int_{\mathbb{R}} |\phi(t, \theta)|^2 d\theta dt.$$

The proof of this fact follows the arguments of [A], with some slight modifications; the adaptation to the quasi-periodic framework can be seen as a version of the Birkhoff ergodic theorem, see [DS]. It is a consequence of the two following claims.

Lemma D.1 *Let ω be a element of \mathbb{R}^r the components of which are rationally independent. Let $\phi \in C^0_{\#}(\mathbb{Y})$. Then $\phi(\omega t/\varepsilon) \rightarrow \int_{\mathbb{Y}} \phi(\theta) d\theta$ weakly- \star in $L^\infty(\mathbb{R})$.*

Proof. We start by proving the result for $\phi(\theta) = \exp(2i\pi k \cdot \theta)$, $k \in \mathbb{Z}^r$. Indeed, let $\psi \in L^1(\mathbb{R})$. We get

$$\int_{\mathbb{R}} \psi(t) e^{2i\pi k \cdot \omega t/\varepsilon} dt \widehat{\psi} \left(-\frac{2\pi k \cdot \omega}{\varepsilon} \right).$$

Therefore, for $k = 0$, this is nothing but

$$\widehat{\psi}(0) = \int_{\mathbb{R}} \psi(t) dt = \int_{\mathbb{R}} \psi(t) dt \int_{\mathbb{Y}} e^{2i\pi 0 \cdot \theta} d\theta,$$

while for $k \neq 0$, the ergodic condition $k \cdot \omega \neq 0$ yields

$$\lim_{\varepsilon \rightarrow 0} \widehat{\psi} \left(-\frac{2\pi k \cdot \omega}{\varepsilon} \right) = 0 = \int_{\mathbb{R}} \psi(t) dt \int_{\mathbb{Y}} e^{2i\pi k \cdot \theta} d\theta.$$

Of course, we immediately deduce that the result also applies to any trigonometric polynomial.

Then, we extend the property to any $\phi \in C^0_{\#}(\mathbb{Y})$. Indeed, such a function can be approached, in the sup norm sense, by a sequence $(p_n)_{n \in \mathbb{N}}$ of trigonometric polynomials. Then, we note that

$$\begin{aligned} & \left| \int_{\mathbb{R}} \psi(t) \phi(\omega t/\varepsilon) dt - \int_{\mathbb{R}} \psi(t) \left(\int_{\mathbb{Y}} \phi(\theta) d\theta \right) dt \right| \\ & \leq \int_{\mathbb{R}} |\psi(t)| |\phi(\omega t/\varepsilon) - p_n(\omega t/\varepsilon)| dt + \left| \int_{\mathbb{R}} \psi(t) p_n(\omega t/\varepsilon) dt - \int_{\mathbb{R}} \psi(t) \left(\int_{\mathbb{Y}} p_n(\theta) d\theta \right) dt \right| \\ & \quad + \int_{\mathbb{R}} |\psi(t)| \int_{\mathbb{Y}} |\phi(\theta) - p_n(\theta)| d\theta dt \\ & \leq 2 \|\psi\|_{L^1(\mathbb{R})} \|\phi - p_n\|_{L^\infty(\mathbb{Y})} + \left| \int_{\mathbb{R}} \psi(t) p_n(\omega t/\varepsilon) dt - \int_{\mathbb{R}} \psi(t) \left(\int_{\mathbb{Y}} p_n(\theta) d\theta \right) dt \right|. \end{aligned}$$

Let $\delta > 0$ be a positive number. Then, there exists $n = n(\delta)$ such that the first term at the right hand side is less than δ . Eventually, the previous step of the proof guarantees that for $0 < \varepsilon < \varepsilon(\delta)$ small enough, the last integral is also less than δ . This ends the proof. \blacksquare

Lemma D.2 *Let ω be an element of \mathbb{R}^r the components of which are rationally independent. Let $\phi \in L^1(\mathbb{R}; C_{\#}^0(\mathbb{Y}))$. Then, we have*

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} \phi(t, \omega t / \varepsilon) dt = \int_{\mathbb{Y}} \int_{\mathbb{R}} \phi(t, \theta) d\theta dt.$$

Proof. Let us introduce a covering of the unit cube of \mathbb{R}^r , made of $I(n)$ open sets O_i with diameter $\leq \alpha_n$, where we assume that $I(n) \rightarrow \infty$ and $\alpha_n \rightarrow 0$ as n goes to ∞ . For each $i \in \{1, \dots, I(n)\}$, Let θ_i be an element of O_i . To this covering, we associate a set of functions ζ_i , $i \in \{1, \dots, I(n)\}$ such that

$$0 \leq \zeta_i(\theta) \leq 1, \quad \text{supp}(\zeta_i) \subset O_i, \quad \sum_{i=1}^{I(n)} \zeta_i(\theta) = 1,$$

and we extend these functions to \mathbb{R}^r by periodicity. Let $\phi \in L^1(\mathbb{R}; C_{\#}^0(\mathbb{Y}))$. We set

$$\phi_n(t, \theta) = \sum_{i=1}^{I(n)} \phi(t, \theta_i) \zeta_i(\theta).$$

Then, we note that

$$\begin{aligned} |\phi(t, \theta) - \phi_n(t, \theta)| &= \left| \sum_{i=1}^{I(n)} \zeta_i(\theta) (\phi(t, \theta_i) - \phi(t, \theta)) \right| \\ &\leq \sum_{i=1}^{I(n)} \zeta_i(\theta) \sup_{\theta \in O_i} |\phi(t, \theta_i) - \phi(t, \theta)|. \end{aligned}$$

Since, for $t \in \mathbb{R}$ a.e., the function $\theta \mapsto \phi(t, \theta)$ is continuous on the compact set \mathbb{Y} , and for $\theta \in O_i$, $|\theta - \theta_i| \leq \alpha_n \rightarrow 0$, we deduce that $\sup_{\theta \in \mathbb{Y}} |\phi(t, \theta) - \phi_n(t, \theta)| \rightarrow 0$ as n goes to ∞ . Besides, we have $\sup_{\theta \in \mathbb{Y}} |\phi(t, \theta) - \phi_n(t, \theta)| \leq 2 \|\phi(t, \cdot)\|_{L^\infty(\mathbb{Y})} \in L^1(\mathbb{R})$. Therefore, the Lebesgue theorem yields

$$\|\phi - \phi_n\|_{L^1(\mathbb{R}, L^\infty(\mathbb{Y}))} \xrightarrow{n \rightarrow \infty} 0. \quad (\text{D.1})$$

Then, for $n \in \mathbb{N}$ fixed, we write

$$\int_{\mathbb{R}} \phi_n(t, \omega t / \varepsilon) dt = \sum_{i=1}^{I(n)} \int_{\mathbb{R}} \phi(t, \theta_i) \zeta_i(\omega t / \varepsilon) dt.$$

Since $t \mapsto \phi(t, \theta_i)$ belongs to $L^1(\mathbb{R})$ and $\zeta_i \in C_{\#}^0(\mathbb{Y})$, Lemma D.1 applies and leads to

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} \phi_n(t, \omega t / \varepsilon) dt = \sum_{i=1}^{I(n)} \int_{\mathbb{R}} \phi(t, \theta_i) \left(\int_{\mathbb{Y}} \zeta_i(\theta) d\theta \right) dt = \int_{\mathbb{R}} \int_{\mathbb{Y}} \phi_n(t, \theta) d\theta dt.$$

Combining this to (D.1) ends the proof. ■

E A simple example

It is worth illustrating the previous developments with a fully explicit computation. This can be performed when considering Hamiltonians based on the harmonic oscillator

$$H_{\text{harm}}(X) = |X|^2/2 = \frac{x^2 + p^2}{2},$$

with $X = (x, p) \in \mathbb{R}^2$ and the simplest perturbation

$$V(t/\varepsilon^2, x) = x \cos(\omega t/\varepsilon^2), \quad \omega \in \mathbb{R} \setminus \{0\}.$$

Let us consider the following Hamiltonian

$$H_0(X) = G(H_{\text{harm}}(X)),$$

with $G : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a C^1 , strictly increasing function. We note that $H_0(X) = E$ iff $|X|^2 = 2G^{-1}(E)$. Therefore, integration over S_E reduces to integration over the sphere of \mathbb{R}^2 with radius $\sqrt{2G^{-1}(E)}$: we write $(x, p) \in S_E$ as $x = \sqrt{2G^{-1}(E)} \cos(\sigma)$, $p = \sqrt{2G^{-1}(E)} \sin(\sigma)$, with $\sigma \in (0, 2\pi)$ and $d\sigma_E$ becomes $\sqrt{2G^{-1}(E)} d\sigma$. Next, we compute

$$\nabla H_0(X) = G'(|X|^2/2) \begin{pmatrix} x \\ p \end{pmatrix},$$

so that $|\nabla H_0(X)| = G'(|X|^2/2) |X| = G'(G^{-1}(E)) \sqrt{2G^{-1}(E)}$. In what follows, we denote

$$\Omega(E) = G'(G^{-1}(E)).$$

Hence, we obtain

$$h_0(E) = \int_{x^2+p^2=2G^{-1}(E)} \frac{d\sigma_E}{|\nabla H_0(x, p)|} \int_0^{2\pi} \frac{\sqrt{2G^{-1}(E)}}{\Omega(E) \sqrt{2G^{-1}(E)}} d\sigma = \frac{2\pi}{\Omega(E)},$$

and

$$\Pi f(E) = \frac{1}{2\pi} \int_0^{2\pi} f\left(\sqrt{2G^{-1}(E)} \cos(\sigma), \sqrt{2G^{-1}(E)} \sin(\sigma)\right) d\sigma.$$

The characteristics $\bar{X}(t; x, p) = (\bar{x}(t; x, p), \bar{p}(t; x, p))$ verify

$$\frac{d}{dt} \bar{X}(t; x, p) = G'(|\bar{X}(t; x, p)|^2/2) \begin{pmatrix} \bar{p}(t; x, p) \\ -\bar{x}(t; x, p) \end{pmatrix}, \quad \bar{X}(0; x, p) = \begin{pmatrix} x \\ p \end{pmatrix}.$$

The keypoint relies on the observation that $X(t; x, p)$ lies on the same sphere of \mathbb{R}^2 than the initial data. Indeed, we have

$$\frac{d}{dt} H_0(\bar{X}(t; x, p)) = 0.$$

Since G is a diffeomorphism, we deduce that

$$\bar{x}(t; x, p)^2 + \bar{p}(t; x, p)^2 = x^2 + p^2 = 2G^{-1}(E).$$

In turn, $\bar{x}(t; x, p)$ satisfies the following simple second order ODE

$$\begin{aligned} \frac{d^2}{dt^2} \bar{x}(t; x, p) &= \frac{d}{dt} \left[G'(|\bar{X}(t; x, p)|^2/2) \bar{p}(t; x, p) \right] G'(|\bar{X}(t; x, p)|^2/2) \frac{d}{dt} \bar{p}(t; x, p) \\ &= -\Omega(E)^2 \bar{x}(t; x, p). \end{aligned}$$

We immediatly solve this ODE, and we finally obtain

$$\bar{X}(t; x, p) = \begin{pmatrix} \cos(\Omega(E)t) & \sin(\Omega(E)t) \\ -\sin(\Omega(E)t) & \cos(\Omega(E)t) \end{pmatrix} \begin{pmatrix} x \\ p \end{pmatrix}, \quad E = G\left(\frac{x^2 + p^2}{2}\right).$$

In particular, we note that

$$\nabla_{x,p} \bar{X}(t; x, p) = \begin{pmatrix} \cos(\Omega(E)t) + \bar{p}(t; x, p) t\Omega\Omega'(E) x & \sin(\Omega(E)t) + \bar{p}(t; x, p) t\Omega\Omega'(E) p \\ -\sin(\Omega(E)t) - \bar{x}(t; x, p) t\Omega\Omega'(E) x & \cos(\Omega(E)t) - \bar{x}(t; x, p) t\Omega\Omega'(E) p \end{pmatrix}.$$

Therefore, Hypothesis 1.3 is satisfied since $E \mapsto \Omega\Omega'(E)$ is locally bounded. Of course, this is also true in the purely harmonic case ($G(h) = h$, $\Omega(E) = 1$).

It remains to compute the effective coefficients. Since $\partial_p H_0(x, p) = \Omega(E) p$, we get

$$\chi(\theta, x, p) \int_0^\infty e^{-\gamma s} \cos(\theta - \omega s) \Omega(E) (x \sin(\Omega(E)s) + p \cos(\Omega(E)s)) ds.$$

Then, we are led to

$$\begin{aligned} b(E) &= \Pi \left(\int_0^{2\pi} \partial_x V(\theta, x) \partial_p H_{\text{harm}} \chi^*(\theta, x, p) d\theta \right) (E) \\ &= \frac{1}{2\pi} \int_0^\infty \int_0^{2\pi} \int_0^{2\pi} \cos(\theta) \Omega(E) \sqrt{2G^{-1}(E)} \sin(\sigma) e^{-\gamma s} \cos(\theta - \omega s) \\ &\quad \times \Omega(E) \sqrt{2G^{-1}(E)} (\cos(\sigma) \sin(\Omega(E)s) + \sin(\sigma) \cos(\Omega(E)s)) d\theta d\sigma ds \\ &= \frac{2G^{-1}(E) \Omega(E)^2}{2\pi} \int_0^\infty e^{-\gamma s} \pi \cos(\Omega(E)s) \left(\int_0^{2\pi} \cos(\theta) \cos(\theta - \omega s) d\theta \right) ds \\ &= \pi G^{-1}(E) \Omega(E)^2 \int_0^\infty e^{-\gamma s} \cos(\Omega(E)s) \cos(\omega s) ds \\ &= \pi \frac{G^{-1}(E) \Omega(E)^2}{2} \left(\frac{\gamma}{(\omega + \Omega(E))^2 + \gamma^2} + \frac{\gamma}{(\omega - \Omega(E))^2 + \gamma^2} \right). \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} a(E) &= \Pi \left(\int_0^{2\pi} \partial_x V(\theta, x) \partial_p \chi^*(\theta, x, p) d\theta \right) (E) \\ &= \frac{1}{h_0(E)} \partial_E (h_0 b^*(E)) \\ &= \frac{\pi}{2} \Omega(E) \partial_E \left[\Omega(E) G^{-1}(E) \left(\frac{\gamma}{(\omega + \Omega(E))^2 + \gamma^2} + \frac{\gamma}{(\omega - \Omega(E))^2 + \gamma^2} \right) \right] \end{aligned}$$

Let us end with a couple of remarks concerning these computations. Notice that the diffusion coefficient $b(E)$ vanishes when $G^{-1}(E)$ or $\Omega(E)$ vanish, which is the case for the harmonic oscillator at the energy $E = 0$. The coefficient becomes infinite when G has an infinite derivative. Remark that the limit $\gamma \rightarrow 0$ reveals resonance phenomena: dealing with the purely harmonic case ($G(h) = h$, $\Omega(E) = 1$), we remark that the coefficients tend to ∞ as $\gamma \rightarrow 0$ if the perturbation V oscillates with the characteristic frequency of the system $\omega = \pm 1$. The situation can be different when dealing with another function G . Indeed, if the equation $\Omega(E) = \pm\omega$ has a finite number of solutions $\{E_1, \dots, E_I\}$, resonances only occur on this finite set of energies.

Of course, it is also interesting to compare with the explicit solution of the kinetic equation

$$\partial_t f^\varepsilon + \frac{1}{\varepsilon^2} \{H_0, f^\varepsilon\} + \frac{1}{\varepsilon} \{V(t/\varepsilon^2), f^\varepsilon\} = 0,$$

that can be obtained in the simplest case $H_0(x, p) = (x^2 + p^2)/2$ and $V(t, x) = x \cos(\omega t)$. Indeed, the characteristics associated with the full Hamiltonian can be readily computed. They satisfy the ODE system

$$\begin{cases} \frac{d}{ds} \tilde{x}(s; t, x, p) = \frac{1}{\varepsilon^2} \tilde{p}(s; t, x, p), & \frac{d}{ds} \tilde{p}(s; t, x, p) = -\frac{1}{\varepsilon^2} \tilde{x}(s; t, x, p) + \frac{1}{\varepsilon} \cos(\omega s/\varepsilon^2), \\ \tilde{x}(t; t, x, p) = x, & \tilde{p}(t; t, x, p) = p. \end{cases}$$

We get for $\omega \neq \pm 1$:

$$\begin{aligned} \tilde{x}(0; t, x, p) &= x \cos(t/\varepsilon^2) - p \sin(t/\varepsilon^2) + \frac{\varepsilon}{2} \left(\frac{1 - \cos((1 + \omega)t/\varepsilon^2)}{1 + \omega} + \frac{1 - \cos((1 - \omega)t/\varepsilon^2)}{1 - \omega} \right), \\ \tilde{p}(0; t, x, p) &= x \sin(t/\varepsilon^2) + p \cos(t/\varepsilon^2) - \frac{\varepsilon}{2} \left(\frac{\sin((1 + \omega)t/\varepsilon^2)}{1 + \omega} + \frac{\sin((1 - \omega)t/\varepsilon^2)}{1 - \omega} \right), \end{aligned}$$

and for $\omega = \pm 1$:

$$\begin{aligned} \tilde{x}(0; t, x, p) &= x \cos(t/\varepsilon^2) - p \sin(t/\varepsilon^2) + \frac{\varepsilon}{2} \frac{1 - \cos(2t/\varepsilon^2)}{2}, \\ \tilde{p}(0; t, x, p) &= x \sin(t/\varepsilon^2) + p \cos(t/\varepsilon^2) - \frac{\varepsilon}{4} \sin(2t/\varepsilon^2) - \frac{t}{2\varepsilon}. \end{aligned}$$

Given an initial data f_0 , we thus have

$$f^\varepsilon(t, x, p) = f_0(\tilde{x}(0; t, x, p), \tilde{p}(0; t, x, p)),$$

which develops different features than solutions of a diffusion equation.

References

- [Ale1] R. Alexandre, *Some results in homogenization tackling memory effects*, Asymptot. Anal. 15 no. 3-4, 229–259 (1997).

- [Ale2] R. Alexandre, *Asymptotic behaviour of transport equations*, Appl. Anal. 70 no. 3-4, 405–430 (1999).
- [A] G. Allaire, *Homogenization and two-scale convergence*, SIAM J. Math. Anal., 23, 1482–1518 (1992).
- [AHZ] Y. Amirat, K. Hamdache, A. Ziani, *On homogenization of ordinary differential equations and linear transport equations* in **Calculus of variations, homogenization and continuum mechanics** (Marseille, 1993) Ser. Adv. Math. Appl. Sci., vol. 18, (World Sci. Publishing, 1994) pp. 29–50.
- [BPR] G. Bal, G. Papanicolaou, L. Ryzhik, *Radiative transport limit for the random Schrödinger equation*, Nonlinearity 15 no. 2, 513–529 (2002).
- [BLP] A. Bensoussan, J.-L. Lions, G. Papanicolaou, **Asymptotic analysis for periodic structures**, Studies in Mathematics and its Applications, 5 (North-Holland, 1978).
- [BDG] C. Bardos, L. Dumas, F. Golse, *Diffusion approximation for billiards with totally accommodating scatterers*, J. Statist. Phys. 86 no. 1-2, 351–375 (1997).
- [BCD] B. Bidégaray, F. Castella, P. Degond, *From Bloch model to the rate equations*, Discrete Contin. Dyn. Syst., 11, no. 1, 1–26 (2004).
- [BCDG] B. Bidégaray, F. Castella, E. Dumas, M. Gisclon, *From Bloch model to the rate equations II: the case of almost degenerate energy levels*, Math. Models Methods Appl. Sci., 14, no. 12, 1785–1817 (2004).
- [BCS] L. A. Bunimovich, N.I. Chernov, Ya. G. Sinai, *Statistical properties of two-dimensional hyperbolic billiards*, Uspekhi Mat. Nauk 46 no. 4, 43–92, (1991); translation in Russian Math. Surveys 46, no. 4, 47–106 (1991).
- [CG] J. Casado-Díaz, I. Gayte, *The two-scale convergence method applied to generalized Besicovitch spaces*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. 458 no. 2028, 2925–2946 (2002).
- [Ca1] F. Castella, *On the derivation of a Quantum Boltzmann Equation from the periodic von Neumann equation*, Mod. Math. An. Num., Vol. 33, N. 2, 329-349 (1999).
- [Ca2] F. Castella, *From the von Neumann equation to the Quantum Boltzmann equation in a deterministic framework*, J. Stat. Phys., Vol. 104, N. 1/2, 387–447 (2001).
- [CD] F. Castella, P. Degond *Convergence de l'équation de von Neumann vers l'équation de Boltzmann Quantique dans un cadre déterministe*, C. R. Acad. Sci., t. 329, sér. I, 231–236 (1999).
- [DS] N. Dunford, J. Schwartz, **Linear operators** (John Wiley & Sons, 1988).

- [EY1] L. Erdős, H. T. Yau, *Linear Boltzmann equation as scaling limit of quantum Lorentz gas*, Advances in Differential Equations and Mathematical Physics, Contemporary Mathematics 217, 135–155 (1998).
- [EY2] L. Erdős, H. T. Yau, *Linear Boltzmann equation as the weak coupling limit of a random Schrödinger equation*, Commun. Pure Appl. Math. 53, 667–735 (2000).
- [Ev1] L. C. Evans, *The perturbed test function method for viscosity solutions of nonlinear PDE*, Proc. Roy. Soc. Edinburgh Sect. A, 111, 359–375 (1989).
- [Ev2] L. C. Evans, *Periodic homogenization of certain fully nonlinear partial differential equations*, Proc. Roy. Soc. Edinburgh Sect. A, 120, 245–265 (1992).
- [FH] E. Frenod, K. Hamdache, *Homogenisation of transport kinetic equations with oscillating potentials*, Proc. Roy. Soc. Edinburgh Sect. A, 126, no. 6, 1247–1275 (1996).
- [GP1] T. Goudon, F. Poupaud, *Approximation by homogenization and diffusion of kinetic equations*, Comm. P.D.E., 26, 537–570 (2001).
- [GP2] T. Goudon, F. Poupaud, *Homogenization of transport equations: weak mean field approximation*, SIAM J. Math. Anal., 36, 856–881 (2004).
- [GP3] T. Goudon, F. Poupaud, *Homogenization of transport equations: A simple PDE approach to the Kubo formula*, Preprint.
- [KPR] J.B. Keller, G. Papanicolaou, L. Ryzhik, *Transport equations for elastic and other waves in random media*, Wave Motion, 24, no. 4, 327–370 (1996).
- [La] O. Lanford III, *The evolution of large classical systems*, in: Moser, J. (ed) Dynamical systems, theory and applications. Lectures Notes in Physics 35 (Springer, 1975) pp. 1–111.
- [LV] G. Loeper, A. Vasseur, *Electric turbulence in a plasma subject to a strong magnetic field*, Asymptot. Anal., 40, no. 1, 51–65 (2004).
- [Lo] R. Loudon, **The quantum theory of light** (Oxford Science Publications, 1983).
- [M] J.W. Milnor, **Topology from the differentiable viewpoint** (The University Press of Virginia, 1963).
- [N] G. Nguetseng, *A general convergence result for a functional related to the theory of homogenization*, SIAM J. Math. Anal., 20, 608–623 (1989).
- [N2] G. Nguetseng, *Almost periodic homogenization: asymptotic analysis of a second order elliptic equation*, Publications Mathématiques du laboratoire d’analyse numérique, Université Yaoundé I.
- [PR] B. Perthame, L. Ryzhik, *The quantum scattering limit for a regularized Wigner equation*, Preprint.

- [Pe] M. Petrini, *Homogenization of a linear transport equation with time depending coefficient*, Rend. Sem. Mat. Univ. Padova 101, 191–207 (1999).
- [PV] F. Poupaud, A. Vasseur, *Classical and quantum transport in random media*, Journal de Mathématiques Pures et Appliquées, 82, no. 6, 711–748 (2003).
- [SP] E. Sánchez-Palencia, **Nonhomogeneous media and vibration theory**, Lecture Notes in Physics, 127 (Springer-Verlag, 1980).
- [SV] J.A. Sanders, F. Verhulst, **Averaging methods in nonlinear dynamical systems**, Applied Mathematical Sciences, Vol. 59 (Springer-Verlag, 1985).
- [Sp] H. Spohn, **Large scale dynamics of interacting particles** (Springer, 1991).
- [T0] L. Tartar, *H-convergence et compacité par compensation*, Cours Peccot, Collège de France, March 1977. Partially written in: F. Murat, *H-convergence*, Séminaire d'Analyse Fonctionnelle et Numérique 1977-78, Université d'Alger, multicopied 34 p. English translation in F. Murat, L. Tartar, *H-convergence*, in **Topics in the mathematical modelling of composite materials**, A. Cherkaev - R. V. Kohn eds., Progress in Nonlinear Differential Equations Appl., 31 (Birkhäuser, 1997) pp. 21-43.
- [T] L. Tartar, *Remarks on homogenization* in **Homogenization and effective moduli of material and media**, IMA Vol. in Math. and Appl. (Springer, 1986) pp. 228–246.