

November 4, 2013

GROUND ENERGY OF THE MAGNETIC LAPLACIAN IN POLYHEDRAL BODIES

VIRGINIE BONNAILLIE-NOËL, MONIQUE DAUGE, NICOLAS POPOFF

Abstract. The asymptotic behavior of the first eigenvalues of magnetic Laplacian operators with large magnetic fields and Neumann realization in polyhedral domains is characterized by a hierarchy of model problems. We investigate properties of the model problems (continuity, semi-continuity, existence of generalized eigenfunctions). We prove estimates for the remainders of our asymptotic formula. Lower bounds are obtained with the help of a classical IMS partition based on adequate coverings of the polyhedral domain, whereas upper bounds are established by a novel construction of quasimodes, qualified as sitting or sliding according to spectral properties of local model problems.

Contents

1. Introduction. Main results	2
1.1. Asymptotic formulas with remainders	3
1.2. Contents of the paper	5
1.3. Notations	5
2. State of the art	6
2.1. Dimension 2	6
2.2. Dimension 3	8
3. Polyhedral domains and their singular chains	9
3.1. Domains and tangent cones	9
3.2. Recursive definition of singular chains	10
3.3. Polyhedral domains	11
4. Taxonomy of model problems	13
4.1. Tangent and model operators	13
4.2. Singular chains and generalized eigenvectors for model problems	13
4.3. Dichotomy for model problems	15
4.4. Full space, $d = 0$	16
4.5. Half space, $d = 1$	16
4.6. Wedges, $d = 2$	18
4.7. Polyhedral cones, $d = 3$	19
4.8. Structure of the admissible generalized eigenvectors	22
5. Continuity properties of the ground energy	24
6. Upper bound for first eigenvalues in polyhedral domains	25
6.1. Change of variables	27

6.2. Construction of quasimodes	28
6.3. Estimation of the quasimodes	30
6.4. Improvement for more regular magnetic fields	35
6.5. Improvement for a straight polyhedron with constant magnetic field	43
7. Lower bound for first eigenvalues	45
Appendix A. Technical lemmas	47
A.1. Gauge transform	47
A.2. Change of variables	49
A.3. Miscelaneous	49
A.4. Cut-off effect	50
Appendix B. Partition of unity suitable for IMS formulas	51
References	56

1. Introduction. Main results

The Schrödinger operator with magnetic field (also called magnetic Laplacian) takes the form

$$(-i\nabla + \mathbf{A})^2$$

where \mathbf{A} is a given vector field that will be assumed to be regular. When set on a domain Ω of \mathbb{R}^n ($n = 2$ or 3) and completed by natural boundary conditions (Neumann), this operator is denoted by $H(\mathbf{A}, \Omega)$. If Ω is bounded with Lipschitz boundary, the form domain of $H(\mathbf{A}, \Omega)$ is the standard Sobolev space $H^1(\Omega)$ and $H(\mathbf{A}, \Omega)$ is positive self-adjoint with compact resolvent. The ground state of $H(\mathbf{A}, \Omega)$ is the eigenpair (λ, ψ)

$$(1.1) \quad \begin{cases} (-i\nabla + \mathbf{A})^2 \psi = \lambda \psi & \text{in } \Omega, \\ (-i\partial_n + \mathbf{n} \cdot \mathbf{A}) \psi = 0 & \text{on } \partial\Omega, \end{cases}$$

associated with the lowest eigenvalue λ . If Ω is simply connected, its eigenvalues only depend on the magnetic field \mathbf{B} defined as

$$(1.2) \quad \mathbf{B} = \text{curl } \mathbf{A}.$$

The eigenvectors corresponding to two different instances of \mathbf{A} for the same \mathbf{B} are deduced from each other by a *gauge transform*.

Introducing a (small) parameter $h > 0$ and setting

$$H_h(\mathbf{A}, \Omega) = (-ih\nabla + \mathbf{A})^2 \quad \text{with Neumann b.c. on } \partial\Omega,$$

we get the relation

$$(1.3) \quad H_h(\mathbf{A}, \Omega) = h^2 H\left(\frac{\mathbf{A}}{h}, \Omega\right)$$

linking the problem with large magnetic field to the semiclassical limit $h \rightarrow 0$. We denote by $\lambda_h(\mathbf{B}, \Omega)$ (or λ_h if no confusion is possible) the smallest eigenvalue of $H_h(\mathbf{A}, \Omega)$ and by ψ_h an associated eigenvector, so that

$$(1.4) \quad \begin{cases} (-ih\nabla + \mathbf{A})^2\psi_h = \lambda_h\psi_h & \text{in } \Omega, \\ (-ih\partial_n + \mathbf{n} \cdot \mathbf{A})\psi_h = 0 & \text{on } \partial\Omega. \end{cases}$$

The behavior of λ_h as $h \rightarrow 0$ clearly provide equivalent information about the lowest eigenvalue of $H(\check{\mathbf{A}}, \Omega)$ when $\check{\mathbf{B}}$ is large, especially in the parametric case when $\check{\mathbf{B}} = B\mathbf{B}$ where the real number B tends to $+\infty$ and \mathbf{B} is a chosen reference magnetic field.

From now on, we consider that \mathbf{B} is fixed. We assume that it is smooth and does not vanish¹ on $\overline{\Omega}$. The question of the semiclassical behavior of λ_h has been considered in many papers for a variety of domains, with constant or variable magnetic fields: Smooth domains [21, 16, 13, 2, 32] and polygons [4, 5, 6] in dimension $n = 2$, and mainly smooth domains [22, 17, 18, 33, 14] in dimension $n = 3$. Until now, three-dimensional non-smooth domains are only addressed in two particular configurations—rectangular cuboids [26] and lenses [28, 31], with special orientation of the (constant) magnetic field. We give more detail about the state of the art in Section 2.

1.1. Asymptotic formulas with remainders. Let us briefly describe our main results in the three-dimensional setting.

Each point \mathbf{x} in the closure of a polyhedral domain Ω is associated with a dilation invariant, tangent open set $\Pi_{\mathbf{x}}$, according to the following cases:

- (1) If \mathbf{x} is an interior point, $\Pi_{\mathbf{x}} = \mathbb{R}^3$,
- (2) If \mathbf{x} belongs to a *face* \mathbf{f} (i.e., a connected component of the smooth part of $\partial\Omega$), $\Pi_{\mathbf{x}}$ is a half-space,
- (3) If \mathbf{x} belongs to an *edge* \mathbf{e} , $\Pi_{\mathbf{x}}$ is an infinite wedge,
- (4) If \mathbf{x} is a *vertex* \mathbf{v} , $\Pi_{\mathbf{x}}$ is an infinite polyhedral cone.

Let $\mathbf{B}_{\mathbf{x}}$ be the magnetic field frozen at \mathbf{x} . Let $E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}})$ be the bottom of the spectrum (ground energy) of the tangent operator $H(\mathbf{A}_{\mathbf{x}}, \Pi_{\mathbf{x}})$ where $\mathbf{A}_{\mathbf{x}}$ is the affine approximation of \mathbf{A} at \mathbf{x} , so that

$$\text{curl } \mathbf{A}_{\mathbf{x}} = \mathbf{B}_{\mathbf{x}}.$$

We introduce the quantity

$$(1.5) \quad \mathcal{E}(\mathbf{B}, \Omega) := \inf_{\mathbf{x} \in \overline{\Omega}} E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}}).$$

In this paper, we prove that this quantity provides the value of the limit of λ_h/h as $h \rightarrow 0$ with some control of the convergence rate as $h \rightarrow 0$, namely

$$(1.6) \quad -Ch^{5/4} \leq \lambda_h(\mathbf{B}, \Omega) - h\mathcal{E}(\mathbf{B}, \Omega) \leq Ch^{5/4},$$

¹Should \mathbf{B} cancel, the situation would be very different, leading to another type of asymptotics [15, 12].

where the constant C is bounded by the norm of \mathbf{A} in $W^{2,\infty}(\Omega)$, as proved in Theorems 6.1 and 7.1. We can also control the constant C by the magnetic field \mathbf{B} as established in Corollaries 6.2 and 7.2. With the point of view of large magnetic fields in the parametric case $\check{\mathbf{B}} = B\mathbf{B}$, (1.6) yields obviously

$$(1.7) \quad -CB^{3/4} \leq \lambda(\check{\mathbf{B}}, \Omega) - B\mathcal{E}(\mathbf{B}, \Omega) \leq CB^{3/4}, \quad \text{as } B \rightarrow +\infty.$$

Note that $B\mathcal{E}(\mathbf{B}, \Omega) = \mathcal{E}(\check{\mathbf{B}}, \Omega)$ by homogeneity (see Lemma A.5).

If the magnetic potential is more regular $\mathbf{A} \in W^{3,\infty}(\overline{\Omega})$, we establish in Theorem 6.3 a better upper bound:

$$(1.8) \quad -Ch^{5/4} \leq \lambda_h(\mathbf{B}, \Omega) - h\mathcal{E}(\mathbf{B}, \Omega) \leq Ch^{4/3},$$

where the constant C is bounded by the norm of \mathbf{A} in $W^{3,\infty}(\Omega)$ and can be controlled by the magnetic field \mathbf{B} as mentioned in Corollary 6.4. Like for (1.7), we deduce the asymptotics for large magnetic fields $\check{\mathbf{B}} = B\mathbf{B}$:

$$(1.9) \quad -CB^{3/4} \leq \lambda(\check{\mathbf{B}}, \Omega) - B\mathcal{E}(\mathbf{B}, \Omega) \leq CB^{2/3}, \quad \text{as } B \rightarrow +\infty.$$

These results are new in this generality. In view of [18, Theorem 1.1] (in the smooth three-dimensional case) the upper bound in (1.9) is optimal. The lower bound coincides with the one obtained in the smooth case in dimensions 2 and 3 when no further assumptions are done. In the literature, improvements of the convergence rates are possible in certain cases when one knows more on $\mathcal{E}(\mathbf{B}, \Omega)$, in particular whether the infimum is attained in some special points.

Our result does not need such extra assumptions, but our proofs have to distinguish cases whether the local ground energies $E(\mathbf{B}_x, \Pi_x)$ are attained or not, and we have to understand the behavior of the function $\mathbf{x} \mapsto E(\mathbf{B}_x, \Pi_x)$ when \mathbf{x} spans the different regions of $\overline{\Omega}$. We have proved very general continuity and semi-continuity properties as described now.

Let \mathfrak{F} be the set of faces \mathbf{f} , \mathfrak{E} the set of edges \mathbf{e} and \mathfrak{V} the set of vertices of Ω . They form a partition of the closure of Ω , called stratification

$$(1.10) \quad \overline{\Omega} = \Omega \cup \left(\bigcup_{\mathbf{f} \in \mathfrak{F}} \mathbf{f} \right) \cup \left(\bigcup_{\mathbf{e} \in \mathfrak{E}} \mathbf{e} \right) \cup \left(\bigcup_{\mathbf{v} \in \mathfrak{V}} \mathbf{v} \right).$$

The sets Ω , \mathbf{f} , \mathbf{e} and \mathbf{v} are called the strata of $\overline{\Omega}$, compare with [23] and [25, Ch. 9]. We denote them by \mathbf{t} and their set by \mathfrak{T} . For each stratum \mathbf{t} , let us denote by $\Lambda_{\mathbf{t}}$ the function

$$(1.11) \quad \Lambda_{\mathbf{t}} : \mathbf{t} \ni \mathbf{x} \mapsto E(\mathbf{B}_x, \Pi_x).$$

We will show the following facts

- a) The function $\mathbf{x} \mapsto E(\mathbf{B}_x, \Pi_x)$ is lower semi-continuous on $\overline{\Omega}$.
- b) For each stratum $\mathbf{t} \in \mathfrak{T}$, the function $\Lambda_{\mathbf{t}}$ is continuous on \mathbf{t} and can be continuously extended to the closure $\bar{\mathbf{t}}$ of \mathbf{t} . Moreover, for each $\mathbf{x}_0 \in \bar{\mathbf{t}}$, $\Lambda_{\mathbf{t}}(\mathbf{x}_0)$ is the bottom of the spectrum $E(\mathbf{B}_x, \Pi_x)$ of a tangent magnetic operator $H(\mathbf{A}_x, \Pi_x)$ associated with a *singular chain* \mathbb{X} originating at \mathbf{x}_0 .

As a consequence, the infimum determining the limit $\mathcal{E}(\mathbf{B}, \Omega)$ in (1.5) is a minimum

$$(1.12) \quad \mathcal{E}(\mathbf{B}, \Omega) = \min_{\mathbf{x} \in \overline{\Omega}} E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}}).$$

1.2. Contents of the paper. In Section 2 we place our results in the framework of existing literature for dimensions 2 and 3. In Section 3 we introduce the wider class of corner domains, alongside with their tangent cones and singular chains. We particularize these notions in the case of three-dimensional polyhedral domains. In Section 4 we introduce and classify magnetic model problems on tangent cones (taxonomy) and extract from the literature related facts. We show that to each point $\mathbf{x} \in \overline{\Omega}$ is associated a singular chain \mathbb{X} originating at \mathbf{x} for which the tangent operator $H(\mathbf{A}_{\mathbb{X}}, \Pi_{\mathbb{X}})$ possesses *admissible generalized eigenvectors* with energy $E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}})$. In Section 5 we prove the semi-continuity and continuity properties of the functions $\mathbf{x} \mapsto E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}})$ on $\overline{\Omega}$ and its strata. In Section 6 we prove the upper bounds $\lambda_h(\mathbf{B}, \Omega) \leq h\mathcal{E}(\mathbf{B}, \Omega) + Ch^\kappa$, with $\kappa = 5/4$ or $\kappa = 4/3$ according to the regularity of \mathbf{A} , by a construction of quasimodes based on admissible generalized eigenvectors for tangent problems. In Section 7 we prove the lower bound $h\mathcal{E}(\mathbf{B}, \Omega) - Ch^{5/4} \leq \lambda_h(\mathbf{B}, \Omega)$ by a classical IMS formula.

1.3. Notations. For a generic (unbounded) self-adjoint operator L we denote by $\text{Dom}(L)$ its domain and $\mathfrak{S}(L)$ its spectrum. Likewise the domain of a quadratic form q is denoted by $\text{Dom}(q)$.

Domains as open simply connected subsets of \mathbb{R}^n are denoted by \mathcal{O} if they are generic, Π if they are invariant by dilatation (cones) and Ω if they are bounded.

In this paper, the quadratic forms of interest are those associated with magnetic Laplacians, namely, for a positive constant h , a smooth magnetic potential \mathbf{A} , and a generic domain \mathcal{O}

$$(1.13) \quad q_h[\mathbf{A}, \mathcal{O}](f) := \int_{\mathcal{O}} (-ih\nabla + \mathbf{A})f \cdot \overline{(-ih\nabla + \mathbf{A})f} \, d\mathbf{x},$$

and its domain

$$(1.14) \quad \text{Dom}(q_h[\mathbf{A}, \mathcal{O}]) = \{f \in L^2(\mathcal{O}), (-ih\nabla + \mathbf{A})f \in L^2(\mathcal{O})\}.$$

For a bounded domain Ω , $\text{Dom}(q_h[\mathbf{A}, \Omega])$ coincides with $H^1(\Omega)$. For $h = 1$, we omit the index h , denoting the quadratic form by $q[\mathbf{A}, \mathcal{O}]$.

In relation with changes of variables, we will also use the more general form with metric:

$$(1.15) \quad q_h[\mathbf{A}, \mathcal{O}, G](\psi) = \int_{\mathcal{O}} (-ih\nabla + \mathbf{A})\psi \cdot G(\overline{(-ih\nabla + \mathbf{A})\psi}) |G|^{-1/2} \, d\mathbf{x},$$

where G is a smooth function with values in 3×3 positive symmetric matrices and $|G| = \det G$. Its domain is (see [18, §5] for more details)

$$\text{Dom}(q_h[\mathbf{A}, \mathcal{O}, G]) = \{\psi \in L_G^2(\mathcal{O}), G^{1/2}(-ih\nabla + \mathbf{A})\psi \in L_G^2(\mathcal{O})\},$$

where $L_G^2(\mathcal{O})$ is the space of the square-integrable functions for the weight $|G|^{-1/2}$ and $G^{1/2}$ is the square root of the matrix G .

The domain of the magnetic Laplacian with Neumann boundary conditions on the set \mathcal{O} is

$$(1.16) \quad \text{Dom}(H_h(\mathbf{A}, \mathcal{O})) = \{\psi \in L^2(\mathcal{O}), \\ (-ih\nabla + \mathbf{A})^2\psi \in L^2(\mathcal{O}) \text{ and } (-ih\partial_n + \mathbf{n} \cdot \mathbf{A})\psi = 0 \text{ on } \partial\mathcal{O}\}.$$

We will also use the space of the functions which are *locally*² in the domain of $H_h(\mathbf{A}, \mathcal{O})$:

$$(1.17) \quad \text{Dom}_{\text{loc}}(H_h(\mathbf{A}, \mathcal{O})) := \{\psi \in H_{\text{loc}}^1(\overline{\mathcal{O}}), \\ (-ih\nabla + \mathbf{A})^2\psi \in H_{\text{loc}}^0(\overline{\mathcal{O}}) \text{ and } (-ih\partial_n + \mathbf{n} \cdot \mathbf{A})\psi = 0 \text{ on } \partial\mathcal{O}\}.$$

When $h = 1$, we omit the index h in (1.16) and (1.17).

2. State of the art

In this section we review the literature about the semiclassical limit for the first eigenvalue $\lambda_h(\mathbf{B}, \Omega)$ of the magnetic Laplacian $H_h(\mathbf{A}, \Omega)$ depending on the dimension $n \in \{2, 3\}$ and the geometry of the domain Ω .

2.1. Dimension 2. In dimension $n = 2$, two classes of domains are considered: the domains with a regular boundary and the polygonal domains.

2.1.1. Regular domains. Let $\Omega \subset \mathbb{R}^2$ be a regular domain and B be a regular non-vanishing scalar magnetic field on $\overline{\Omega}$. To each $\mathbf{x} \in \overline{\Omega}$ is associated a tangent problem. According to whether \mathbf{x} is an interior point or a boundary point, the tangent problem is the magnetic Laplacian on the plane \mathbb{R}^2 or the half-plane $\Pi_{\mathbf{x}}$ tangent to Ω at \mathbf{x} , with the constant magnetic field $B_{\mathbf{x}} \equiv B(\mathbf{x})$. The associated spectral quantities $E(B_{\mathbf{x}}, \mathbb{R}^2)$ and $E(B_{\mathbf{x}}, \Pi_{\mathbf{x}})$ are respectively equal to $|B_{\mathbf{x}}|$ and $|B_{\mathbf{x}}|\Theta_0$ where $\Theta_0 := E(1, \mathbb{R}_+^2)$ is a universal constant whose value is close to 0.59 (see [34]). With the quantities

$$(2.1) \quad b = \inf_{\mathbf{x} \in \Omega} |B(\mathbf{x})| \quad \text{and} \quad b' = \inf_{\mathbf{x} \in \partial\Omega} |B(\mathbf{x})|,$$

we find

$$\mathcal{E}(B, \Omega) = \min(b, b'\Theta_0).$$

In this generality, the asymptotic limit

$$(2.2) \quad \lim_{h \rightarrow 0} \frac{\lambda_h(B, \Omega)}{h} = \mathcal{E}(B, \Omega)$$

is proved by Lu and Pan in [21]. Improvements of this result depend on the geometry and the variation of the magnetic field as we describe now.

²Here $H_{\text{loc}}^m(\overline{\mathcal{O}})$ denotes for $m = 0, 1$ the space of functions which are in $H^m(\mathcal{O} \cap \mathcal{B})$ for any ball \mathcal{B} .

- *Constant magnetic field.* If the magnetic field is constant and normalized to 1, then $\mathcal{E}(B, \Omega) = \Theta_0$. The following estimate is proved by Helffer and Morame:

$$\exists C > 0, \quad -Ch^{3/2} \leq \lambda_h(1, \Omega) - h\Theta_0 \leq Ch^{3/2},$$

for h small enough [16, §10], while the upper bound was already given by Bernoff and Sternberg [3]. This result is improved in [16, §11] in which a two-term asymptotics is proved, showing that a remainder in $O(h^{3/2})$ is optimal. Under the additional assumption that the curvature of the boundary admits a unique and non-degenerate maximum, a complete expansion of $\lambda_h(1, \Omega)$ is provided by Fournais and Helffer [13].

- *Variable magnetic field.* Here we recall results from [16, §9] for variable magnetic fields (we use the notation (2.1))

$$\begin{aligned} \text{If } b < \Theta_0 b', \quad & \exists C > 0, & |\lambda_h(B, \Omega) - hb| &\leq Ch^2, \\ \text{If } b > \Theta_0 b', \quad & \exists C > 0, & -Ch^{5/4} \leq \lambda_h(B, \Omega) - h\Theta_0 b' &\leq Ch^{3/2}, \\ \text{If } b = \Theta_0 b', \quad & \exists C > 0, & -Ch^{5/4} \leq \lambda_h(B, \Omega) - hb &\leq Ch^2, \end{aligned}$$

for h small enough. Under further assumptions, more precise asymptotic expansions are given by Aramaki [2] and Raymond [32]. They show that the upper bounds above are sharp.

2.1.2. *Polygonal domains.* Let Ω be a curvilinear polygon and let \mathfrak{V} be the (finite) set of its vertices. In this case, new model operators appear on infinite sectors $\Pi_{\mathbf{x}}$ tangent to Ω at vertices $\mathbf{x} \in \mathfrak{V}$. By homogeneity $E(B_{\mathbf{x}}, \Pi_{\mathbf{x}}) = |B(\mathbf{x})|E(1, \Pi_{\mathbf{x}})$ and by rotation invariance, $E(1, \Pi_{\mathbf{x}})$ only depends on the opening $\alpha(\mathbf{x})$ of the sector $\Pi_{\mathbf{x}}$. Let \mathcal{S}_{α} be a model sector of opening $\alpha \in (0, 2\pi)$. Then

$$\mathcal{E}(B, \Omega) = \min(b, b'\Theta_0, \min_{\mathbf{x} \in \mathfrak{V}} |B(\mathbf{x})| E(1, \mathcal{S}_{\alpha(\mathbf{x})})) .$$

In [4, §11], it is proved that

$$\exists C > 0, \quad -Ch^{5/4} \leq \lambda_h(B, \Omega) - h\mathcal{E}(B, \Omega) \leq Ch^{9/8}.$$

This estimate can be improved under the assumption that

$$(2.3) \quad \mathcal{E}(B, \Omega) < \min(b, b'\Theta_0),$$

which means that at least one of the corners makes the energy lower than in the regular case: The asymptotic expansions provided in [5] then yield the sharp estimates

$$\exists C > 0, \quad |\lambda_h(B, \Omega) - h\mathcal{E}(B, \Omega)| \leq Ch^{3/2} .$$

From [19, 4] follows that for all $\alpha \in (0, \frac{\pi}{2}]$ there holds

$$(2.4) \quad E(1, \mathcal{S}_{\alpha}) < \Theta_0.$$

Therefore condition (2.3) holds for constant magnetic fields as soon as there is an angle opening $\alpha_{\mathbf{x}} \leq \frac{\pi}{2}$. Finite element computations by Galerkin projection as presented in [6] suggest that (2.4) still holds for all $\alpha \in (0, \pi)$. Let us finally mention that if Ω has straight

sides and B is constant, the convergence of $\lambda_h(B, \Omega)$ to $h\mathcal{E}(B, \Omega)$ is exponential: Their difference is bounded by $C \exp(-\beta h^{-1/2})$ for suitable positive constants C and β (see [5]).

2.2. Dimension 3. In dimension $n = 3$ we still distinguish the regular and singular domains.

2.2.1. Regular domains. Here $\Omega \subset \mathbb{R}^3$ is assumed to be regular. For a continuous magnetic field \mathbf{B} it is known ([22] and [17]) that (2.2) holds. In that case

$$\mathcal{E}(\mathbf{B}, \Omega) = \min \left(\inf_{\mathbf{x} \in \Omega} |\mathbf{B}(\mathbf{x})|, \inf_{\mathbf{x} \in \partial\Omega} |\mathbf{B}(\mathbf{x})| \sigma(\theta(\mathbf{x})) \right),$$

where $\theta(\mathbf{x}) \in [0, \frac{\pi}{2}]$ denotes the unoriented angle between the magnetic field and the boundary at the point $\mathbf{x} \in \partial\Omega$, and the quantity $\sigma(\theta)$ is the bottom of the spectrum of a model problem, see Section 4. Let us simply mention that σ is increasing on $[0, \frac{\pi}{2}]$ and that $\sigma(0) = \Theta_0$, $\sigma(\pi/2) = 1$.

- *Constant magnetic field.* Here the magnetic field \mathbf{B} is assumed to be constant and unitary. There exists a non-empty set Σ of $\partial\Omega$ on which $\mathbf{B}(\mathbf{x})$ is tangent to the boundary. In that case we have

$$\mathcal{E}(\mathbf{B}, \Omega) = \Theta_0 .$$

Theorem 1.1 of [18] states that

$$\exists C > 0, \quad |\lambda_h(\mathbf{B}, \Omega) - h\Theta_0| \leq Ch^{4/3},$$

for h small enough. Under some extra assumptions on Σ , Theorem 1.2 of [18] yields a two-term asymptotics for $\lambda_h(\mathbf{B}, \Omega)$ showing the optimality of the previous estimate.

- *Variable magnetic field.* Let \mathbf{B} be a smooth non-vanishing magnetic field. There holds [14, Theorem 9.1.1]

$$\exists C > 0, \quad -Ch^{5/4} \leq \lambda_h(\mathbf{B}, \Omega) - h\mathcal{E}(\mathbf{B}, \Omega) \leq Ch^{5/4} .$$

The proof of this result was already sketched in [22]. In [18, Remark 6.2], the upper bound is improved to $O(h^{4/3})$.

Under the following two extra assumptions

- The inequality $\inf_{\mathbf{x} \in \partial\Omega} |\mathbf{B}(\mathbf{x})| \sigma(\theta(\mathbf{x})) < \inf_{\mathbf{x} \in \Omega} |\mathbf{B}(\mathbf{x})|$ holds,
- The function $\mathbf{x} \mapsto |\mathbf{B}(\mathbf{x})| \sigma(\theta(\mathbf{x}))$ reaches its minimum at a point \mathbf{x}_0 where \mathbf{B} is neither normal nor tangent to the boundary,

a two-term asymptotics is valid [33], providing the sharp estimate:

$$\exists C > 0, \quad |\lambda_h(\mathbf{B}, \Omega) - h\mathcal{E}(\mathbf{B}, \Omega)| \leq Ch^{3/2} .$$

2.2.2. Singular domains. Until now, two examples of non-smooth domains have been addressed in the literature. In both cases, the magnetic field \mathbf{B} is assumed to be constant.

- *Rectangular cuboids.* The case where Ω is a rectangular cuboid (i.e., the product of three bounded intervals) is considered by Pan [26]: The asymptotic limit (2.2) holds for such a domain and there exists a vertex $\mathbf{v} \in \mathfrak{V}$ such that $\mathcal{E}(\mathbf{B}, \Omega) = E(\mathbf{B}, \Pi_{\mathbf{v}})$. Moreover, in the case where the magnetic field is tangent to a face but is not tangent to any edge, there holds

$$E(\mathbf{B}, \Pi_{\mathbf{v}}) < \inf_{\mathbf{x} \in \overline{\Omega} \setminus \mathfrak{V}} E(\mathbf{B}, \Pi_{\mathbf{x}})$$

and eigenfunctions associated to $\lambda_h(\mathbf{B}, \Omega)$ concentrate near corners as $h \rightarrow 0$.

- *Lenses.* The case where Ω has the shape of a lens is treated in [28] and [31]. The domain Ω is supposed to have two faces separated by an edge \mathbf{e} that is a regular loop contained in the plane $x_3 = 0$. The magnetic field considered is $\mathbf{B} = (0, 0, 1)$.

It is proved in [28] that, if the opening angle α of the lens is constant and $\leq 0.38\pi$,

$$\inf_{\mathbf{x} \in \mathbf{e}} E(\mathbf{B}, \Pi_{\mathbf{x}}) < \inf_{\mathbf{x} \in \overline{\Omega} \setminus \mathbf{e}} E(\mathbf{B}, \Pi_{\mathbf{x}})$$

and that the asymptotic limit (2.2) holds with the following estimate:

$$\exists C > 0, \quad |\lambda_h(\mathbf{B}, \Omega) - h\mathcal{E}(\mathbf{B}, \Omega)| \leq Ch^{5/4}.$$

When the opening angle of the lens is variable and under some non-degeneracy hypotheses, a complete eigenvalue asymptotics is obtained in [31] resulting into the optimal estimate

$$\exists C > 0, \quad |\lambda_h(\mathbf{B}, \Omega) - h\mathcal{E}(\mathbf{B}, \Omega)| \leq Ch^{3/2}.$$

3. Polyhedral domains and their singular chains

For the sake of completeness and for ease of further discussion, in the same spirit as in [10, Section 2], we introduce here a recursive definition of two intertwining classes of domains

- \mathfrak{P}_n , a class of infinite open cones in \mathbb{R}^n .
- $\mathfrak{D}(M)$, a class of bounded connected open subsets of a smooth manifold without boundary—actually, $M = \mathbb{R}^n$ or $M = \mathbb{S}^n$, with \mathbb{S}^n the unit sphere of \mathbb{R}^{n+1} ,

3.1. **Domains and tangent cones.** We call a *cone* any open subset Π of \mathbb{R}^n satisfying

$$\forall \rho > 0 \text{ and } \mathbf{x} \in \Pi, \quad \rho\mathbf{x} \in \Pi,$$

and the *section* of the cone Π is its subset $\Pi \cap \mathbb{S}^{n-1}$. Note that $\mathbb{S}^0 = \{-1, 1\}$.

Initialization: \mathfrak{P}_0 has one element, $\{0\}$. $\mathfrak{D}(\mathbb{S}^0)$ is formed by all subsets of \mathbb{S}^0 .

Recurrence: For $n \geq 1$,

- (1) $\Pi \in \mathfrak{P}_n$ if and only if the section of Π belongs to $\mathfrak{D}(\mathbb{S}^{n-1})$,

- (2) $\Omega \in \mathfrak{D}(M)$ if and only if for any $\mathbf{x} \in \overline{\Omega}$, there exists a cone $\Pi_{\mathbf{x}} \in \mathfrak{P}_n$ and a local \mathcal{C}^∞ diffeomorphism $U_{\mathbf{x}}$ which maps a neighborhood $\mathcal{U}_{\mathbf{x}}$ of \mathbf{x} in M onto a neighborhood $\mathcal{V}_{\mathbf{x}}$ of $\mathbf{0}$ in \mathbb{R}^n and such that

$$(3.1) \quad U_{\mathbf{x}}(\mathcal{U}_{\mathbf{x}} \cap \Omega) = \mathcal{V}_{\mathbf{x}} \cap \Pi_{\mathbf{x}} \quad \text{and} \quad U_{\mathbf{x}}(\mathcal{U}_{\mathbf{x}} \cap \partial\Omega) = \mathcal{V}_{\mathbf{x}} \cap \partial\Pi_{\mathbf{x}}.$$

We assume without restriction that the differential of $U_{\mathbf{x}}$ at the point \mathbf{x} is the identity matrix \mathbb{I}_n . The cone $\Pi_{\mathbf{x}}$ is said to be tangent to Ω at \mathbf{x} .

Examples:

- The elements of \mathfrak{P}_1 are \mathbb{R} , \mathbb{R}_+ and \mathbb{R}_- .
- The elements of $\mathfrak{D}(\mathbb{S}^1)$ are \mathbb{S}^1 and all open intervals $\mathcal{I} \subset \mathbb{S}^1$ such that $\overline{\mathcal{I}} \neq \mathbb{S}^1$.
- The elements of \mathfrak{P}_2 are \mathbb{R}^2 and all sectors with opening $\alpha \in (0, 2\pi)$, including half-spaces ($\alpha = \pi$).
- The elements of $\mathfrak{D}(\mathbb{R}^2)$ are curvilinear polygons with piecewise non-tangent smooth sides. Note that corner angles do not take the values 0 or 2π , and that $\mathfrak{D}(\mathbb{R}^2)$ includes smooth domains.

Definition 3.1. Let \mathfrak{O}_n denote the group of orthogonal linear transformations of \mathbb{R}^n .

- a) We say that a cone Π is *equivalent* to another cone Π' and denote

$$\Pi \equiv \Pi'$$

if there exists $\underline{U} \in \mathfrak{O}_n$ such that $\underline{U}\Pi = \Pi'$.

- b) Let $\Pi \in \mathfrak{P}_n$. If Π is equivalent to $\mathbb{R}^{n-d} \times \Gamma$ with $\Gamma \in \mathfrak{P}_d$ and d is minimal for such an equivalence, Γ is said to be a *minimal reduced cone* associated with Π .
- c) Let $\mathbf{x} \in \overline{\Omega}$ and let $\Pi_{\mathbf{x}}$ be its tangent cone. We denote by $d_0(\mathbf{x})$ the dimension of the minimal reduced cone associated with $\Pi_{\mathbf{x}}$.

3.2. Recursive definition of singular chains. A singular chain $\mathbb{X} = (\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_\nu) \in \mathfrak{C}(\Omega)$ (with a natural number ν) is a finite collection of points according to the following recursive definition.

Initialization: $\mathbf{x}_0 \in \overline{\Omega}$,

- Let $C_{\mathbf{x}_0}$ be the tangent cone to Ω at \mathbf{x}_0 (here $C_{\mathbf{x}_0} = \Pi_{\mathbf{x}_0}$),
- Let $\Gamma_{\mathbf{x}_0} \in \mathfrak{P}_{d_0}$ be its minimal reduced cone: $C_{\mathbf{x}_0} = \underline{U}_0(\mathbb{R}^{n-d_0} \times \Gamma_{\mathbf{x}_0})$.
- Alternative:
 - If $\nu = 0$, stop here.
 - If $\nu > 0$, then $d_0 > 0$ and let $\Omega_{\mathbf{x}_0} \in \mathfrak{D}(\mathbb{S}^{d_0-1})$ be the section of $\Gamma_{\mathbf{x}_0}$

Recurrence: $\mathbf{x}_j \in \overline{\Omega_{\mathbf{x}_0, \dots, \mathbf{x}_{j-1}}} \in \mathfrak{D}(\mathbb{S}^{d_{j-1}-1})$. If $d_{j-1} = 1$, stop here ($\nu = j$). If not:

- Let $C_{\mathbf{x}_0, \dots, \mathbf{x}_j}$ be the tangent cone to $\Omega_{\mathbf{x}_0, \dots, \mathbf{x}_{j-1}}$ at \mathbf{x}_j ,
- Let $\Gamma_{\mathbf{x}_0, \dots, \mathbf{x}_j} \in \mathfrak{P}_{d_j}$ be its minimal reduced cone: $C_{\mathbf{x}_0, \dots, \mathbf{x}_j} = \underline{U}_j(\mathbb{R}^{d_{j-1}-1-d_j} \times \Gamma_{\mathbf{x}_0, \dots, \mathbf{x}_j})$.

- Alternative:
 - If $j = \nu$, stop here.
 - If $j < \nu$, then $d_j > 0$ and let $\Omega_{\mathbf{x}_0, \dots, \mathbf{x}_j} \in \mathfrak{D}(\mathbb{S}^{d_j-1})$ be the section of $\Gamma_{\mathbf{x}_0, \dots, \mathbf{x}_j}$.

Note that $n \geq d_0 > d_1 > \dots > d_\nu$. Hence $\nu \leq n$. Note also that for $\nu = 0$, we obtain the trivial one element chain (\mathbf{x}_0) for any $\mathbf{x}_0 \in \overline{\Omega}$.

While $\mathfrak{C}(\Omega)$ is the set of all singular chains, for any $\mathbf{x} \in \overline{\Omega}$, we denote by $\mathfrak{C}_{\mathbf{x}}(\Omega)$ the subset of chains originating at \mathbf{x} , i.e., the set of chains $\mathbb{X} = (\mathbf{x}_0, \dots, \mathbf{x}_\nu)$ with $\mathbf{x}_0 = \mathbf{x}$. Note that the one element chain (\mathbf{x}) belongs to $\mathfrak{C}_{\mathbf{x}}(\Omega)$. We also set

$$(3.2) \quad \mathfrak{C}_{\mathbf{x}}^*(\Omega) = \{\mathbb{X} \in \mathfrak{C}_{\mathbf{x}}(\Omega), \nu > 0\} = \mathfrak{C}_{\mathbf{x}}(\Omega) \setminus \{(\mathbf{x})\}.$$

We set finally, with the notation $\langle \mathbf{y} \rangle$ for the vector space generated by \mathbf{y} ,

$$(3.3) \quad \Pi_{\mathbb{X}} = \begin{cases} C_{\mathbf{x}_0} = \Pi_{\mathbf{x}_0} & \text{if } \nu = 0, \\ \bigcup_0 (\mathbb{R}^{n-d_0} \times \langle \mathbf{x}_1 \rangle \times C_{\mathbf{x}_0, \mathbf{x}_1}) & \text{if } \nu = 1, \\ \bigcup_0 (\mathbb{R}^{n-d_0} \times \langle \mathbf{x}_1 \rangle \times \dots \times \bigcup_{\nu-1} (\mathbb{R}^{d_{\nu-2}-1-d_{\nu-1}} \times \langle \mathbf{x}_\nu \rangle \times C_{\mathbf{x}_0, \dots, \mathbf{x}_\nu}) \dots) & \text{if } \nu \geq 2. \end{cases}$$

Note that if $d_\nu = 0$, the cone $C_{\mathbf{x}_0, \dots, \mathbf{x}_\nu}$ coincides with $\mathbb{R}^{d_{\nu-1}-1}$, leading to $\Pi_{\mathbb{X}} = \mathbb{R}^n$.

Definition 3.2. Let $\mathbb{X} = (\mathbf{x}_0, \dots, \mathbf{x}_\nu)$ and $\mathbb{X}' = (\mathbf{x}'_0, \dots, \mathbf{x}'_\nu)$ be two chains in $\mathfrak{C}(\Omega)$. We say that \mathbb{X} is equivalent to \mathbb{X}' if $\mathbf{x}_0 = \mathbf{x}'_0$ and $\Pi_{\mathbb{X}} = \Pi_{\mathbb{X}'}$.

Special subsets of $\overline{\Omega}$: For $d \in \{0, \dots, n\}$, let

$$(3.4) \quad \mathfrak{A}_d(\Omega) = \{\mathbf{x} \in \overline{\Omega}, d_0(\mathbf{x}) = d\}.$$

The strata of $\overline{\Omega}$ are the connected components of $\mathfrak{A}_d(\Omega)$, for $d \in \{0, \dots, n\}$. They are denoted by \mathbf{t} and their set by \mathfrak{T} .

Examples:

- $\mathfrak{A}_0(\Omega)$ coincides with Ω .
- $\mathfrak{A}_1(\Omega)$ is the subset of $\partial\Omega$ of the regular points of the boundary.
- If $n = 2$, $\mathfrak{A}_2(\Omega)$ is the set of corners.
- If $n = 3$, $\mathfrak{A}_2(\Omega)$ is the set of edge points.
- If $n = 3$, $\mathfrak{A}_3(\Omega)$ is the set of corners.

3.3. Polyhedral domains. Polyhedral domains and polyhedral cones form subclasses of $\mathfrak{D}(M)$ and \mathfrak{P}_n , denoted by $\overline{\mathfrak{D}}(M)$ and $\overline{\mathfrak{P}}_n$, respectively:

- The cone $\Pi \in \mathfrak{P}_n$ is a polyhedral cone if its boundary is contained in a finite union of (hyper)surfaces. We write $\Pi \in \overline{\mathfrak{P}}_n$.
- The domain $\Omega \in \mathfrak{D}(M)$ is a polyhedral domain if all its tangent cones $\Pi_{\mathbf{x}}$ are polyhedral. We write $\Omega \in \overline{\mathfrak{D}}(M)$.

This allows to make precise the definition of faces, edges and corners in dimension 3, in connection with singular chains.

- (1) Interior point $\mathbf{x} \in \Omega$. Only one chain in $\mathfrak{C}_{\mathbf{x}}(\Omega)$: $\mathbb{X} = (\mathbf{x})$.
- (2) The faces \mathbf{f} are the connected components of $\mathfrak{A}_1(\Omega)$. The set of faces is denoted by \mathfrak{F} . Let \mathbf{x} belong to a face. There are two chains in $\mathfrak{C}_{\mathbf{x}}(\Omega)$:
 - (a) $\mathbb{X} = (\mathbf{x})$ with $\Pi_{\mathbb{X}} = \Pi_{\mathbf{x}}$, the tangent half-space. $\Pi_{\mathbb{X}} \equiv \mathbb{R}^2 \times \mathbb{R}_+$.
 - (b) $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1)$ where $\mathbf{x}_1 = 1$ is the only element in $\mathbb{R}_+ \cap \mathbb{S}^0$. Thus $\Pi_{\mathbb{X}} = \mathbb{R}^3$.
- (3) The edges \mathbf{e} are the connected components of $\mathfrak{A}_2(\Omega)$. The set of edges is denoted by \mathfrak{E} . Let \mathbf{x} belong to an edge. There are three types of chains in $\mathfrak{C}_{\mathbf{x}}(\Omega)$:
 - (a) $\mathbb{X} = (\mathbf{x})$ with $\Pi_{\mathbb{X}} = \Pi_{\mathbf{x}}$, the tangent wedge (which is not a half-plane). The reduced cone of $\Pi_{\mathbf{x}}$ is a sector $\Gamma_{\mathbf{x}}$ the section of which is an interval $\mathcal{I}_{\mathbf{x}} \subset \mathbb{S}^1$.
 - (b) $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1)$ where $\mathbf{x}_1 \in \overline{\mathcal{I}_{\mathbf{x}}}$.
 - (i) If \mathbf{x}_1 is interior to $\mathcal{I}_{\mathbf{x}}$, $\Pi_{\mathbb{X}} = \mathbb{R}^3$. No further chain.
 - (ii) If \mathbf{x}_1 is a boundary point of $\mathcal{I}_{\mathbf{x}}$, $\Pi_{\mathbb{X}}$ is a half-space, containing one of the two faces $\partial^{\pm}\Pi_{\mathbf{x}}$ of the wedge $\Pi_{\mathbf{x}}$.
 - (c) $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1, \mathbf{x}_2)$ where $\mathbf{x}_1 \in \partial\mathcal{I}_{\mathbf{x}}$, $\mathbf{x}_2 = 1$ and $\Pi_{\mathbb{X}} = \mathbb{R}^3$.

To sum up, there are 4 equivalence classes in $\mathfrak{C}_{\mathbf{x}}(\Omega)$ in the case of an edge point \mathbf{x} :

- $\mathbb{X} = (\mathbf{x})$
- $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1^{\pm})$ with $\{\mathbf{x}_1^-, \mathbf{x}_1^+\} = \partial\mathcal{I}_{\mathbf{x}}$
- $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1^{\circ})$ with \mathbf{x}_1° any chosen point in $\mathcal{I}_{\mathbf{x}}$.

- (4) The corners \mathbf{v} are the connected components of $\mathfrak{A}_3(\Omega)$. The set of corners is denoted by \mathfrak{V} . There are four types of chains in $\mathfrak{C}_{\mathbf{x}}(\Omega)$:
 - (a) $\mathbb{X} = (\mathbf{x})$ with $\Pi_{\mathbb{X}} = \Pi_{\mathbf{x}}$, the tangent cone (that is not a wedge). It coincides with its reduced cone. Its section $\Omega_{\mathbf{x}}$ is a polygonal domain in \mathbb{S}^2 .
 - (b) $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1)$ where $\mathbf{x}_1 \in \overline{\Omega_{\mathbf{x}}}$.
 - (i) If \mathbf{x}_1 is interior to $\Omega_{\mathbf{x}}$, $\Pi_{\mathbb{X}} = \mathbb{R}^3$. No further chain.
 - (ii) If \mathbf{x}_1 is in a side of $\Omega_{\mathbf{x}}$, $\Pi_{\mathbb{X}}$ is a half-space, containing one of the faces of the cone $\Pi_{\mathbf{x}}$.
 - (iii) If \mathbf{x}_1 is a corner of $\Omega_{\mathbf{x}}$, $\Pi_{\mathbb{X}}$ is a wedge. Its edge contains one of the edges of $\Pi_{\mathbf{x}}$.
 - (c) $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1, \mathbf{x}_2)$ where $\mathbf{x}_1 \in \partial\Omega_{\mathbf{x}}$
 - (i) If \mathbf{x}_1 is in a side of $\Omega_{\mathbf{x}}$, $\mathbf{x}_2 = 1$, $\Pi_{\mathbb{X}} = \mathbb{R}^3$. No further chain.
 - (ii) If \mathbf{x}_1 is a corner of $\Omega_{\mathbf{x}}$, $C_{\mathbf{x}, \mathbf{x}_1}$ is plane sector, and $\mathbf{x}_2 \in \overline{\mathcal{I}_{\mathbf{x}, \mathbf{x}_1}}$ where the interval $\mathcal{I}_{\mathbf{x}, \mathbf{x}_1}$ is its section. If \mathbf{x}_2 is an interior point, then $\Pi_{\mathbb{X}} = \mathbb{R}^3$.
 - (d) $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$ where \mathbf{x}_1 is a corner of $\Omega_{\mathbf{x}}$, $\mathbf{x}_2 \in \partial\mathcal{I}_{\mathbf{x}, \mathbf{x}_1}$ and $\mathbf{x}_3 = 1$. Then $\Pi_{\mathbb{X}} = \mathbb{R}^3$.

Let \mathbf{x}_1^j , $1 \leq j \leq N$, be the corners of $\Omega_{\mathbf{x}}$, and \mathbf{f}_1^j , $1 \leq j \leq N$, be its sides (notice that there are as many corners as sides). There are $2N + 2$ equivalence classes in $\mathfrak{C}_{\mathbf{x}}(\Omega)$:

- $\mathbb{X} = (\mathbf{x})$ (vertex)
- $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1^j)$ with $1 \leq j \leq N$ (edge points limit)

- $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1^{\circ j})$ with $\mathbf{x}_1^{\circ j}$ any chosen point inside \mathbf{f}_1^j (face points limit)
- $\mathbb{X} = (\mathbf{x}, \mathbf{x}_1^\circ)$ with \mathbf{x}_1° any chosen point in $\Omega_{\mathbf{x}}$ (interior points limit).

Remark 3.3. For polyhedral domains Ω , it is a consequence of the description above that chains $(\mathbf{x}_0, \mathbf{x}_1)$ of length 2 are enough to describe all equivalence classes of the set of chains $\mathfrak{C}_{\mathbf{x}_0}^*(\Omega)$ (3.2). This does not hold anymore if general corner domains are considered. Besides, the notion of equivalence classes as introduced in Definition 3.2 is sufficient for the analysis of operators $H_h(\mathbf{A}, \Omega)$ in the case of magnetic fields \mathbf{B} smooth in Cartesian variables. Should \mathbf{B} be smooth in polar variables only, the whole hierarchy of singular chains would be needed.

4. Taxonomy of model problems

4.1. Tangent and model operators. We recall that \mathbf{A} is a magnetic potential associated with the magnetic field \mathbf{B} on the polyhedral domain $\Omega \in \overline{\mathfrak{D}}(\mathbb{R}^3)$. For each singular chain $\mathbb{X} = (\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_\nu) \in \mathfrak{C}(\Omega)$ we set

$$(4.1) \quad \mathbf{B}_{\mathbb{X}} = \mathbf{B}(\mathbf{x}_0) \quad \text{and} \quad \mathbf{A}_{\mathbb{X}} = \mathbf{A}(\mathbf{x}_0) + (\nabla \mathbf{A}(\mathbf{x}_0))(\mathbf{x} - \mathbf{x}_0).$$

We have obviously

$$\text{curl } \mathbf{A}_{\mathbb{X}} = \mathbf{B}_{\mathbb{X}},$$

so that the tangent magnetic operator $H(\mathbf{A}_{\mathbb{X}}, \Pi_{\mathbb{X}})$ and its ground energy $E(\mathbf{B}_{\mathbb{X}}, \Pi_{\mathbb{X}})$ make sense. By homogeneity, there holds for tangent problems (cf. Lemma A.5)

Lemma 4.1. *Let $\Pi \in \mathfrak{P}_3$ be a cone, \mathbf{B} be a constant magnetic field with norm $b > 0$. There holds*

$$(4.2) \quad E(\mathbf{B}, \Pi) = b E\left(\frac{\mathbf{B}}{b}, \Pi\right).$$

Moreover, let \mathbf{A} be a linear potential associated with \mathbf{B} . Then $\mathbf{x} \mapsto \Psi(\sqrt{b}\mathbf{x})$ is an eigenfunction of $H(\mathbf{A}, \Pi)$ associated with $E(\mathbf{B}, \Pi)$ if and only if $\mathbf{x} \mapsto \Psi(\mathbf{x})$ is an eigenfunction of $H(\mathbf{A}/b, \Pi)$ associated with $E(\mathbf{B}/b, \Pi)$.

That is why we can reduce to consider *model problems* on cones $\Pi \in \overline{\mathfrak{P}}_3$ with unitary constant magnetic fields.

4.2. Singular chains and generalized eigenvectors for model problems. Let $\Pi \in \overline{\mathfrak{P}}_3$ be a polyhedral cone and \mathbf{B} be a unitary constant magnetic field associated with an affine potential \mathbf{A} . Let $\Gamma \in \overline{\mathfrak{P}}_d$ be a minimal reduced cone associated with Π . We recall that this means that $\Pi \equiv \mathbb{R}^{3-d} \times \Gamma$ and that the dimension d is minimal for such an equivalence.

Let $\mathfrak{C}_0(\Pi)$ denote the singular chains of Π originating at its vertex $\mathbf{0}$ and let $\mathfrak{C}_0^*(\Pi)$ be the subset of chains of length ≥ 2 (see (3.2)). Note that $\mathfrak{C}_0^*(\Pi)$ is empty if and only if $\Pi = \mathbb{R}^3$, i.e., if $d = 0$. We introduce the energy along higher chains:

Definition 4.2 (Energy along higher chains). We define the quantity

$$(4.3) \quad \mathcal{E}^*(\mathbf{B}, \Pi) := \begin{cases} \inf_{\mathbb{X} \in \mathfrak{C}_0^*(\Pi)} E(\mathbf{B}, \Pi_{\mathbb{X}}) & \text{if } d > 0, \\ +\infty & \text{if } d = 0, \end{cases}$$

which is the infimum of the ground energy of the magnetic Laplacian over all the singular chains of length ≥ 2 .

If $d > 0$, let $\Omega_0 \in \overline{\mathfrak{D}}(\mathbb{S}^{d-1})$ be the section of Γ . Since Π is a polyhedral cone, we have (cf. Remark 3.3)

$$(4.4) \quad \mathcal{E}^*(\mathbf{B}, \Pi) = \inf_{\mathbf{x}_1 \in \overline{\Omega}_0} E(\mathbf{B}, \Pi_{(\mathbf{0}, \mathbf{x}_1)}),$$

i.e., among all chains $\mathbb{X} \in \mathfrak{C}_0^*(\Pi)$, we can restrict to those of length 2, $\mathbb{X} = (\mathbf{0}, \mathbf{x}_1)$.

Since the cone Π is unbounded, it is relevant to define $\lambda_{\text{ess}}(\mathbf{B}, \Pi)$ as the bottom of the essential spectrum of the operator $H(\mathbf{A}, \Pi)$. When $d \leq 2$, due to translation invariance we have $E(\mathbf{B}, \Pi) = \lambda_{\text{ess}}(\mathbf{B}, \Pi)$. When $d = 3$, the operator $H(\mathbf{A}, \Pi)$ may have discrete spectrum.

With the aim of constructing quasimodes for our original problem on Ω , we need generalized eigenvectors for its tangent problems. For this we make use of the localized domain $\text{Dom}_{\text{loc}}(H(\mathbf{A}, \Pi))$ of the model magnetic Laplacian $H(\mathbf{A}, \Pi)$ as introduced in (1.17).

Definition 4.3 (Generalized eigenvector). Let $\Pi \in \overline{\mathfrak{P}}_3$ be a polyhedral cone and \mathbf{A} an affine magnetic potential. We call *generalized eigenvector* for $H(\mathbf{A}, \Pi)$ a nonzero function $\Psi \in \text{Dom}_{\text{loc}}(H(\mathbf{A}, \Pi))$ associated with a real number Λ , so that

$$(4.5) \quad \begin{cases} (-i\nabla + \mathbf{A})^2 \Psi = \Lambda \Psi & \text{in } \Pi, \\ (-i\partial_n + \mathbf{n} \cdot \mathbf{A}) \Psi = 0 & \text{on } \partial\Pi. \end{cases}$$

Definition 4.4 (Admissible generalized eigenvector). Under the hypothesis of Definition 4.3, a generalized eigenvector Ψ for $H(\mathbf{A}, \Pi)$ is said to be *admissible* if there exist a rotation $\underline{U} : \Pi \mapsto \mathbb{R}^{3-k} \times \Upsilon$ with $k \geq d$ and $\Upsilon \in \overline{\mathfrak{P}}_k$, and a system of coordinates $(\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \mathbb{R}^k$ such that

$$(4.6) \quad \Psi \circ \underline{U}^{-1}(\mathbf{y}, \mathbf{z}) = e^{i\varphi(\mathbf{y}, \mathbf{z})} \Phi(\mathbf{z}),$$

with some real polynomial function φ of degree ≤ 2 and some exponentially decreasing function Φ , namely there exist positive constants c_Ψ and C_Ψ such that

$$(4.7) \quad \|e^{c_\Psi|\mathbf{z}|} \Phi\|_{L^2(\Upsilon)} \leq C_\Psi \|\Phi\|_{L^2(\Upsilon)}.$$

We will denote by $\mathbf{x}^{\natural} = (\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \Upsilon$ the coordinates associated with Ψ .

Remark 4.5. By Lemma A.4 it is straightforward that in coordinates \mathbf{x}^{\natural} the magnetic potential and the magnetic field are transformed into

$$\mathbf{A}^{\natural} = J^{\top}(\mathbf{A} \circ \underline{U}^{-1}) \quad \text{and} \quad \mathbf{B}^{\natural} = J^{\top}(\mathbf{B} \circ \underline{U}^{-1})$$

where the orthogonal 3×3 matrix J is such that $\underline{U}^{-1}(\mathbf{x}^{\natural}) = J\mathbf{x}^{\natural}$. Note also that $\underline{U}(\mathbf{x}) = J^{\top}\mathbf{x}$. Therefore Ψ is an admissible generalized eigenfunction of $H(\mathbf{A}, \Pi)$ associated with the value Λ if and only if $\Psi \circ \underline{U}^{-1}$ is a generalized eigenfunction of $H(\mathbf{A}^{\natural}, \mathbb{R}^{3-k} \times \Upsilon)$ associated with the same value Λ .

Lemma 4.6. *If Ψ is an admissible generalized eigenvector for $H(\mathbf{A}, \Pi)$ associated with Λ , for any other affine magnetic potential \mathbf{A}' such that $\text{curl } \mathbf{A}' = \text{curl } \mathbf{A}$, the operator $H(\mathbf{A}', \Pi)$ possesses an admissible generalized eigenvector Ψ' associated with the same value Λ .*

Proof. If $\text{curl } \mathbf{A} = \text{curl } \mathbf{A}'$ and \mathbf{A}, \mathbf{A}' are both affine, there exists a polynomial function ϕ of degree 2 such that $\mathbf{A} = \mathbf{A}' - \nabla\phi$. Using a change of gauge (Lemma A.1), we find that Ψ' defined as

$$\Psi'(\mathbf{x}) = e^{i\phi(\mathbf{x})} e^{i\varphi(\mathbf{y}, \mathbf{z})} \Phi(\mathbf{z}), \quad \mathbf{x} \in \Pi,$$

is an admissible generalized eigenvector for $H(\mathbf{A}', \Pi)$. \square

4.3. Dichotomy for model problems. The main result which we prove in this section is a dichotomy statement, as follows.

Theorem 4.7. *Let $\Pi \in \overline{\mathfrak{P}}_3$ be a polyhedral cone and \mathbf{B} be a constant nonzero magnetic field. Let \mathbf{A} be any associated affine magnetic potential. Recall that $E(\mathbf{B}, \Pi)$ is the ground energy of $H(\mathbf{A}, \Pi)$ and $\mathcal{E}^*(\mathbf{B}, \Pi)$ is the energy along higher chains introduced in (4.4). Then,*

$$(4.8) \quad E(\mathbf{B}, \Pi) \leq \mathcal{E}^*(\mathbf{B}, \Pi)$$

and we have the dichotomy:

- (i) *If $E(\mathbf{B}, \Pi) < \mathcal{E}^*(\mathbf{B}, \Pi)$, then $H(\mathbf{A}, \Pi)$ admits an admissible generalized eigenvector associated with the value $E(\mathbf{B}, \Pi)$.*
- (ii) *If $E(\mathbf{B}, \Pi) = \mathcal{E}^*(\mathbf{B}, \Pi)$, then there exists a singular chain $\mathbb{X} \in \mathfrak{C}_0^*(\Pi)$ such that*

$$E(\mathbf{B}, \Pi_{\mathbb{X}}) = E(\mathbf{B}, \Pi) \quad \text{and} \quad E(\mathbf{B}, \Pi_{\mathbb{X}}) < \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbb{X}}).$$

Remark 4.8. In the case (ii), we note that by statement (i) applied to the cone $\Pi_{\mathbb{X}}$, $H(\mathbf{A}, \Pi_{\mathbb{X}})$ admits an admissible generalized eigenvector associated with the value $E(\mathbf{B}, \Pi)$.

- *Outline of the proof of Theorem 4.7.* Owing to Lemma 4.1 we may assume that $\mathbf{B} := \text{curl } \mathbf{A}$ is unitary. The proof relies on an exhaustion of cases, organized according increasing values of d , the dimension of the reduced cone of Π . First note that quantities $E(\mathbf{B}, \Pi)$ and $\mathcal{E}^*(\mathbf{B}, \Pi)$, like the definition of admissible generalized eigenvectors, are independent of a choice of Cartesian coordinates. Thus, for each value of d , ranging from 0 to 3, once Π and a constant unitary magnetic field \mathbf{B} are chosen, we exhibit a system of Cartesian coordinates $\mathbf{x} = (x_1, x_2, x_3)$ that allows the simplest possible description of the configuration (\mathbf{B}, Π) . In these coordinates, the magnetic field can be viewed as a reference one and we denote it by $\underline{\mathbf{B}}$. We also choose a corresponding reference linear potential $\underline{\mathbf{A}}$, since we have gauge independence by virtue of Lemma 4.6. Then relying on various results corresponding

to each case, we prove that we are either in situation (i) and exhibit coordinates³ (\mathbf{y}, \mathbf{z}) and admissible generalized eigenvectors, or in situation (ii).

In each of Sections 4.4–4.7 we consider one value of d , from 0 to 3, which will achieve the proof of Theorem 4.7. In Section 4.8, we collect all possible structures of admissible generalized eigenvectors Ψ , organized according to increasing values of k , the number of directions in which Ψ has exponential decay.

4.4. Full space, $d = 0$. Π is the full space. We take coordinates $\mathbf{x} = (x_1, x_2, x_3)$ so that

$$\Pi = \mathbb{R}^3 \quad \text{and} \quad \underline{\mathbf{B}} = (1, 0, 0),$$

and choose as reference potential

$$\underline{\mathbf{A}} = (0, -\frac{x_3}{2}, \frac{x_2}{2}).$$

Hence

$$H(\underline{\mathbf{A}}, \Pi) = H(\underline{\mathbf{A}}, \mathbb{R}^3) = D_1^2 + (D_2 - \frac{x_3}{2})^2 + (D_3 + \frac{x_2}{2})^2 \quad \text{with} \quad D_j = -i\partial_{x_j}.$$

It is classical (see [20]) that the spectrum of $H(\underline{\mathbf{A}}, \mathbb{R}^3)$ is $[1, +\infty)$. Therefore

$$(4.9) \quad E(\underline{\mathbf{B}}, \mathbb{R}^3) = 1.$$

An admissible generalized eigenfunction associated to the ground energy is

$$\Psi(\mathbf{x}) = e^{-(x_2^2 + x_3^2)/4},$$

which has the form (4.6) with $\mathbf{y} = x_1$, $\mathbf{z} = (x_2, x_3)$, and $\varphi \equiv 0$.

4.5. Half space, $d = 1$. Π is a half-space. We take coordinates $\mathbf{x} = (x_1, x_2, x_3)$ so that

$$\Pi = \mathbb{R}_+^3 := \{(x_1, x_2, x_3) \in \mathbb{R}^3, x_3 > 0\} \quad \text{and} \quad \underline{\mathbf{B}} = (0, b_1, b_2) \quad \text{with} \quad b_1^2 + b_2^2 = 1,$$

and choose as reference potential

$$\underline{\mathbf{A}} = (b_1 x_3 - b_2 x_2, 0, 0).$$

Hence

$$H(\underline{\mathbf{A}}, \Pi) = H(\underline{\mathbf{A}}, \mathbb{R}_+^3) = (D_1 + b_1 x_3 - b_2 x_2)^2 + D_2^2 + D_3^2.$$

We note that

$$(4.10) \quad \mathcal{E}^*(\underline{\mathbf{B}}, \mathbb{R}_+^3) = E(\underline{\mathbf{B}}, \mathbb{R}^3) = 1.$$

There exists $\theta \in [0, 2\pi)$ such that $b_1 = \cos \theta$ and $b_2 = \sin \theta$, so that θ is the angle between the magnetic field and the boundary of \mathbb{R}_+^3 . Due to symmetries we can reduce to $\theta \in [0, \frac{\pi}{2}]$. Denote by \mathcal{F}_1 the Fourier transform in x_1 -variable, τ the Fourier variable associated with x_1 , and

$$\widehat{H}_\tau(\underline{\mathbf{A}}, \mathbb{R}_+^3) := (\tau + b_1 x_3 - b_2 x_2)^2 + D_2^2 + D_3^2,$$

³As our system of Cartesian coordinates $\mathbf{x} = (x_1, x_2, x_3)$ will be already chosen in some optimal way, coordinates (\mathbf{y}, \mathbf{z}) will simply be a splitting of (x_1, x_2, x_3) . But it is noticeable that the dimension k of \mathbf{z} variables may be strictly larger than d .

acting on $L^2(\mathbb{R} \times \mathbb{R}_+)$ with natural boundary condition. There holds

$$\mathcal{F}_1 H(\mathbf{A}, \mathbb{R}_+^3) \mathcal{F}_1^* = \int_{\tau \in \mathbb{R}}^{\oplus} \widehat{H}_\tau(\mathbf{A}, \mathbb{R}_+^3) d\tau.$$

We discriminate three cases:

- *Tangent field.* $\theta = 0$, then $\widehat{H}_\tau(\mathbf{A}, \mathbb{R}_+^3) := D_2^2 + D_3^2 + (\tau + x_3)^2$, let ξ be the partial Fourier variable associated with x_2 and define the new operators

$$\widehat{H}_{\xi, \tau}(\mathbf{A}, \mathbb{R}_+^3) := \xi^2 + D_3^2 + (\tau + x_3)^2, \quad \mathcal{H}(\tau) = D_3^2 + (\tau + x_3)^2,$$

where $\mathcal{H}(\tau)$ (sometimes called the de Gennes operator) acts on $L^2(\mathbb{R}_+)$ with Neumann boundary condition. There holds

$$\inf \mathfrak{S}(\mathcal{H}(\tau)) = \mu(\tau), \quad \inf \mathfrak{S}(\widehat{H}_{\tau, \xi}(\mathbf{A}, \mathbb{R}_+^3)) = \mu(\tau) + \xi^2,$$

in which the behavior of the first eigenvalue $\mu(\tau)$ is well-known (see [11]): The function μ admits a unique minimum denoted by $\Theta_0 \simeq 0.59$ for the value $\tau = -\sqrt{\Theta_0}$. Hence

$$E(\mathbf{B}, \mathbb{R}_+^3) = \Theta_0 < \mathcal{E}^*(\mathbf{B}, \mathbb{R}_+^3).$$

We are in case (i) of Theorem 4.7. If Φ denotes an eigenvector of $\mathcal{H}(\tau)$ associated with Θ_0 (function of $x_3 \in \mathbb{R}_+$), a corresponding admissible generalized eigenvector is

$$(4.11) \quad \Psi(\mathbf{x}) = e^{-i\sqrt{\Theta_0}x_1} \Phi(x_3).$$

which has the form (4.6) with $\mathbf{y} = (x_1, x_2)$, $\mathbf{z} = x_3$, and $\varphi(\mathbf{y}, \mathbf{z}) = -x_1\sqrt{\Theta_0}$.

- *Normal field.* $\theta = \frac{\pi}{2}$, then $\widehat{H}_\tau(\mathbf{A}, \mathbb{R}_+^3) := D_2^2 + D_3^2 + (\tau - x_2)^2$. There holds for all $\tau \in \mathbb{R}$, $\inf \mathfrak{S}(\widehat{H}_\tau(\mathbf{A}, \mathbb{R}_+^3)) = 1$ (see [22, Theorem 3.1]), hence

$$E(\mathbf{B}, \mathbb{R}_+^3) = 1 = \mathcal{E}^*(\mathbf{B}, \mathbb{R}_+^3).$$

We are in case (ii) of Theorem 4.7. The chain \mathbb{X} is given by $(\mathbf{0}, 1)$ and $\Pi_{\mathbb{X}} = \mathbb{R}^3$, for which $E(\mathbf{B}, \mathbb{R}^3) < \mathcal{E}^*(\mathbf{B}, \mathbb{R}^3)$.

- *Neither tangent nor normal.* $\theta \in (0, \frac{\pi}{2})$. Then for any $\tau \in \mathbb{R}$, $\widehat{H}_\tau(\mathbf{A}, \mathbb{R}_+^3)$ is isospectral to $\widehat{H}_0(\mathbf{A}, \mathbb{R}_+^3)$ the ground energy of which is an eigenvalue $\sigma(\theta) < 1$ (see [17]). We deduce

$$E(\mathbf{B}, \mathbb{R}_+^3) = \sigma(\theta) \quad \text{with} \quad \sigma(\theta) < 1.$$

We are in case (i) of Theorem 4.7. We recall:

Lemma 4.9. *The function $\theta \mapsto \sigma(\theta)$ is continuous and increasing on $(0, \frac{\pi}{2})$ ([17, 22]). Set $\sigma(0) = \Theta_0$ and $\sigma(\frac{\pi}{2}) = 1$. Then the function $\theta \mapsto \sigma(\theta)$ is of class \mathcal{C}^1 on $[0, \frac{\pi}{2}]$ ([7]).*

The first eigenvalue of $\widehat{H}_0(\mathbf{A}, \mathbb{R}_+^3)$ is associated with an exponentially decreasing eigenvector Φ which is a function of $(x_2, x_3) \in \mathbb{R} \times \mathbb{R}_+$. An admissible generalized eigenvector for $H(\mathbf{A}, \mathbb{R}_+^3)$ is given by

$$(4.12) \quad \Psi(\mathbf{x}) = \Phi(x_2, x_3),$$

which has the form (4.6) with $\mathbf{y} = x_1$, $\mathbf{z} = (x_2, x_3)$, and $\varphi \equiv 0$.

Thus Theorem 4.7 is proved for half-spaces.

4.6. **Wedges**, $d = 2$. Π is a wedge and let $\alpha \in (0, \pi) \cup (\pi, 2\pi)$ denote its opening. Let us introduce the model sector \mathcal{S}_α and the model wedge \mathcal{W}_α

$$(4.13) \quad \mathcal{S}_\alpha = \begin{cases} \{x = (x_2, x_3), x_2 \tan \frac{\alpha}{2} > |x_3|\} & \text{if } \alpha \in (0, \pi) \\ \{x = (x_2, x_3), x_2 \tan \frac{\alpha}{2} > -|x_3|\} & \text{if } \alpha \in (\pi, 2\pi) \end{cases} \quad \text{and} \quad \mathcal{W}_\alpha = \mathbb{R} \times \mathcal{S}_\alpha.$$

We take coordinates $\mathbf{x} = (x_1, x_2, x_3)$ so that

$$\Pi = \mathcal{W}_\alpha \quad \text{and} \quad \underline{\mathbf{B}} = (b_1, b_2, b_3) \quad \text{with} \quad b_1^2 + b_2^2 + b_3^2 = 1,$$

and choose as reference potential

$$\underline{\mathbf{A}} = (b_2 x_3 - b_3 x_2, 0, b_1 x_2).$$

Hence

$$H(\underline{\mathbf{A}}, \Pi) = H(\underline{\mathbf{A}}, \mathcal{W}_\alpha) = (D_1 + b_2 x_3 - b_3 x_2)^2 + D_2^2 + (D_3 + b_1 x_2)^2.$$

Denote by τ the Fourier variable associated with x_1 , and

$$(4.14) \quad \hat{H}_\tau(\underline{\mathbf{A}}, \mathcal{W}_\alpha) := (\tau + b_2 x_3 - b_3 x_2)^2 + D_2^2 + (D_3 + b_1 x_2)^2$$

acting on $L^2(\mathcal{S}_\alpha)$ with natural Neumann boundary condition. We introduce the notation:

$$s(\underline{\mathbf{B}}, \mathcal{S}_\alpha; \tau) := \inf \mathfrak{S}(\hat{H}_\tau(\underline{\mathbf{A}}, \mathcal{W}_\alpha)),$$

so that we have the direct Fourier integral decomposition

$$\mathcal{F}_1 H(\underline{\mathbf{A}}, \mathcal{W}_\alpha) \mathcal{F}_1^* = \int_{\tau \in \mathbb{R}}^{\oplus} \hat{H}_\tau(\underline{\mathbf{A}}, \mathcal{W}_\alpha) d\tau$$

and the relation

$$(4.15) \quad E(\underline{\mathbf{B}}, \mathcal{W}_\alpha) = \inf_{\tau \in \mathbb{R}} s(\underline{\mathbf{B}}, \mathcal{S}_\alpha; \tau).$$

The singular chains of $\mathfrak{C}_0^*(\mathcal{W}_\alpha)$ have three equivalence classes, cf. Definition 3.2 and Section 3.3 (3), corresponding to three distinct model operators, associated to half-spaces Π_α^\pm corresponding to the faces $\partial^\pm \mathcal{W}_\alpha$ of \mathcal{W}_α , and to the full space \mathbb{R}^3 . Thus

$$\mathcal{E}^*(\underline{\mathbf{B}}, \mathcal{W}_\alpha) = \min\{E(\underline{\mathbf{B}}, \Pi_\alpha^+), E(\underline{\mathbf{B}}, \Pi_\alpha^-), E(\underline{\mathbf{B}}, \mathbb{R}^3)\}.$$

Let $\theta^\pm \in [0, \frac{\pi}{2}]$ be the angle between $\underline{\mathbf{B}}$ and the face $\partial \Pi_\alpha^\pm$. We have, cf. Lemma 4.9,

$$(4.16) \quad \mathcal{E}^*(\underline{\mathbf{B}}, \mathcal{W}_\alpha) = \min\{\sigma(\theta^+), \sigma(\theta^-), 1\} = \sigma(\min\{\theta^+, \theta^-\}).$$

When $\Pi = \mathcal{W}_\alpha$, Theorem 4.7 relies on the following result [30, Theorem 3.5]:

Lemma 4.10. *We have $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha) \leq \mathcal{E}^*(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$.*

Moreover, if $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha) < \mathcal{E}^(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$, then the function $\tau \mapsto s(\underline{\mathbf{B}}, \mathcal{S}_\alpha; \tau)$ reaches its infimum. Let τ^* be a minimizer. Then $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$ is a discrete eigenvalue for the operator $\hat{H}_{\tau^*}(\underline{\mathbf{A}}, \mathcal{W}_\alpha)$ and the associated eigenfunctions have exponential decay.*

From the previous lemma we deduce

- (i) If $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha) < \mathcal{E}^*(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$, there exists τ^* such that the operator $\widehat{H}_{\tau^*}(\underline{\mathbf{A}}, \mathcal{W}_\alpha)$ admits an exponential decaying eigenfunction Φ of $(x_2, x_3) \in \mathcal{S}_\alpha$ associated with $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$. The function

$$\Psi(\mathbf{x}) = e^{i\tau^*x_1}\Phi(x_2, x_3)$$

is an admissible generalized eigenvector for the operator $H(\underline{\mathbf{A}}, \mathcal{W}_\alpha)$ associated with $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$. It has the form (4.6) with $\mathbf{y} = x_1$, $\mathbf{z} = (x_2, x_3)$, and $\varphi(\mathbf{y}, \mathbf{z}) = \tau^*\mathbf{y}$.

- (ii) If $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha) = \mathcal{E}^*(\underline{\mathbf{B}}, \mathcal{W}_\alpha)$, let $\circ \in \{-, +\}$ satisfy $\theta^\circ = \min(\theta^-, \theta^+)$ and Π_α° be the corresponding face. We have $\theta^\circ \in [0, \frac{\pi}{2})$ and $\mathcal{E}^*(\underline{\mathbf{B}}, \mathcal{W}_\alpha) = \sigma(\theta^\circ)$. We deduce from Section 4.5 that $\mathcal{E}^*(\underline{\mathbf{B}}, \Pi_\alpha^\circ) = 1$, hence $E(\underline{\mathbf{B}}, \mathcal{W}_\alpha) = E(\underline{\mathbf{B}}, \Pi_\alpha^\circ) < \mathcal{E}^*(\underline{\mathbf{B}}, \Pi_\alpha^\circ)$.

Thus Theorem 4.7 is proved for wedges. We extend the definition of \mathcal{W}_α to $\alpha = \pi$ by setting $\mathcal{W}_\pi := \mathbb{R}_+^3$. Let us quote now the continuity result of [30, Theorem 4.5]:

Lemma 4.11. *The function $(\mathbf{B}, \alpha) \mapsto E(\mathbf{B}, \mathcal{W}_\alpha)$ is continuous on $\mathbb{S}^2 \times (0, 2\pi)$.*

We end this paragraph by a few examples.

Example 4.12. Let $\mathbf{B} \in \mathbb{S}^2$ be a constant magnetic field. Let α be chosen in $(0, \pi) \cup (\pi, 2\pi)$.

a) For α small enough $E(\mathbf{B}, \mathcal{W}_\alpha) < \mathcal{E}^*(\mathbf{B}, \mathcal{W}_\alpha)$ (see [30] when the magnetic field is not tangent to the plane of symmetry of the wedge and [28, Ch. 7] otherwise).

b) Let $\mathbf{B} = (0, 0, 1)$ be tangent to the edge. Then $\mathcal{E}^*(\mathbf{B}, \mathcal{W}_\alpha) = \Theta_0$ and $E(\mathbf{B}, \mathcal{W}_\alpha) = E(1, \mathcal{S}_\alpha)$, cf. Section 2.1.2. According to whether the ground energy $E(1, \mathcal{S}_\alpha)$ of the plane sector \mathcal{S}_α is less than Θ_0 or equal to Θ_0 , we are in case (i) or (ii) of the dichotomy.

c) Let \mathbf{B} be tangent to a face of the wedge and normal to the edge. Then $\mathcal{E}^*(\mathbf{B}, \mathcal{W}_\alpha) = \Theta_0$. It is proved in [29] that there holds $E(\mathbf{B}, \mathcal{W}_\alpha) < \Theta_0$ for α small enough, whereas $E(\mathbf{B}, \mathcal{W}_\alpha) = \Theta_0$ for $\alpha \in [\frac{\pi}{2}, \pi)$.

As a direct consequence of the whole description performed in the previous Sections 4.4-4.6, we obtain the following continuity statements.

Theorem 4.13. *Let Π be a cone in $\overline{\mathfrak{P}}_3$ with $d < 3$ (i.e. Π is the full space, a half-space or a wedge). Then the function $\mathbf{B} \mapsto E(\mathbf{B}, \Pi)$ is continuous on \mathbb{S}^2 .*

Corollary 4.14. *Let Π be a cone in $\overline{\mathfrak{P}}_3$. The function $\mathbf{B} \mapsto \mathcal{E}^*(\mathbf{B}, \Pi)$ is continuous on \mathbb{S}^2 .*

4.7. Polyhedral cones, $d = 3$. The main result of this paragraph is the characterization of the bottom $\lambda_{\text{ess}}(\mathbf{B}, \Pi)$ of the essential spectrum of $H(\mathbf{A}, \Pi)$.

Proposition 4.15. *Let $\Pi \in \overline{\mathfrak{P}}_3$ be a polyhedral cone with $d = 3$, which means that Π is not a wedge, nor a half-space, nor the full space. Let \mathbf{B} be a constant magnetic field. With the quantity $\mathcal{E}^*(\mathbf{B}, \Pi)$ introduced in (4.3), there holds*

$$\lambda_{\text{ess}}(\mathbf{B}, \Pi) = \mathcal{E}^*(\mathbf{B}, \Pi).$$

Before writing proof details, let us specify what is $\mathcal{E}^*(\mathbf{B}, \Pi)$ in the case of a polyhedral cone. Let Ω_0 be the section of Π , i.e., $\Omega_0 = \Pi \cap \mathbb{S}^2$. We recall from (4.4) that

$$(4.17) \quad \mathcal{E}^*(\mathbf{B}, \Pi) = \inf_{\mathbf{x}_1 \in \overline{\Omega}_0} E(\mathbf{B}, \Pi_{(\mathbf{0}, \mathbf{x}_1)}).$$

In fact the set of equivalence classes (Definition 3.2) of the chains $\mathbb{X} = (\mathbf{0}, \mathbf{x}_1)$ is finite. Let us describe this set, cf. Section 3.3 (4). Let \mathfrak{F} and \mathfrak{E} be the set of faces \mathbf{f} and edges \mathbf{e} of Π . For $\mathbf{f} \in \mathfrak{F}$, let $\Pi_{\mathbf{f}}$ be the half-space whose boundary contains \mathbf{f} and containing points of Π near any point of \mathbf{f} . For $\mathbf{e} \in \mathfrak{E}$, there are two faces \mathbf{f}_e^\pm adjacent to \mathbf{e} . Let $\Pi_{\mathbf{e}}$ be the wedge whose boundary contains $\mathbf{e} \cup \mathbf{f}_e^+ \cup \mathbf{f}_e^-$ and containing points of Π near any point of $\mathbf{f}_e^+ \cup \mathbf{f}_e^-$.

Let $\mathbf{x}_1 \in \overline{\Omega}_0$. There are three possibilities:

- (1) \mathbf{x}_1 is interior to Ω_0 . Then $\Pi_{(\mathbf{0}, \mathbf{x}_1)} = \mathbb{R}^3$.
- (2) \mathbf{x}_1 belongs to a side of Ω_0 . This side is contained in a face \mathbf{f} of Π . Then $\Pi_{(\mathbf{0}, \mathbf{x}_1)} = \Pi_{\mathbf{f}}$.
- (3) \mathbf{x}_1 belongs to a vertex of Ω_0 . This vertex is contained in an edge \mathbf{e} of Π . Then $\Pi_{(\mathbf{0}, \mathbf{x}_1)} = \Pi_{\mathbf{e}}$.

We have that

$$(4.18) \quad \mathcal{E}^*(\mathbf{B}, \Pi) = \min \left\{ \min_{\mathbf{e} \in \mathfrak{E}} E(\mathbf{B}, \Pi_{\mathbf{e}}), \min_{\mathbf{f} \in \mathfrak{F}} E(\mathbf{B}, \Pi_{\mathbf{f}}), 1 \right\}.$$

Since (4.8) is proved for $d = 2$, we have $E(\mathbf{B}, \Pi_{\mathbf{e}}) \leq \min\{E(\mathbf{B}, \Pi_{\mathbf{f}_e^+}), E(\mathbf{B}, \Pi_{\mathbf{f}_e^-})\}$. Therefore equation (4.18) becomes

$$(4.19) \quad \mathcal{E}^*(\mathbf{B}, \Pi) = \min_{\mathbf{e} \in \mathfrak{E}} \{E(\mathbf{B}, \Pi_{\mathbf{e}})\}.$$

We recall the Persson Lemma that gives a characterization of the bottom of the essential spectrum (see [27]):

Lemma 4.16. *We have*

$$\lambda_{\text{ess}}(\mathbf{B}, \Pi) = \lim_{R \rightarrow +\infty} \Sigma(\mathbf{B}, \Pi, R)$$

with

$$\Sigma(\mathbf{B}, \Pi, R) := \inf_{\substack{\psi \in \mathcal{C}_0^\infty(\overline{\Pi} \setminus \mathcal{B}_R) \\ \psi \neq 0}} \frac{q[\mathbf{A}, \Pi](\psi)}{\|\psi\|^2}$$

where \mathcal{B}_R is the ball of radius R centered at the origin and $\mathcal{C}\mathcal{B}_R$ its complementary in \mathbb{R}^3 .

Proof. (of Proposition 4.15). Let \mathbf{A} be a linear potential associated with \mathbf{B} .

Upper bound: We denote by \mathbf{x}_{e^*} a vertex of $\overline{\Omega}_0$ and \mathbf{e}^* the associated edge such that $\mathcal{E}^*(\mathbf{B}, \Pi) = E(\mathbf{B}, \Pi_{\mathbf{e}^*})$, cf. (4.19). Let $\varepsilon > 0$, there exists $\psi_\varepsilon \in \mathcal{C}_0^\infty(\overline{\Pi}_{\mathbf{e}^*})$ a normalized function such that

$$q[\mathbf{A}, \Pi_{\mathbf{e}^*}](\psi_\varepsilon) \leq E(\mathbf{B}, \Pi_{\mathbf{e}^*}) + \varepsilon.$$

For $r > 0$ we define

$$\psi_\varepsilon^r(\mathbf{x}) := e^{i\langle \mathbf{x}, \mathbf{A}(r\mathbf{x}_{e^*}) \rangle} \psi_\varepsilon(\mathbf{x} - r\mathbf{x}_{e^*}),$$

so that we have, due to gauge invariance and translation effect, cf. Lemma A.2,

$$\text{supp}(\psi_\varepsilon^r) = \text{supp}(\psi_\varepsilon) + r\mathbf{x}_{\mathbf{e}^*} \quad \text{and} \quad q[\mathbf{A}, \Pi](\psi_\varepsilon^r) = q[\mathbf{A}, \Pi_{\mathbf{e}^*}](\psi_\varepsilon).$$

Let $R > 0$, for r large enough we have $\text{supp}(\psi_\varepsilon^r) \subset \mathbb{C}\mathcal{B}_R$ and $\psi_\varepsilon^r \in \text{Dom}(q[\mathbf{A}, \Pi])$. We get

$$q[\mathbf{A}, \Pi](\psi_\varepsilon^r) = q[\mathbf{A}, \Pi_{\mathbf{e}^*}](\psi_\varepsilon) \leq E(\mathbf{B}, \Pi_{\mathbf{e}^*}) + \varepsilon.$$

We deduce

$$\forall \varepsilon > 0, \forall R > 0, \quad \Sigma(\mathbf{B}, \Pi, R) \leq E(\mathbf{B}, \Pi_{\mathbf{e}^*}) + \varepsilon$$

and Lemma 4.16 provides the upper bound of Proposition 4.15: $\lambda_{\text{ess}}(\mathbf{B}, \Pi) \leq \mathcal{E}^*(\mathbf{B}, \Pi)$.

Lower bound: Let

$$\mathcal{U}_0 \cup \left(\bigcup_{\mathbf{f} \in \mathfrak{F}} \mathcal{U}_{\mathbf{f}} \right) \cup \left(\bigcup_{\mathbf{e} \in \mathfrak{E}} \mathcal{U}_{\mathbf{e}} \right)$$

a covering of $\overline{\Omega}_0$ according to its stratification, which means that

$$\mathcal{U}_0 \subset \Omega_0, \quad \mathcal{U}_{\mathbf{f}} \cap \overline{\Omega}_0 = \Pi_{\mathbf{f}} \cap \overline{\Omega}_0 \quad (\forall \mathbf{f} \in \mathfrak{F}), \quad \text{and} \quad \mathcal{U}_{\mathbf{e}} \cap \overline{\Omega}_0 = \Pi_{\mathbf{e}} \cap \overline{\Omega}_0 \quad (\forall \mathbf{e} \in \mathfrak{E}).$$

Let $\hat{\chi}_j, j \in \mathfrak{J} := \{0\} \cup \mathfrak{F} \cup \mathfrak{E}$, be an associated partition of unity of the section $\overline{\Omega}_0$ such that

$$\sum_{j \in \mathfrak{J}} (\hat{\chi}_j)^2 = 1 \quad \text{and} \quad \text{supp}(\hat{\chi}_j) \subset \mathcal{U}_j, \quad \forall j \in \mathfrak{J}.$$

Let $\chi \in \mathcal{C}^\infty(\mathbb{R}_+)$ such that $\chi \equiv 0$ on $[0, \frac{1}{2}]$ and $\chi \equiv 1$ on $[1, +\infty)$. We now define a partition of the unity of $\mathbb{C}\mathcal{B}_R \cap \overline{\Pi}$ by setting

$$\chi_j^R(\mathbf{x}) = \chi\left(\frac{|\mathbf{x}|}{R}\right) \hat{\chi}_j\left(\frac{\mathbf{x}}{|\mathbf{x}|}\right), \quad j \in \mathfrak{J}.$$

We have $\sum_j (\chi_j^R)^2 = 1$ on $\mathbb{C}\mathcal{B}_R \cap \overline{\Pi}$ and

$$\forall R > 0, \quad \sum_{j \in \mathfrak{J}} |\nabla \chi_j^R|^2 \leq CR^{-2}.$$

Moreover we have $\text{supp}(\chi_j^R) \cap \overline{\Pi} \subset \overline{\Pi}_j, j \in \mathfrak{J} = \{0\} \cup \mathfrak{F} \cup \mathfrak{E}$, where we have set $\Pi_0 = \mathbb{R}^3$. The IMS formula for quadratic forms (see Lemma A.7) provides

$$\begin{aligned} \forall \psi \in \mathcal{C}_0^\infty(\overline{\Pi} \cap \mathbb{C}\mathcal{B}_R), \quad q[\mathbf{A}, \Pi](\psi) &\geq \sum_{j \in \mathfrak{J}} q[\mathbf{A}, \Pi](\chi_j^R \psi) - CR^{-2} \|\psi\|^2 \\ &= \sum_{j \in \mathfrak{J}} q[\mathbf{A}, \Pi_j](\chi_j^R \psi) - CR^{-2} \|\psi\|^2 \\ &\geq \sum_{j \in \mathfrak{J}} E(\mathbf{B}, \Pi_j) \|\chi_j^R \psi\|^2 - CR^{-2} \|\psi\|^2 \\ &\geq (\mathcal{E}^*(\mathbf{B}, \Pi) - CR^{-2}) \|\psi\|^2. \end{aligned}$$

Thus we deduce the lower bound of Proposition 4.15 by using Lemma 4.16. \square

Then it is clear that Theorem 4.7 in the case of polyhedral cones is a consequence of Proposition 4.15:

- if $E(\mathbf{B}, \Pi) < \mathcal{E}^*(\mathbf{B}, \Pi) = \lambda_{\text{ess}}(\mathbf{B}, \Pi)$, then there exists an eigenvector for $H(\mathbf{A}, \Pi)$, which by standard arguments based on Agmon estimates is exponentially decreasing, see [1].
- if $E(\mathbf{B}, \Pi) = \mathcal{E}^*(\mathbf{B}, \Pi) = \lambda_{\text{ess}}(\mathbf{B}, \Pi) = \min_{\mathbf{e} \in \mathcal{E}} \{E(\mathbf{B}, \Pi_{\mathbf{e}})\}$, we are reduced to the previous cases in lower values of d .

Example 4.17 (Octant). Let $\Pi = (\mathbb{R}_+)^3$ be the model octant and \mathbf{B} be a constant magnetic field with $|\mathbf{B}| = 1$. We quote from [26, §8]:

- If the magnetic field \mathbf{B} is tangent to a face but not to an edge of Π , there exists an edge \mathbf{e} such that $\mathcal{E}^*(\mathbf{B}, \Pi) = E(\mathbf{B}, \Pi_{\mathbf{e}})$ and there holds $E(\mathbf{B}, \Pi) < E(\mathbf{B}, \Pi_{\mathbf{e}})$.
- If the magnetic field \mathbf{B} is tangent to an edge \mathbf{e} of Π , $\mathcal{E}^*(\mathbf{B}, \Pi) = E(\mathbf{B}, \Pi_{\mathbf{e}}) = E(\mathbf{B}, \Pi)$. Moreover by [26, §4], $E(\mathbf{B}, \Pi_{\mathbf{e}}) = E(1, \mathcal{S}_{\pi/2}) < \Theta_0 = \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{e}})$.

4.8. Structure of the admissible generalized eigenvectors. We have described in the previous Sections 4.4-4.7 admissible generalized eigenvectors in every situation. In this section we list the model configurations (\mathbf{B}, Π) owning admissible generalized eigenvectors and give a comprehensive overview of their structure in a table. We also prove some stability properties of the generalized eigenvectors and the associated energy under perturbation of the magnetic field \mathbf{B} .

Let \mathbf{B} be a constant magnetic field and Π a cone in $\overline{\mathfrak{P}}_3$. Let us assume that $E(\mathbf{B}, \Pi) < \mathcal{E}^*(\mathbf{B}, \Pi)$. Therefore by Theorem 4.7 there exist admissible generalized eigenvectors Ψ that have the form (4.6). We recall the discriminant parameter $k \in \{1, 2, 3\}$ which is the number of directions in which the generalized eigenvector has an exponential decay. For further use we call (G1), (G2), and (G3) the situation where $k = 1, 2$, and 3, respectively.

In Table 1 we gather all possible situations for (k, d) where d is the dimension of the reduced cone of Π . We assume that the magnetic field \mathbf{B} is unitary, similar formulas can be found using Lemma 4.1 for any non-zero constant magnetic field. We provide the explicit form of an admissible generalized eigenfunction Ψ of $H(\underline{\mathbf{A}}, \Pi)$ in variables $(\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \Upsilon$ where $\underline{\mathbf{A}}$ is a model linear potential associated with \mathbf{B} in these variables. We also give the cone Υ on which the generalized eigenfunction has exponential decay (note that Υ does not always coincide with the reduced cone Γ of Π).

Remark 4.18. In the case where Π is a half-space and \mathbf{B} is normal to $\partial\Pi$, we have $E(\mathbf{B}, \Pi) = \mathcal{E}^*(\mathbf{B}, \Pi) = 1$ and we are in case (ii) of Theorem 4.7, therefore there exists an admissible generalized eigenvector for the strict singular chain \mathbb{R}^3 of Π . However there also exists an admissible generalized eigenfunction for the operator on the half-plane Π . Let $\mathbf{z} = (z_1, z_2)$ be coordinates of $\partial\Pi$ and y coordinate normal to $\partial\Pi$. Let $\underline{\mathbf{A}}(y, z_1, z_2) := (0, -\frac{z_2}{2}, \frac{z_1}{2})$. As described in [22, Lemma 4.3], the function $\Psi : (y, z_1, z_2) \mapsto e^{-(z_1^2+z_2^2)/4}$ is an admissible generalized eigenvector for $H(\underline{\mathbf{A}}, \Pi)$ associated with 1, indeed it satisfies the Neumann boundary condition at the boundary $y = 0$ since it is constant in the y direction and it is a solution of the eigenvalue equation $H(\underline{\mathbf{A}}, \Pi)\Psi = \Psi$, see Section 4.4.

(k, d)	Model (\mathbf{B}, Π)	Potential \mathbf{A}	Υ	Explicit Ψ	Φ eigenvector of
(3,3)	_____	_____	$\Pi = \Gamma$	$\Phi(\mathbf{z})$	$H(\mathbf{A}, \Pi)$
(2,2)	(b_0, b_1, b_2) $\Pi = \mathbb{R} \times \mathcal{S}_\alpha$	$(b_1 z_2 - b_2 z_1, 0, b_0 z_1)$	$\mathcal{S}_\alpha = \Gamma$	$e^{i\tau^* y} \Phi(\mathbf{z})$	$\widehat{H}_\tau(\mathbf{A}, \mathcal{W}_\alpha)$, cf (4.14)
(2,1)	$(0, b_1, b_2), b_2 \neq 0$ $\Pi = \mathbb{R}^2 \times \mathbb{R}_+$	$(b_1 z_2 - b_2 z_1, 0, 0)$	$\mathbb{R} \times \mathbb{R}_+$	$\Phi(\mathbf{z})$	$-\Delta_{\mathbf{z}} + (b_1 z_2 - b_2 z_1)^2$
(2,0)	$(1, 0, 0)$ $\Pi = \mathbb{R}^3$	$(0, -\frac{1}{2}z_2, \frac{1}{2}z_1)$	\mathbb{R}^2	$e^{- \mathbf{z} ^2/4}$	$-\Delta_{\mathbf{z}} + i\mathbf{z} \times \nabla_{\mathbf{z}} + \frac{ \mathbf{z} ^2}{4}$
(1,1)	$(0, 1, 0)$ $\Pi = \mathbb{R}^2 \times \mathbb{R}_+$	$(z, 0, 0)$	$\mathbb{R}_+ = \Gamma$	$e^{-iy_1 \sqrt{\Theta_0}} \Phi(\mathbf{z})$	$-\partial_z^2 + (z - \sqrt{\Theta_0})^2$

Table 1. Generalized eigenfunctions of $H(\mathbf{A}, \Pi)$ depending on the geometry (\mathbf{B}, Π) with $E(\mathbf{B}, \Pi) < \mathcal{E}^*(\mathbf{B}, \Pi)$ written in variables $(\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \Upsilon$.

A perturbation of the magnetic field has distinct effects according to the situation : (G1) is not stable whereas (G2) is. We prove this in the following lemma together with local uniform estimates for exponential decay.

Lemma 4.19. *Let \mathbf{B}_0 be a non zero constant magnetic field and Π be a cone in $\overline{\mathfrak{P}}_3$ with $d < 3$. Assume that $E(\mathbf{B}_0, \Pi) < \mathcal{E}^*(\mathbf{B}_0, \Pi)$.*

(a) *In a ball $\mathcal{B}(\mathbf{B}_0, \varepsilon)$, the function $\mathbf{B} \mapsto E(\mathbf{B}, \Pi)$ is Lipschitz-continuous and*

$$E(\mathbf{B}, \Pi) < \mathcal{E}^*(\mathbf{B}, \Pi).$$

(b) *We suppose moreover that (\mathbf{B}_0, Π) is in situation (G2). For $\mathbf{B} \in \mathcal{B}(\mathbf{B}_0, \varepsilon)$, we denote by $\Psi^{\mathbf{B}}$ an admissible generalized eigenfunction given by Theorem 4.7. For ε small enough, (\mathbf{B}, Π) is still in situation (G2) and $\Psi^{\mathbf{B}}$ has the form*

$$\Psi^{\mathbf{B}}(\mathbf{x}) = e^{i\varphi^{\mathbf{B}}(\mathbf{y}, \mathbf{z})} \Phi^{\mathbf{B}}(\mathbf{z}) \quad \text{for } \underline{\mathbf{U}}_{\mathbf{B}} \mathbf{x} = (\mathbf{y}, \mathbf{z}) \in \mathbb{R} \times \Upsilon,$$

with $\underline{\mathbf{U}}_{\mathbf{B}}$ a suitable rotation, and there exist constants $c > 0$ and $C > 0$ such that

$$(4.20) \quad \forall \mathbf{B} \in \mathcal{B}(\mathbf{B}_0, \varepsilon), \quad \|\Phi^{\mathbf{B}} e^{c|\mathbf{z}|}\|_{L^2(\Upsilon)} \leq C \|\Phi^{\mathbf{B}}\|_{L^2(\Upsilon)}.$$

Proof. Let us distinguish the three possible situations according to the value of d :

$d = 0$: When $\Pi = \mathbb{R}^3$, we have $E(\mathbf{B}, \Pi) = |\mathbf{B}|$ and $\mathcal{E}^*(\mathbf{B}, \Pi) = +\infty$. The admissible generalized eigenvector $\Psi^{\mathbf{B}}$ is explicit as explained above. Thus (a) and (b) are established in this case.

$d = 1$: When Π is a half-space, we denote by $\theta(\mathbf{B})$ the unoriented angle in $[0, \frac{\pi}{2}]$ between \mathbf{B} and the boundary. The function $\mathbf{B} \mapsto \theta(\mathbf{B})$ is Lipschitz. Moreover the function σ is \mathcal{C}^1 on $[0, \pi/2]$ (see Lemma 4.9). We deduce that the function $\mathbf{B} \mapsto \sigma(\theta(\mathbf{B}))$ is Lipschitz outside any neighborhood of $\mathbf{B} = 0$. Thus point (a) is proved. Assuming

furthermore that (Π, \mathbf{B}_0) is in situation (G2), we have $\theta(\mathbf{B}_0) \in (0, \frac{\pi}{2})$ and there exist $\varepsilon > 0$, θ_{\min} and θ_{\max} such that

$$\forall \mathbf{B} \in \mathcal{B}(\mathbf{B}_0, \varepsilon), \quad \theta(\mathbf{B}) \in [\theta_{\min}, \theta_{\max}] \subset (0, \frac{\pi}{2}).$$

The admissible generalized eigenvector is constructed above. The uniform exponential estimate is proved in [7, §2].

$d = 2$: When Π is a wedge, point (a) is proved in [30, §4]. Due to the continuity of $\mathbf{B} \mapsto E(\mathbf{B}, \Pi)$ there exist $c > 0$ and $\varepsilon > 0$ such that

$$\forall \mathbf{B} \in \mathcal{B}(\mathbf{B}_0, \varepsilon), \quad \mathcal{E}^*(\mathbf{B}, \Pi) - E(\mathbf{B}, \Pi) > c.$$

Point (b) is then a direct consequence of [30, Proposition 4.2].

The proof of Lemma 4.19 is complete. □

5. Continuity properties of the ground energy

Let $\Omega \in \overline{\mathcal{D}}(\mathbb{R}^3)$ and let $\mathbf{B} \in \mathcal{C}^0(\overline{\Omega})$ be a continuous magnetic field. In this section we investigate the continuity properties on $\overline{\Omega}$ of the application $\Lambda : \mathbf{x} \mapsto E(\mathbf{B}_x, \Pi_x)$. Let \mathbf{t} be a stratum of $\overline{\Omega}$ (see (1.10)). We have denoted by $\Lambda_{\mathbf{t}}$ the restriction of Λ to \mathbf{t} (see (1.11)). Combining (4.2), (4.9), Lemma 4.9 and Lemma 4.11 we get that $\Lambda_{\mathbf{t}}$ is continuous.

Let us assume that \mathbf{t} is not reduced to a point. We now describe how we extend $\Lambda_{\mathbf{t}}$ to the boundary of \mathbf{t} . Let $\mathbf{x} \in \partial\mathbf{t}$ and $\Pi_x \in \overline{\mathfrak{P}}_3$ be its tangent cone. Let \mathcal{U}_x , \mathcal{V}_x and U_x be the open sets and the diffeomorphism introduced in Section 3.1. Let $\tilde{\mathbf{t}}$ be the stratum of Π_x such that

$$U_x(\mathbf{t} \cap \mathcal{U}_x) = \tilde{\mathbf{t}} \cap \mathcal{V}_x.$$

To $\tilde{\mathbf{t}}$ is associated the singular chain $\mathbb{X} \in \mathcal{C}_0^*(\Pi_x)$ such that $\Pi_{\mathbb{X}}$ is the tangent cone to Π_x at any point of $\tilde{\mathbf{t}}$.

We extend $\Lambda_{\mathbf{t}}$ in \mathbf{x} by setting

$$(5.1) \quad \Lambda_{\mathbf{t}}(\mathbf{x}) = E(\mathbf{B}_x, \Pi_{\mathbb{X}}).$$

Lemma 5.1. *Let $\Omega \in \overline{\mathcal{D}}(\mathbb{R}^3)$ and let $\mathbf{B} \in \mathcal{C}^0(\overline{\Omega})$. Let \mathbf{t} a stratum of Ω which is not a vertex. Then formula (5.1) defines a continuous extension of the function $\Lambda_{\mathbf{t}}$ to $\tilde{\mathbf{t}}$.*

Proof. For $\mathbf{x} \in \partial\mathbf{t}$ we show that the extension defined by (5.1) is continuous in \mathbf{x} . Let $\mathbf{y} \in \mathcal{U}_x \cap \mathbf{t}$ and Π_y be the tangent cone to Ω at \mathbf{y} . In the following we will prove that

$$(5.2) \quad \lim_{\mathbf{y} \rightarrow \mathbf{x}} E(\mathbf{B}_y, \Pi_y) = E(\mathbf{B}_x, \Pi_{\mathbb{X}}).$$

For a tangent cone Π we denote by $d(\Pi)$ the dimension of its reduced cone. Since \mathbf{t} is not reduced to a point, we have $d(\Pi_{\mathbb{X}}) = d(\Pi_y) \leq 2$ and we distinguish several cases:

- $d = 0$. $\Pi_{\mathbb{X}} = \Pi_y = \mathbb{R}^3$. It follows from Section 4.4 that $E(\mathbf{B}_y, \Pi_y) = |\mathbf{B}_y|$ and $E(\mathbf{B}_x, \Pi_{\mathbb{X}}) = |\mathbf{B}_x|$. Therefore (5.2).

- $d = 1$. $\Pi_{\mathbf{x}}$ and $\Pi_{\mathbf{y}}$ are half-spaces. We denote by $\theta_{\mathbf{x}}$ (respectively $\theta_{\mathbf{y}}$) the angle between $\mathbf{B}_{\mathbf{x}}$ and $\Pi_{\mathbf{x}}$ (respectively $\mathbf{B}_{\mathbf{y}}$ and $\Pi_{\mathbf{y}}$). We have $E(\mathbf{B}_{\mathbf{y}}, \Pi_{\mathbf{y}}) = |\mathbf{B}_{\mathbf{y}}|\sigma(\theta_{\mathbf{y}})$ and $E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}}) = |\mathbf{B}_{\mathbf{x}}|\sigma(\theta_{\mathbf{x}})$ (see Section 4.5). Since $\theta_{\mathbf{y}}$ goes to $\theta_{\mathbf{x}}$ when \mathbf{y} goes to \mathbf{x} , (5.2) follows from the continuity of the function σ , see Lemma 4.9.
- $d = 2$. $\Pi_{\mathbf{x}}$ and $\Pi_{\mathbf{y}}$ are wedges. We denote by $\alpha_{\mathbf{x}}$ and $\alpha_{\mathbf{y}}$ their openings. We denote by $\underline{U}_{\mathbf{x}} \in \mathfrak{D}_3$ (respectively $\underline{U}_{\mathbf{y}} \in \mathfrak{D}_3$) the linear orthogonal transformation which maps $\Pi_{\mathbf{x}}$ on $\mathcal{W}_{\alpha_{\mathbf{x}}}$ (respectively $\Pi_{\mathbf{y}}$ on $\mathcal{W}_{\alpha_{\mathbf{y}}}$). We have

$$(5.3) \quad E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}}) = E(\underline{\mathbf{B}}_{\mathbf{x}}, \mathcal{W}_{\alpha_{\mathbf{x}}}) \quad \text{and} \quad E(\mathbf{B}_{\mathbf{y}}, \Pi_{\mathbf{y}}) = E(\underline{\mathbf{B}}_{\mathbf{y}}, \mathcal{W}_{\alpha_{\mathbf{y}}}).$$

where we have denoted $\underline{\mathbf{B}}_{\mathbf{x}} = \underline{U}_{\mathbf{x}}(\mathbf{B}_{\mathbf{x}})$ and $\underline{\mathbf{B}}_{\mathbf{y}} = \underline{U}_{\mathbf{y}}(\mathbf{B}_{\mathbf{y}})$. We have

$$\lim_{\mathbf{y} \rightarrow \mathbf{x}} \|\underline{U}_{\mathbf{x}} - \underline{U}_{\mathbf{y}}\| = 0 \quad \text{and} \quad \lim_{\mathbf{y} \rightarrow \mathbf{x}} |\alpha_{\mathbf{x}} - \alpha_{\mathbf{y}}| = 0,$$

therefore we deduce (5.2) from (5.3) and Lemma 4.11.

Hence we have proved (5.2) in all cases. □

Let $\mathbf{x} \in \partial\mathbf{t}$, we deduce from (4.8) that

$$\Lambda_{\mathbf{t}}(\mathbf{x}) = E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}}) \geq E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}}) = \Lambda(\mathbf{x}).$$

Combining this with Lemma 5.1, we obtain the following:

Theorem 5.2. *Let $\Omega \in \overline{\mathfrak{D}}(\mathbb{R}^3)$ and let $\mathbf{B} \in \mathcal{C}^0(\overline{\Omega})$ be a continuous magnetic field. Then the function $\Lambda : \mathbf{x} \mapsto E(\mathbf{B}_{\mathbf{x}}, \Pi_{\mathbf{x}})$ is lower semi-continuous on $\overline{\Omega}$.*

6. Upper bound for first eigenvalues in polyhedral domains

In this section we prove general upper bounds for the first eigenvalue $\lambda_h(\mathbf{B}, \Omega)$ by $h\mathcal{E}(\mathbf{B}, \Omega)$ up to a remainder of size h^κ with $\kappa > 1$. The first theorem provides an upper bound with $\kappa = \frac{5}{4}$ using a novel construction of quasimodes depending on the geometry. Under more regularity assumptions on the potential and using more knowledge on the model problems we refine the quasimodes and we reach an upper bound with $\kappa = \frac{4}{3}$.

Here is our first result:

Theorem 6.1. *Let $\Omega \in \overline{\mathfrak{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{A} \in W^{2,\infty}(\overline{\Omega})$ be a twice differentiable magnetic potential such that the associated magnetic field \mathbf{B} does not vanish on $\overline{\Omega}$. Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that*

$$(6.1) \quad \forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \leq h\mathcal{E}(\mathbf{B}, \Omega) + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) h^{5/4}.$$

We recall that the quantity $\mathcal{E}(\mathbf{B}, \Omega)$ is the lowest local energy defined in (1.5).

It is possible to obtain an upper bound in (6.1) depending on the magnetic field \mathbf{B} and not on the magnetic potential. For this, we consider \mathbf{B} as a datum and associate a potential \mathbf{A} with it. Operators $\mathcal{A} : \mathbf{B} \mapsto \mathbf{A}$ lifting the curl (i.e. such that $\text{curl} \circ \mathcal{A} = \mathbb{I}$) and satisfying

suitable estimates do exist in the literature. We quote [8] in which it is proved that such lifting can be constructed as a pseudo-differential operator of order -1 . As a consequence \mathcal{A} is continuous between Hölder classes of non integer order:

$$\forall \alpha \in (0, 1), \quad \exists C_\alpha > 0, \quad \|\mathcal{A}\mathbf{B}\|_{W^{2+\alpha,\infty}(\Omega)} \leq C_\alpha \|\mathbf{B}\|_{W^{1+\alpha,\infty}(\Omega)}.$$

Choosing $\mathbf{A} = \mathcal{A}\mathbf{B}$ in Theorem 6.1, we deduce the following.

Corollary 6.2. *Let $\Omega \in \overline{\mathcal{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{B} \in W^{1+\alpha,\infty}(\overline{\Omega})$ be a non-vanishing Hölder continuous magnetic field of order $1 + \alpha$ with some $\alpha \in (0, 1)$. Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that*

$$(6.2) \quad \forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \leq h \mathcal{E}(\mathbf{B}, \Omega) + C(\Omega)(1 + \|\mathbf{B}\|_{W^{1+\alpha,\infty}(\Omega)}^2) h^{5/4}.$$

Theorem 6.1 is proved in Sections 6.1–6.3 according to the following plan. Since the energy $\mathbf{x} \mapsto E(\mathbf{B}_\mathbf{x}, \Pi_\mathbf{x})$ is lower semi-continuous (see Theorem 5.2), it reaches its infimum over the compact $\overline{\Omega}$. We denote by $\mathbf{x}_0 \in \overline{\Omega}$ a point such that

$$(6.3) \quad E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) = \mathcal{E}(\mathbf{B}, \Omega)$$

where $\Pi_{\mathbf{x}_0}$ is the tangent cone at \mathbf{x}_0 . Starting from this, the proof is organized in three main steps, developed in Sections 6.1, 6.2, and 6.3 respectively:

- (1) By the local diffeomorphism $U_{\mathbf{x}_0}$ in the neighborhood $\mathcal{U}_{\mathbf{x}_0}$ of \mathbf{x}_0 introduced in (3.1), we reduce to a local magnetic operator set on the tangent cone $\Pi_{\mathbf{x}_0}$.
- (2) We construct quasimodes for the tangent magnetic operator at the vertex $\mathbf{0}$ of $\Pi_{\mathbf{x}_0}$. Here we use Theorem 4.7 which provides us with a suitable admissible generalized eigenfunction associated with the energy $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$. This generalized eigenfunction will be scaled, truncated and translated in order to give a quasimode for the local magnetic operator.
- (3) The estimation of various terms in the associated Rayleigh quotient and the min-max principle will finally prove Theorem 6.1.

If steps (1) and (3) are very classical, step (2) reveals much more originality, because our constructions are valid in any configuration: We do not need to know *a priori* whether $(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$ is in situation (i) or (ii) of the dichotomy theorem. If we are in situation (i) – the most classical one – our quasimodes will be classical too, and qualified as “sitting”. If we are in situation (ii), we define our quasimodes on a higher structure $\Pi_\mathbf{x}$ and make then “slide” towards the vertex $\mathbf{0}$ of $\Pi_{\mathbf{x}_0}$.

Our second result is also general, the unique additional assumption is a supplementary regularity on the magnetic potential (or equivalently on the magnetic field).

Theorem 6.3. *Let $\Omega \in \overline{\mathcal{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{A} \in W^{3,\infty}(\overline{\Omega})$ be a magnetic potential such that the associated magnetic field does not vanish. Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that*

$$(6.4) \quad \forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \leq h \mathcal{E}(\mathbf{B}, \Omega) + C(\Omega)(1 + \|\mathbf{A}\|_{W^{3,\infty}(\Omega)}^2) h^{4/3}.$$

Like for Corollary 6.2, we can deduce an upper bound where the constant depends on the magnetic field.

Corollary 6.4. *Let $\Omega \in \overline{\mathfrak{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{B} \in W^{2+\alpha,\infty}(\overline{\Omega})$ be a non-vanishing Hölder continuous magnetic field of order $2 + \alpha$ with some $\alpha \in (0, 1)$. Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that*

$$(6.5) \quad \forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \leq h \mathcal{E}(\mathbf{B}, \Omega) + C(\Omega)(1 + \|\mathbf{B}\|_{W^{2+\alpha,\infty}(\Omega)}^2) h^{4/3}.$$

Note that the $h^{4/3}$ bound was known for smooth three-dimensional domains, [18, Proposition 6.1 & Remark 6.2] and that our result extends this result to polyhedral domains without loss.

Theorem 6.3 is proved in Section 6.4 and we give now some hint on the proof. Like for Theorem 6.1 we start from admissible generalized eigenfunctions and construct sitting or sliding quasimodes adapted to the geometry. However, unlike for the proof of Theorem 6.1, we are going to actually take advantage of the decaying property of the generalized eigenfunctions and adopt different strategies depending on the number k of directions in which the generalized eigenfunction has exponential decay: A Feynman-Hellmann formula if $k = 1$, a refined Taylor expansion of the potential if $k = 2$, and an Agmon decay estimate if $k = 3$.

6.1. Change of variables. Let us recall from Section 3.1 that the local smooth diffeomorphism $U_{\mathbf{x}_0}$ maps a neighborhood $\mathcal{U}_{\mathbf{x}_0}$ of \mathbf{x}_0 onto a neighborhood $\mathcal{V}_{\mathbf{x}_0}$ of $\mathbf{0}$ so that

$$(6.6) \quad U_{\mathbf{x}_0}(\mathcal{U}_{\mathbf{x}_0} \cap \Omega) = \mathcal{V}_{\mathbf{x}_0} \cap \Pi_{\mathbf{x}_0} \quad \text{and} \quad U_{\mathbf{x}_0}(\mathcal{U}_{\mathbf{x}_0} \cap \partial\Omega) = \mathcal{V}_{\mathbf{x}_0} \cap \partial\Pi_{\mathbf{x}_0}.$$

The differential of $U_{\mathbf{x}_0}$ at the point \mathbf{x}_0 is the identity matrix \mathbb{I}_3 . Let

$$J := d(U_{\mathbf{x}_0}^{-1}) \quad \text{and} \quad G := J^{-1}(J^{-1})^\top$$

be the jacobian matrix of the inverse of $U_{\mathbf{x}_0}$ and the associated metric. Lemma A.4 tells us that the transformed magnetic potential $\tilde{\mathbf{A}}$ and magnetic field $\tilde{\mathbf{B}} = \text{curl } \tilde{\mathbf{A}}$ in $\mathcal{V}_{\mathbf{x}_0} \cap \Pi_{\mathbf{x}_0}$ are given by

$$(6.7) \quad \tilde{\mathbf{A}} := J^\top(\mathbf{A} \circ U_{\mathbf{x}_0}^{-1}) \quad \text{and} \quad \tilde{\mathbf{B}} := |\det J| J^{-1}(\mathbf{B} \circ U_{\mathbf{x}_0}^{-1}).$$

Let moreover f be a function in $H^1(\Omega)$ with support in $\mathcal{U}_{\mathbf{x}_0}$ and $\psi := f \circ U_{\mathbf{x}_0}^{-1}$ defined in $\Pi_{\mathbf{x}_0}$, with support in $\mathcal{V}_{\mathbf{x}_0}$. For any $h > 0$ we have

$$(6.8) \quad q_h[\mathbf{A}, \Omega](f) = q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}, G](\psi) \quad \text{and} \quad \|f\|_{L^2(\Omega)} = \|\psi\|_{L^2_G(\Pi_{\mathbf{x}_0})},$$

where the quadratic forms $q_h[\mathbf{A}, \Omega]$ and $q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}, G]$ are defined in (1.13) and (1.15), respectively.

Let $\tilde{\mathbf{A}}_0$ be the affine part of $\tilde{\mathbf{A}}$ at the vertex $\mathbf{0}$ of $\Pi_{\mathbf{x}_0}$. Since $dU_{\mathbf{x}_0}(\mathbf{x}_0) = \mathbb{I}_3$, the affine part of $\tilde{\mathbf{A}}$ at $\mathbf{0}$ coincides with the affine part of \mathbf{A} at \mathbf{x}_0 :

$$(6.9) \quad \tilde{\mathbf{A}}_0 = \mathbf{A}_{\mathbf{x}_0}.$$

Lemma 6.5. *Let $r_0 > 0$ be such that $\mathcal{V}_{\mathbf{x}_0}$ contains the ball $\mathcal{B}(\mathbf{0}, r_0)$ of center $\mathbf{0}$ and radius r_0 . Then there exists a constant $C(\Omega)$ such that for all $r \in (0, r_0]$, if $\psi \in H^1(\Pi_{\mathbf{x}_0})$ with $\text{supp}(\psi) \subset \mathcal{B}(\mathbf{0}, r)$ we have the two estimates*

$$(6.10) \quad |q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi) - q_h[\mathbf{A}, \Omega](\psi \circ U_{\mathbf{x}_0})| \leq C(\Omega) r q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi),$$

$$(6.11) \quad \left| \|\psi\|_{L^2(\Pi_{\mathbf{x}_0})} - \|\psi \circ U_{\mathbf{x}_0}\|_{L^2(\Omega)} \right| \leq C(\Omega) r \|\psi\|_{L^2(\Pi_{\mathbf{x}_0})}.$$

Proof. Recall that $J_{\mathbf{x}_0}(\mathbf{x}_0) = \mathbb{I}_3$ and therefore $G_{\mathbf{x}_0}(\mathbf{x}_0) = \mathbb{I}_3$. We deduce

$$(6.12) \quad \|G_{\mathbf{x}_0} - \mathbb{I}_3\|_{L^\infty(\mathcal{B}(0,r))} \leq r \|G_{\mathbf{x}_0}\|_{W^{1,\infty}(\mathcal{V}_{\mathbf{x}_0})}.$$

Since Ω is assumed to be polyhedral, its curvature (curvature of the faces and curvature of the edges) is bounded, therefore there exists $C(\Omega)$ such that

$$\|G_{\mathbf{x}_0}\|_{W^{1,\infty}(\mathcal{V}_{\mathbf{x}_0})} \leq C(\Omega), \quad \forall \mathbf{x}_0 \in \bar{\Omega}.$$

Notice that $q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}] = q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}, \mathbb{I}_3]$, we deduce the Lemma by using (6.12) in (6.8). \square

Therefore we are reduced to study the Laplacian with magnetic potential $\tilde{\mathbf{A}}$ on the cone $\Pi_{\mathbf{x}_0}$ with the identity metric.

6.2. Construction of quasimodes. Let $\mathbf{x}_0 \in \bar{\Omega}$ be a point satisfying (6.3). Thus \mathbf{x}_0 minimizes the local ground energy. For shortness we denote by Λ_0 this energy:

$$(6.13) \quad \Lambda_0 = E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}).$$

In order to prove Theorem 6.1, we are going to construct a family of quasimodes $f_h \in H^1(\Omega)$ satisfying the estimate for $h \leq h_0$ (with some chosen positive h_0)

$$(6.14) \quad \frac{q_h[\mathbf{A}, \Omega](f_h)}{\|f_h\|^2} \leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)h^{5/4}.$$

We recall that $\mathbf{A}_{\mathbf{x}_0}$ is the affine part of \mathbf{A} at \mathbf{x}_0 . Let $\tilde{\mathbf{A}}$ be the magnetic potential in the tangent cone $\Pi_{\mathbf{x}_0} \cap \mathcal{V}_{\mathbf{x}_0}$ given by (6.7). We recall that since $dU_{\mathbf{x}_0}(\mathbf{x}_0) = \mathbb{I}_3$, the affine part $\tilde{\mathbf{A}}_0$ of $\tilde{\mathbf{A}}$ at $\mathbf{0}$ coincides with $\mathbf{A}_{\mathbf{x}_0}$.

Theorem 4.7 provides a singular chain⁴ \mathbb{X} in $\mathfrak{C}_{\mathbf{x}_0}(\Omega) \equiv \mathfrak{C}_0(\Pi_{\mathbf{x}_0})$ and its associated cone $\Pi_{\mathbb{X}}$ such that the operator $H(\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}})$ has an admissible generalized eigenfunction associated with the energy $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$.

According as $\Pi_{\mathbf{x}_0}$ equals $\Pi_{\mathbb{X}}$ or not (cases (i) or (ii) in Theorem 4.7, respectively), our constructions will be different, leading to two types of quasimodes qualified as “sitting” or “sliding”. For this, we need to introduce a new geometrical object $\boldsymbol{\tau} \in \mathcal{B}(0, 1)$:

Definition 6.6. We define the vector $\boldsymbol{\tau} \in \mathcal{B}(0, 1)$ according to the dichotomy given by Theorem 4.7:

⁴In the case (i) of Theorem 4.7, \mathbb{X} is the trivial chain (\mathbf{x}_0) and $\Pi_{\mathbb{X}} = \Pi_{\mathbf{x}_0}$.

- (i) If $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) < \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$, then $\mathbb{X} = (\mathbf{x}_0)$ and $\Pi_{\mathbb{X}} = \Pi_{\mathbf{x}_0}$. We set $\boldsymbol{\tau} = 0$.
- (ii) If $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) = \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$, then $\mathbb{X} \in \mathcal{C}_{\mathbf{x}_0}^*(\Omega)$. Let $\underline{U}_0 \in \mathfrak{D}_3$ such that $\Pi_{\mathbf{x}_0} = \underline{U}_0(\mathbb{R}^{3-d} \times \Gamma)$ where Γ is the reduced cone of $\Pi_{\mathbf{x}_0}$. Let $\Omega_0 = \Gamma \cap \mathbb{S}^{d-1}$ be the section of Γ . According to Remark 3.3, there exists $\mathbf{x}_1 \in \overline{\Omega}_0$ so that $\mathbb{X} = (\mathbf{x}_0, \mathbf{x}_1)$. Let $\underline{\boldsymbol{\tau}} := (0, \mathbf{x}_1) \in \mathbb{R}^{3-d} \times \Gamma$. We set $\boldsymbol{\tau} = \underline{U}_0 \underline{\boldsymbol{\tau}} \in \mathbb{S}^2$.

Remark 6.7. In case (ii), the cone $\Pi_{\mathbb{X}}$ can be the full space, a half-space or a wedge, and $\boldsymbol{\tau}$ gives a direction associated with $\Pi_{\mathbb{X}}$ starting from the origin of $\Pi_{\mathbf{x}_0}$:

- (1) If $\Pi_{\mathbb{X}} \equiv \mathbb{R}^3$, $\boldsymbol{\tau}$ belongs to the interior of $\Pi_{\mathbf{x}_0}$.
- (2) If $\Pi_{\mathbb{X}} \equiv \mathbb{R}_+^3$, $\boldsymbol{\tau}$ belongs to a face of $\Pi_{\mathbf{x}_0}$.
- (3) If $\Pi_{\mathbb{X}} \equiv \mathcal{W}_\alpha$, $\boldsymbol{\tau}$ belongs to an edge of $\Pi_{\mathbf{x}_0}$.

Note that unless we are in the latter case ($\Pi_{\mathbb{X}}$ is a wedge), the choice of $\boldsymbol{\tau}$ is not unique.

Let Ψ be an admissible generalized eigenvector for $H(\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}})$ associated with Λ_0 .

There exists a rotation \underline{U} which transforms $\Pi_{\mathbb{X}}$ into $\mathbb{R}^{3-k} \times \Upsilon$ so that in coordinates $(\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \Upsilon$, with $\Upsilon \in \overline{\mathfrak{P}}_k$, the generalized eigenfunction writes:

$$(6.15) \quad \Psi(\mathbf{x}) = e^{i\varphi(\mathbf{y}, \mathbf{z})} \Phi(\mathbf{z}) \quad \text{with} \quad \underline{U}\mathbf{x} = (\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \Upsilon,$$

where Φ is an exponentially decreasing function. The function Ψ satisfies

$$(6.16) \quad \begin{cases} (-i\nabla + \tilde{\mathbf{A}}_0)^2 \Psi = \Lambda_0 \Psi & \text{in } \Pi_{\mathbb{X}}, \\ (-i\partial_n + \mathbf{n} \cdot \tilde{\mathbf{A}}_0) \Psi = 0 & \text{on } \partial\Pi_{\mathbb{X}}. \end{cases}$$

Then the scaled function

$$(6.17) \quad \Psi_h(\mathbf{x}) := \Psi\left(\frac{\mathbf{x}}{\sqrt{h}}\right), \quad \text{for } \mathbf{x} \in \Pi_{\mathbb{X}},$$

defines a generalized eigenfunction for the operator $H_h(\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}})$ associated with $h\Lambda_0$.

Let us choose, once for all, a model cut-off function $\underline{\chi} \in \mathcal{C}^\infty(\mathbb{R}^+)$ such that

$$(6.18) \quad \underline{\chi}(r) = \begin{cases} 1 & \text{if } r \leq 1, \\ 0 & \text{if } r \geq 2. \end{cases}$$

For any $R > 0$, let $\underline{\chi}_R$ be the cut-off function defined by

$$(6.19) \quad \underline{\chi}_R(r) = \underline{\chi}\left(\frac{r}{R}\right),$$

and, finally

$$(6.20) \quad \chi_h(\mathbf{x}) = \underline{\chi}_R\left(\frac{|\mathbf{x}|}{h^\delta}\right) = \underline{\chi}\left(\frac{|\mathbf{x}|}{Rh^\delta}\right) \quad \text{with} \quad 0 \leq \delta \leq \frac{1}{2}.$$

We set

$$(6.21) \quad \psi_h(\mathbf{x}) := \chi_h(\mathbf{x})\Psi_h(\mathbf{x})$$

which provides a quasimode for $q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}}]$ satisfying $(-ih\partial_n + \mathbf{n} \cdot \tilde{\mathbf{A}}_0)\psi_h = 0$ on $\partial\Pi_{\mathbb{X}}$.

At this point, let us emphasize that, in order to obtain better cut-off estimates, cf. Lemma A.8, we need that our quasimodes on the tangent cone $\Pi_{\mathbf{x}_0}$ satisfy the Neumann boundary conditions. Here the vector $\boldsymbol{\tau}$ introduced in Definition 6.6 comes into play and we define our quasimode by

$$(6.22) \quad \psi_h^{\boldsymbol{\tau}}(\mathbf{x}) = e^{i\langle \tilde{\mathbf{A}}_0(\mathbf{0}) - \tilde{\mathbf{A}}_0(\boldsymbol{\tau}h^\delta), \mathbf{x}/h \rangle} \psi_h(\mathbf{x} - \boldsymbol{\tau}h^\delta), \quad \mathbf{x} \in \Pi_{\mathbf{x}_0}.$$

We check that

$$(6.23) \quad (-ih\nabla + \tilde{\mathbf{A}}_0(\mathbf{x}))\psi_h^{\boldsymbol{\tau}}(\mathbf{x}) = e^{i\langle \tilde{\mathbf{A}}_0(\mathbf{0}) - \tilde{\mathbf{A}}_0(\boldsymbol{\tau}h^\delta), \mathbf{x}/h \rangle} (-ih\nabla + \tilde{\mathbf{A}}_0(\mathbf{x} - \boldsymbol{\tau}h^\delta))\psi_h(\mathbf{x} - \boldsymbol{\tau}h^\delta).$$

In case (i), note that $\psi_h^{\boldsymbol{\tau}} = \psi_h$. We choose $R = 1$ in (6.19).

In case (ii), we choose $R > 0$ in (6.19) such that $\mathcal{B}(\boldsymbol{\tau}, 2R) \cap \Pi_{\mathbf{x}_0} = \mathcal{B}(\boldsymbol{\tau}, 2R) \cap \Pi_{\mathbb{X}}$. Hence

$$(6.24) \quad \text{supp}(\underline{\chi}_R(\cdot - \boldsymbol{\tau})) \cap \Pi_{\mathbf{x}_0} = \text{supp}(\underline{\chi}_R(\cdot - \boldsymbol{\tau})) \cap \Pi_{\mathbb{X}}.$$

Note that R depends only on the geometry of Ω near \mathbf{x}_0 . It follows by scaling that

$$\forall h > 0, \quad \text{supp}(\psi_h^{\boldsymbol{\tau}}) \cap \Pi_{\mathbf{x}_0} = \text{supp}(\psi_h^{\boldsymbol{\tau}}) \cap \Pi_{\mathbb{X}}.$$

Therefore $\psi_h^{\boldsymbol{\tau}}$ satisfies Neumann boundary conditions on $\partial\Pi_{\mathbf{x}_0}$ and we have

$$(6.25) \quad q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbf{x}_0}](\psi_h^{\boldsymbol{\tau}}) = q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}}](\psi_h^{\boldsymbol{\tau}}).$$

In virtue of (6.23) and the translation invariance of $\Pi_{\mathbb{X}}$ by $\boldsymbol{\tau}$, we have

$$(6.26) \quad q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}}](\psi_h^{\boldsymbol{\tau}}) = q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}}](\psi_h).$$

Definition 6.8. The functions $\psi_h^{\boldsymbol{\tau}} \in \text{Dom}(H_h(\tilde{\mathbf{A}}_0, \Pi_{\mathbf{x}_0}))$ constructed above along formulas (6.15)-(6.24) and defined by

$$(6.27) \quad \psi_h^{\boldsymbol{\tau}}(\mathbf{x}) = e^{i\langle \tilde{\mathbf{A}}_0(\mathbf{0}) - \tilde{\mathbf{A}}_0(\boldsymbol{\tau}h^\delta), \mathbf{x}/h \rangle} \underline{\chi}_R\left(\frac{|\mathbf{x} - \boldsymbol{\tau}h^\delta|}{h^\delta}\right) \Psi\left(\frac{\mathbf{x} - \boldsymbol{\tau}h^\delta}{h^{1/2}}\right), \quad \mathbf{x} \in \Pi_{\mathbf{x}_0},$$

are called

- (i) *sitting quasimodes* if $\boldsymbol{\tau} = 0$: Their supports contain the vertex $\mathbf{0}$ of $\Pi_{\mathbf{x}_0}$,
- (ii) *sliding quasimodes* if $\boldsymbol{\tau} \neq 0$: Their supports do not contain $\mathbf{0}$ but get closer as $h \rightarrow 0$.

The localization of the sliding quasimodes can be deduced from Remark 6.7:

- (1) If $\Pi_{\mathbb{X}} \equiv \mathbb{R}^3$, $\psi_h^{\boldsymbol{\tau}}$ is centered in the interior of $\Pi_{\mathbf{x}_0}$.
- (2) If $\Pi_{\mathbb{X}} \equiv \mathbb{R}_+^3$, $\psi_h^{\boldsymbol{\tau}}$ is centered on the face of $\Pi_{\mathbf{x}_0}$ associated to $\boldsymbol{\tau}$.
- (3) If $\Pi_{\mathbb{X}} \equiv \mathcal{W}_\alpha$, $\psi_h^{\boldsymbol{\tau}}$ is centered on the edge of $\Pi_{\mathbf{x}_0}$ associated to $\boldsymbol{\tau}$.

6.3. Estimation of the quasimodes. We separately estimate the cut-off errors, the linearization errors, and the error due to the change of metric.

6.3.1. *Cut-off effect.* Applying Lemma A.8 with $\tilde{\mathbf{A}}_0$, $G = \mathbb{I}_3$, $\chi = \chi_h$ and $\psi = \Psi_h$, we obtain for the Rayleigh quotient of ψ_h^τ :

$$(6.28) \quad \frac{q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbf{x}_0}](\psi_h^\tau)}{\|\psi_h^\tau\|^2} = \frac{q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbb{X}}](\chi_h \Psi_h)}{\|\chi_h \Psi_h\|^2} = h\Lambda_0 + h^2\rho_h \quad \text{with} \quad \rho_h = \frac{\|\nabla\chi_h \Psi_h\|^2}{\|\chi_h \Psi_h\|^2}.$$

The following lemma estimates the remainder due to the cut-off effect:

Lemma 6.9. *Let Ψ be an admissible generalized eigenvector given by (6.15) and Ψ_h the rescaled associated function given by (6.17). Let χ_h be the cut-off function defined by (6.20) involving parameters $R > 0$ and $\delta \in [0, \frac{1}{2}]$. Then there exist constants $C_0 > 0$ and $c_0 > 0$ depending only on $h_0 > 0$, $R_0 > 0$ and Ψ such that*

$$\rho_h = \frac{\|\nabla\chi_h \Psi_h\|^2}{\|\chi_h \Psi_h\|^2} \leq \begin{cases} C_0 h^{-2\delta} & \text{if } k < 3, \\ C_0 h^{-2\delta} e^{-c_0 h^{\delta-1/2}} & \text{if } k = 3, \end{cases} \quad \forall R \geq R_0, \forall h \leq h_0, \forall \delta \in [0, \frac{1}{2}].$$

Proof. By assumption $\Psi(\mathbf{x}) = e^{i\varphi(\mathbf{y}, \mathbf{z})} \Phi(\mathbf{z})$ for $\underline{\mathbf{x}} = (\mathbf{y}, \mathbf{z}) \in \mathbb{R}^{3-k} \times \Upsilon$ and there exist positive constants c_Ψ, C_Ψ such that

$$(6.29) \quad \int_{\Upsilon} e^{2c_\Psi|\mathbf{z}|} |\Phi(\mathbf{z})|^2 d\mathbf{z} \leq C_\Psi \|\Phi\|_{L^2(\Upsilon)}^2.$$

Let us set $T = Rh^\delta$, so that $\chi_h(\mathbf{x}) = \underline{\chi}(|\mathbf{x}|/T)$.

Let us first give an upper bound for $\|\nabla\chi_h \Psi_h\|$:

If $k < 3$, then

$$\begin{aligned} \|\nabla\chi_h \Psi_h\|^2 &\leq CT^{-2} \int_{|\mathbf{y}| \leq 2T} d\mathbf{y} \int_{\Upsilon \cap \{|\mathbf{z}| \leq 2T\}} \left| \Phi\left(\frac{\mathbf{z}}{\sqrt{h}}\right) \right|^2 d\mathbf{z} \\ &\leq CT^{-2} T^{3-k} h^{k/2} \|\Phi\|_{L^2(\Upsilon)}^2, \end{aligned}$$

else, if $k = 3$

$$\begin{aligned} \|\nabla\chi_h \Psi_h\|^2 &\leq CT^{-2} \int_{\Upsilon \cap \{T \leq |\mathbf{z}| \leq 2T\}} \left| \Phi\left(\frac{\mathbf{z}}{\sqrt{h}}\right) \right|^2 d\mathbf{z} \\ &\leq CT^{-2} h^{k/2} \int_{\Upsilon \cap \{Th^{-\frac{1}{2}} \leq |\mathbf{z}| \leq 2Th^{-\frac{1}{2}}\}} |\Phi(\mathbf{z})|^2 d\mathbf{z} \\ &\leq CT^{-2} h^{k/2} e^{-2c_\Psi T/\sqrt{h}} \int_{\Upsilon \cap \{Th^{-\frac{1}{2}} \leq |\mathbf{z}| \leq 2Th^{-\frac{1}{2}}\}} e^{2c|\mathbf{z}|} |\Phi(\mathbf{z})|^2 d\mathbf{z} \\ &\leq CT^{-2} h^{k/2} e^{-2c_\Psi T/\sqrt{h}} \|\Phi\|_{L^2(\Upsilon)}^2. \end{aligned}$$

Let us now consider $\|\chi_h \Psi_h\|$ (we use that $2|\mathbf{y}| < R$ and $2|\mathbf{z}| < R$ implies $|\mathbf{x}| < R$):

$$\begin{aligned} \|\chi_h \Psi_h\|^2 &\geq \int_{2|\mathbf{y}| \leq T} d\mathbf{y} \int_{\Upsilon \cap \{2|\mathbf{z}| \leq T\}} \left| \Phi\left(\frac{\mathbf{z}}{\sqrt{h}}\right) \right|^2 d\mathbf{z} \\ (6.30) \quad &\geq CT^{3-k} h^{k/2} \int_{\Upsilon \cap \{2|\mathbf{z}| \leq Th^{-\frac{1}{2}}\}} |\Phi(\mathbf{z})|^2 d\mathbf{z} \end{aligned}$$

$$(6.31) \quad \geq CT^{3-k} h^{k/2} \mathcal{I}(Th^{-\frac{1}{2}}) \|\Phi\|_{L^2(\Upsilon)}^2$$

where we have set for any $S \geq 0$

$$\mathcal{I}(S) := \left(\int_{\Upsilon \cap \{2|\mathbf{z}| \leq S\}} |\Phi(\mathbf{z})|^2 d\mathbf{z} \right) \left(\int_{\Upsilon} |\Phi(\mathbf{z})|^2 d\mathbf{z} \right)^{-1}.$$

The function $S \mapsto \mathcal{I}(S)$ is continuous, non-negative and non-decreasing on $[0, +\infty)$. It is moreover *increasing and positive* on $(0, \infty)$ since Φ , as a solution of an elliptic equation with polynomial coefficients and null right hand side, is analytic inside Υ . Consequently, $\mathcal{I}(Th^{-\frac{1}{2}}) = \mathcal{I}(Rh^{\delta-\frac{1}{2}})$ is uniformly bounded from below for $R \geq R_0$, $h \in (0, h_0)$, $\delta \in [0, \frac{1}{2}]$ and thus

$$\rho_h \leq \begin{cases} CT^{-2} \{\mathcal{I}(Th^{-\frac{1}{2}})\}^{-1} \leq C_0 h^{-2\delta} & \text{if } k < 3, \\ CT^{-2} e^{-2c_\psi T/\sqrt{h}} \{\mathcal{I}(Th^{-\frac{1}{2}})\}^{-1} \leq C_0 h^{-2\delta} e^{-c_0 h^{\delta-1/2}} & \text{if } k = 3, \end{cases}$$

where the constants C_0 and c_0 in the above estimation depend only on the lower bound R_0 on R , the upper bound h_0 on h , and on the model problem associated with \mathbf{x}_0 , provided $\delta \in [0, \frac{1}{2}]$. Lemma 6.9 is proved. \square

If the exponent δ is bounded from above by a number $\delta_0 < \frac{1}{2}$, we obtain the following improvement of the previous lemma.

Lemma 6.10. *Under the conditions of Lemma 6.9, let $\delta_0 < \frac{1}{2}$ be a positive number. Let $R_0 > 0$. Then there exist constants $h_0 > 0$, $C_1 > 0$ and $c_1 > 0$ depending only on R_0 , δ_0 and on the constants c_ψ , C_ψ in (6.29) such that*

$$\rho_h = \frac{\|\nabla \chi_h \Psi_h\|^2}{\|\chi_h \Psi_h\|^2} \leq \begin{cases} C_1 h^{-2\delta} & \text{if } k < 3, \\ C_1 e^{-c_1 h^{\delta-1/2}} & \text{if } k = 3, \end{cases} \quad \forall R \geq R_0, \forall h \leq h_0, \forall \delta \in [0, \delta_0].$$

Proof. We obtain an upper bound of $\|\nabla \chi_h \Psi_h\|^2$ as in the proof of Lemma 6.9. Let us now deal with the lower-bound of $\|\chi_h \Psi_h\|^2$. With $T = Rh^\delta$, we have

$$\begin{aligned} \|\chi_h \Psi_h\|^2 &\geq CT^{3-k} h^{k/2} \int_{\Upsilon \cap \{2|\mathbf{z}| \leq Th^{-\frac{1}{2}}\}} |\Phi(\mathbf{z})|^2 d\mathbf{z} \\ (6.32) \quad &\geq CT^{3-k} h^{k/2} \left(1 - C_\psi e^{-c_\psi R h^{\delta-1/2}}\right) \|\Phi\|_{L^2(\Upsilon)}^2. \end{aligned}$$

Since $0 \leq \delta \leq \delta_0 < \frac{1}{2}$, there holds $C_\psi e^{-c_\psi R h^{\delta-1/2}} < \frac{1}{2}$ for h small enough or R large enough. Thus we deduce the lemma. \square

6.3.2. *Linearization.* Note that $\psi_h^\tau \in \text{Dom}(q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}])$. We can compare the quadratic form for the magnetic potential and its affine part by using (A.7) with $\mathbf{A} = \tilde{\mathbf{A}}$, $\mathbf{A}' = \tilde{\mathbf{A}}_0$, $\mathcal{O} = \Pi_{\mathbf{x}_0}$, and $\psi = \psi_h^\tau$:

$$(6.33) \quad q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi_h^\tau) = q_h[\tilde{\mathbf{A}}_0, \Pi_{\mathbf{x}_0}](\psi_h^\tau) + 2 \operatorname{Re} \int_{\Pi_{\mathbf{x}_0}} (-ih\nabla + \tilde{\mathbf{A}}_0)\psi_h^\tau(\mathbf{x}) \cdot (\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)(\mathbf{x}) \overline{\psi_h^\tau(\mathbf{x})} \, d\mathbf{x} + \|(\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)\psi_h^\tau\|^2.$$

Combining with (6.28) we get

$$(6.34) \quad \frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi_h^\tau)}{\|\psi_h^\tau\|^2} = h\Lambda_0 + h^2\rho_h + \frac{2 \operatorname{Re} \int_{\Pi_{\mathbf{x}_0}} (-ih\nabla + \tilde{\mathbf{A}}_0)\psi_h^\tau(\mathbf{x}) \cdot (\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)(\mathbf{x}) \overline{\psi_h^\tau(\mathbf{x})} \, d\mathbf{x}}{\|\psi_h^\tau\|^2} + \frac{\|(\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)\psi_h^\tau\|^2}{\|\psi_h^\tau\|^2}.$$

By Cauchy-Schwarz inequality, we obtain easily

$$(6.35) \quad \frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi_h^\tau)}{\|\psi_h^\tau\|^2} \leq h\Lambda_0 + h^2\rho_h + 2\sqrt{h}\sqrt{\Lambda_0 + h\rho_h} a_h + a_h^2$$

where we have set

$$(6.36) \quad a_h = \frac{\|(\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)\psi_h^\tau\|}{\|\psi_h^\tau\|}.$$

We now estimate the remainder due to the linearization of $\tilde{\mathbf{A}}$.

Lemma 6.11. *Let $r_0 > 0$ be such that $\mathcal{V}_{\mathbf{x}_0} \supset \mathcal{B}(\mathbf{0}, r_0)$. For any $r \in (0, r_0]$, we have*

$$(6.37) \quad \forall \mathbf{x} \in \mathcal{B}(\mathbf{0}, r), \quad |\tilde{\mathbf{A}}(\mathbf{x}) - \tilde{\mathbf{A}}_0(\mathbf{x})| \leq \frac{1}{2} \|\tilde{\mathbf{A}}\|_{W^{2,\infty}(\mathcal{B}(0,r))} |\mathbf{x}|^2.$$

Proof. Since $\tilde{\mathbf{A}}_0$ is the affine part of $\tilde{\mathbf{A}}$, the formula is deduced from the Taylor expansion of $\tilde{\mathbf{A}}$ at $\mathbf{x} = \mathbf{0}$. \square

By construction, there exists $C(\Omega) > 0$ such that the support of ψ_h^τ is included in $\mathcal{B}(\mathbf{0}, C(\Omega)h^\delta)$. Consequently, we obtain immediately

$$(6.38) \quad a_h \leq C(\Omega) \|\tilde{\mathbf{A}}\|_{W^{2,\infty}(\operatorname{supp}(\psi_h^\tau))} h^{2\delta}.$$

Moreover using the definition of $\tilde{\mathbf{A}}$ (see (A.4)) we get

$$\begin{aligned} \|\tilde{\mathbf{A}}\|_{W^{2,\infty}(\operatorname{supp}(\psi_h^\tau))} &\leq (1 + \|\mathbb{I}_3 - J_{\mathbf{x}_0}\|_{L^\infty(\mathcal{V}_{\mathbf{x}_0})} h^\delta) \|\mathbf{A}\|_{W^{2,\infty}(\mathcal{U}_{\mathbf{x}_0})} \\ &\leq (1 + C(\Omega)h^\delta) \|\mathbf{A}\|_{W^{2,\infty}(\Omega)} \end{aligned}$$

Thus, putting this last inequality in (6.38), we deduce

$$(6.39) \quad a_h \leq C(\Omega) \|\mathbf{A}\|_{W^{2,\infty}(\Omega)} h^{2\delta}.$$

Combining (6.35), (6.39) and Lemma 6.9 we get

$$(6.40) \quad \frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi_h^\tau)}{\|\psi_h^\tau\|^2} \leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)(h^{2-2\delta} + h^{1/2+\delta} + h^{4\delta}).$$

Note that we have also used $\Lambda_0 \leq \|\mathbf{B}\|_{L^\infty(\Omega)} \leq \|\mathbf{A}\|_{W^{1,\infty}(\Omega)}$ (since $\mathbf{B} = \text{curl } \mathbf{A}$) in order to control the cross term $\sqrt{h}\sqrt{\Lambda_0 + h\rho_h a_h}$ in the right hand side of (6.35).

6.3.3. *Quasimode on Ω and estimation of the remainders.* We now define a quasimode for $q_h[\mathbf{A}, \Omega]$. Let us note that for h small enough, ψ_h^τ is compactly supported in $\mathcal{V}_{\mathbf{x}_0}$. Therefore we can define $f_h \in H^1(\Omega)$ by

$$(6.41) \quad f_h := \psi_h^\tau \circ U_{\mathbf{x}_0}.$$

Combining (6.40) with Lemma 6.5 for $r = h^\delta$ we get

$$\frac{q_h[\mathbf{A}, \Omega](f_h)}{\|f_h\|^2} \leq \left(h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)(h^{2-2\delta} + h^{1/2+2\delta} + h^{4\delta}) \right) (1 + C(\Omega)h^\delta).$$

Therefore there exists a constant $C(\Omega) > 0$ such that

$$(6.42) \quad \frac{q_h[\mathbf{A}, \Omega](f_h)}{\|f_h\|^2} \leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)(h^{2-2\delta} + h^{1/2+2\delta} + h^{4\delta} + h^{1+\delta}).$$

We optimize this upper bound by taking $\delta = \frac{3}{8}$, which provides immediately estimate (6.14). The min-max principle then yields Theorem 6.1.

6.3.4. *Improvement in case of corner concentration.* When the geometry minimizing the energy is given by a corner whose tangent problem has an eigenvalue under its essential spectrum, we get a better upper bound by improving the estimate on a_h :

Proposition 6.12. *Let $\Omega \in \overline{\mathfrak{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{A} \in W^{2,\infty}(\overline{\Omega})$ be a twice differentiable magnetic potential such that the associated magnetic field \mathbf{B} does not vanish on $\overline{\Omega}$. We assume moreover that there exists a corner $\mathbf{x}_0 \in \overline{\Omega}$ such that*

$$\mathcal{E}(\mathbf{B}, \Omega) = E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) < \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}).$$

Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that

$$\forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \leq h\mathcal{E}(\mathbf{B}, \Omega) + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) h^{3/2} |\log h|.$$

Remark 6.13. As in Corollary 6.2 we can express this upper bound in function of the Hölder norm of the magnetic field.

Proof. Since $\Pi_{\mathbf{x}_0}$ is a polyhedral corner and $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) < \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$, by Proposition 4.15 the generalized eigenfunction Ψ of $H(\mathbf{A}, \Pi_{\mathbf{x}_0})$ provided by Theorem 4.7 is an eigenfunction and has exponential decay when $|\mathbf{x}| \rightarrow +\infty$. The quasimode defined in (6.22) is sitting. Using (6.37) we get $C(\Omega) > 0$ such that

$$\forall \mathbf{x} \in \text{supp}(\psi_h), \quad |(\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)(\mathbf{x})| \leq C(\Omega) \|\tilde{\mathbf{A}}\|_{W^{2,\infty}(\text{supp}(\psi_h))} |\mathbf{x}|^2.$$

Using the change of variable $\mathbf{X} = \mathbf{x}h^{-1/2}$ and the exponential decay of Ψ we get

$$a_h = \frac{\|(\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_0)\psi_h\|}{\|\psi_h\|} \leq C(\Omega)\|\tilde{\mathbf{A}}\|_{W^{2,\infty}(\text{supp}(\psi_h))}h,$$

where a_h is set in (6.36). Now (6.35) provides, with Lemma 6.9, for any $\delta \in (0, \frac{1}{2}]$

$$\begin{aligned} \frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi_h)}{\|\psi_h\|^2} &\leq h\Lambda_0 + C(\Omega)h^{2-2\delta}e^{-ch^{\delta-\frac{1}{2}}} + C(\Omega)\|\mathbf{A}\|_{W^{2,\infty}(\Omega)}h^{3/2} + C(\Omega)\|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2h^2 \\ &\leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)(h^{2-2\delta}e^{-ch^{\delta-\frac{1}{2}}} + h^{3/2}). \end{aligned}$$

Therefore the quasimode f_h defined in (6.41) satisfies

$$\begin{aligned} \frac{q_h[\mathbf{A}, \Omega](f_h)}{\|f_h\|^2} &\leq (1 + C(\Omega)h^\delta)\{h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)(h^{2-2\delta}e^{-ch^{\delta-\frac{1}{2}}} + h^{3/2})\} \\ &\leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)\{h^{1+\delta} + h^{2-2\delta}e^{-ch^{\delta-\frac{1}{2}}} + h^{3/2}\}. \end{aligned}$$

Here $C(\Omega)$ denotes various constants independent from $h \leq h_0$ and $\delta \leq \frac{1}{2}$. We optimize this by taking $\delta = 1/2 - \varepsilon(h)$ with $\varepsilon(h)$ so that $h^{1+\delta} = h^{2-2\delta}e^{-ch^{\delta-\frac{1}{2}}}$, i.e.

$$h^{\frac{3}{2}-\varepsilon(h)} = h^{1+2\varepsilon(h)}e^{-ch^{-\varepsilon(h)}}.$$

We find

$$e^{ch^{-\varepsilon(h)}} = h^{-\frac{1}{2}+3\varepsilon(h)}, \quad \text{i.e.} \quad h^{-\varepsilon(h)} = \frac{1}{c}(-\frac{1}{2} + 3\varepsilon(h)) \log h.$$

The latter equation has one solution $\varepsilon(h)$ which tends to 0 as h tends to 0. Replacing $h^{-\varepsilon(h)}$ by the value above in $h^{\frac{3}{2}-\varepsilon(h)}$, we find that the remainder is a $O(h^{3/2}|\log h|)$ and the min-max principle provides the proposition. \square

6.4. Improvement for more regular magnetic fields. The object of this section is the proof of Theorem 6.3. In fact, our proof of the $h^{5/4}$ upper bound as done in previous sections weakly uses the exponential decay of generalized eigenfunctions in some directions. It would also work with purely oscillating generalized eigenfunctions.

Now the proof of the $h^{4/3}$ upper bound makes a more extensive use of fine properties of the model problems: First, the decay properties of admissible generalized eigenvectors and their stability upon perturbation, and second, the Lipschitz regularity of the ground energy depending on the magnetic field, cf. Lemma 4.19.

We recall that $\mathbf{x}_0 \in \overline{\Omega}$ is a point such that $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) = \mathcal{E}(\mathbf{B}, \Omega) =: \Lambda_0$. We apply Theorem 4.7 (and Remark 4.8) with $(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$: we denote by \mathbb{X} the corresponding singular chain which satisfies

$$\mathcal{E}(\mathbf{B}, \Omega) = E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) = E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbb{X}}) < \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbb{X}}).$$

We now split our analysis between three geometric configurations depending on the number of variables k in which the generalized eigenfunction has exponential decay (see Section 4.8):

(G1) $\Pi_{\mathbb{X}}$ is a half-space and $\mathbf{B}_{\mathbf{x}_0}$ is tangent to the boundary.

(G2) We are in one of the following situations:

- Either $\Pi_{\mathbb{X}}$ is a wedge,
- or $\Pi_{\mathbb{X}}$ is a half-space and $\mathbf{B}_{\mathbf{x}_0}$ is neither tangent nor normal to the boundary.
- or $\Pi_{\mathbb{X}} = \mathbb{R}^3$.

(G3) $\Pi_{\mathbb{X}}$ is a polyhedral cone of dimension 3 and coincides with $\Pi_{\mathbf{x}_0}$.

Let us now deal with each situation. The arguments are specific to each case.

• *Assume that we are in situation (G3).* This means that \mathbf{x}_0 is a corner and that $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) < \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$. In that case we can rely on Proposition 6.12 in which we have already proved a better upper bound for $\lambda_h(\mathbf{B}, \Omega)$, even with a weaker regularity assumption on the magnetic field.

• *Assume that we are in situation (G2).* The generalized eigenfunction Ψ associated with $H(\mathbf{A}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$ has two directions of decay, z_1 and z_2 , leaving one direction y with a purely oscillating character. In this case, we are going to improve the linearization error: Until now we have used that $\tilde{\mathbf{A}}(\mathbf{x}) - \tilde{\mathbf{A}}_0(\mathbf{x})$ is a $O(|\mathbf{x}|^2)$. Here, by a suitable phase shift (which corresponds to a change of gauge), we can eliminate from this error the term in $O(|y|^2)$, replacing it by a $O(|y|^3)$. The other terms containing at least one power of $|\mathbf{z}|$, we can take advantage of the decay of Ψ . The sitting modes will be constructed following exactly this strategy, whereas concerning sliding modes, we have to linearize the potential at the point $\mathbf{p} := h^\delta \boldsymbol{\tau}$, instead of $\mathbf{0}$ as previously. Let us develop details now.

We recall that we have introduced a vector $\boldsymbol{\tau}$ in Definition 6.6 depending on $\Pi_{\mathbf{x}_0}$ and $\Pi_{\mathbb{X}}$. Mimicking Definition 6.8 for quasimodes $\psi_h^\boldsymbol{\tau}$, we are going to construct new quasimodes with two adaptations: first, we linearize the potential at the point $\mathbf{p} := h^\delta \boldsymbol{\tau}$, second, we shift the phase to optimize remainders.

In a first step, we consider $\mathbf{p} \in \overline{\Pi_{\mathbf{x}_0}} \cap \mathcal{V}_{\mathbf{x}_0}$ we denote by $\tilde{\mathbf{A}}_{\mathbf{p}}$ the affine part of $\tilde{\mathbf{A}}$ at \mathbf{p} :

$$\tilde{\mathbf{A}}_{\mathbf{p}}(\mathbf{x}) = \tilde{\mathbf{A}}(\mathbf{p}) + \nabla \tilde{\mathbf{A}}(\mathbf{p})(\mathbf{x} - \mathbf{p}).$$

We have $\text{curl } \tilde{\mathbf{A}}_{\mathbf{p}} = \tilde{\mathbf{B}}_{\mathbf{p}}$ where the constant $\tilde{\mathbf{B}}_{\mathbf{p}}$ is the magnetic field $\tilde{\mathbf{B}}$ frozen at \mathbf{p} .

We have $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbb{X}}) < \mathcal{E}^*(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbb{X}})$. Due to Lemma 4.19, we have

$$(6.43) \quad \exists \varepsilon > 0, \quad \forall \mathbf{p} \in \mathcal{B}(0, \varepsilon) \cap \overline{\Pi_{\mathbf{x}_0}}, \quad E(\tilde{\mathbf{B}}_{\mathbf{p}}, \Pi_{\mathbb{X}}) < \mathcal{E}^*(\tilde{\mathbf{B}}_{\mathbf{p}}, \Pi_{\mathbb{X}}),$$

and $(\tilde{\mathbf{B}}_{\mathbf{p}}, \Pi_{\mathbb{X}})$ is still in situation (G2). There exists an admissible generalized eigenfunction $\Psi^{\mathbf{p}}$ for the operator $H(\tilde{\mathbf{A}}_{\mathbf{p}}, \Pi_{\mathbb{X}})$ associated with $E(\tilde{\mathbf{B}}_{\mathbf{p}}, \Pi_{\mathbb{X}})$ (denoted by $\Lambda_{\mathbf{p}}$ for shortness):

$$(6.44) \quad \Psi^{\mathbf{p}} \circ \underline{U}_{\mathbf{p}}^{-1}(\mathbf{y}, \mathbf{z}) = e^{i\varphi^{\mathbf{p}}(\mathbf{y}, \mathbf{z})} \Phi^{\mathbf{p}}(\mathbf{z}) \quad \text{for} \quad (\mathbf{y}, \mathbf{z}) \in \mathbb{R} \times \Upsilon,$$

where the rotation⁵ \underline{U}_p maps $\Pi_{\mathbb{X}}$ onto $\mathbb{R} \times \Upsilon$. Here Υ is a cone in 2 dimensions (namely a sector, a half-space or \mathbb{R}^2). We have

$$(6.45) \quad \begin{cases} (-i\nabla + \tilde{\mathbf{A}}_p)^2 \Psi^p = \Lambda_p \Psi^p & \text{in } \Pi_{\mathbb{X}}, \\ (-i\partial_n + \mathbf{n} \cdot \tilde{\mathbf{A}}_p) \Psi^p = 0 & \text{on } \partial\Pi_{\mathbb{X}}. \end{cases}$$

An important point is that, choosing $\varepsilon > 0$ small enough, we may assume that, in virtue of Lemma 4.19, the functions Φ^p are uniformly exponentially decreasing

$$(6.46) \quad \exists c > 0, C > 0, \quad \forall \mathbf{p} \in \mathcal{B}(\mathbf{0}, \varepsilon), \quad \|\Phi^p e^{c|\mathbf{z}|}\|_{L^2(\Upsilon)} \leq C \|\Phi^p\|_{L^2(\Upsilon)}.$$

We set $\Psi_h^p(\mathbf{x}) := \Psi^p(\frac{\mathbf{x}}{h})$.

Now we take $\mathbf{p} = \boldsymbol{\tau} h^\delta$ and the function $\psi_h^{\boldsymbol{\tau}, p}$ is defined in a similar way as in (6.27)

$$(6.47) \quad \psi_h^{\boldsymbol{\tau}, p}(\mathbf{x}) = e^{i\langle \tilde{\mathbf{A}}_p(\mathbf{0}) - \tilde{\mathbf{A}}_p(\boldsymbol{\tau} h^\delta), \mathbf{x}/h \rangle} \underline{\chi}_R \left(\frac{|\mathbf{x} - \boldsymbol{\tau} h^\delta|}{h^\delta} \right) \Psi^p \left(\frac{\mathbf{x} - \boldsymbol{\tau} h^\delta}{h^{1/2}} \right), \quad \mathbf{x} \in \Pi_{\mathbf{x}_0}.$$

Note that, compared with (6.27), the function $\psi_h^{\boldsymbol{\tau}, p}$ is adapted to the potential $\tilde{\mathbf{A}}_p$ instead of $\tilde{\mathbf{A}}_0$:

$$(6.48) \quad q_h[\tilde{\mathbf{A}}_p, \Pi_{\mathbf{x}_0}](\psi_h^{\boldsymbol{\tau}, p}) = q_h[\tilde{\mathbf{A}}_p, \Pi_{\mathbb{X}}](\psi_h^{\boldsymbol{\tau}, p}) = q_h[\tilde{\mathbf{A}}_p, \Pi_{\mathbb{X}}](\psi_h^{0, p}),$$

with

$$\psi_h^{0, p} = \underline{\chi}_R \left(\frac{\mathbf{x}}{h^\delta} \right) \Psi^p \left(\frac{\mathbf{x}}{h^{1/2}} \right) = \underline{\chi}_R \left(\frac{\mathbf{x}}{h^\delta} \right) \Psi_h^p(\mathbf{x}), \quad \mathbf{x} \in \Pi_{\mathbb{X}}.$$

We are going to apply Lemma A.3 with the variables $(\mathbf{y}, \mathbf{z}) = \mathbf{x}^{\natural}$ adapted to the structure of the generalized eigenvector Ψ^p (6.44). We recall that \underline{U}_p is the rotation (J its associated matrix) such that $\underline{U}_p(\mathbf{x} - \mathbf{p}) = J^T(\mathbf{x} - \mathbf{p}) = \mathbf{x}^{\natural}$. The magnetic potential \mathbf{A}^{\natural} associated with $\tilde{\mathbf{A}}$ in variables \mathbf{x}^{\natural} is defined as, cf. Remark 4.5

$$\mathbf{A}^{\natural}(\mathbf{x}^{\natural}) = J^T(\tilde{\mathbf{A}}(\mathbf{x})).$$

Note that there holds the corresponding identity for the affine parts

$$\mathbf{A}_0^{\natural}(\mathbf{x}^{\natural}) = J^T(\tilde{\mathbf{A}}_p(\mathbf{x})).$$

Lemma A.3 in variables $(u_1, u_2, u_3) = (y, z_1, z_2)$ with $\ell = 1$ then gives us a function F^{\natural} such that $\partial_y^2(\mathbf{A}^{\natural} - \nabla_{\natural} F^{\natural})(\mathbf{0}) = 0$ leading to the estimates

$$(6.49) \quad |(\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural} - \nabla_{\natural} F^{\natural})(\mathbf{x}^{\natural})| \leq C(\mathcal{V}_{\mathbf{x}_0}) \|\tilde{\mathbf{A}}\|_{W^{3,\infty}(\mathcal{V}_{\mathbf{x}_0})} (|y|^3 + |y||\mathbf{z}| + |\mathbf{z}|^2).$$

Now we set

$$F_p(\mathbf{x}) = F^{\natural}(\mathbf{x}^{\natural}) = F^{\natural}(\underline{U}_p(\mathbf{x} - \mathbf{p})).$$

Note that we have

$$(6.50) \quad (\tilde{\mathbf{A}} - \tilde{\mathbf{A}}_p - \nabla F_p)(\mathbf{x}) = J(\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural} - \nabla_{\natural} F^{\natural}) \circ \underline{U}_p(\mathbf{x} - \mathbf{p}).$$

⁵The rotation \underline{U}_p may depend on \mathbf{p} but that does not hamper our analysis.

We then define our new quasimode:

$$(6.51) \quad \widehat{\psi}_h^\tau(\mathbf{x}) := e^{iF_p(\mathbf{x})/h} \psi_h^{\tau,p}(\mathbf{x}) \quad \text{with} \quad \mathbf{p} = h^\delta \boldsymbol{\tau}.$$

We have obviously $\|\widehat{\psi}_h^\tau\| = \|\psi_h^{\tau,p}\|$. Owing to Lemma A.1 we have

$$q_h[\widetilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\widehat{\psi}_h^\tau) = q_h[\widetilde{\mathbf{A}} - \nabla F_p, \Pi_{\mathbf{x}_0}](\psi_h^{\tau,p}),$$

and combining with formula (A.7) we get:

$$(6.52) \quad \begin{aligned} q_h[\widetilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\widehat{\psi}_h^\tau) &= q_h[\widetilde{\mathbf{A}} - \nabla F_p, \Pi_{\mathbf{x}_0}](\psi_h^{\tau,p}) \\ &= q_h[\widetilde{\mathbf{A}}_p, \Pi_{\mathbb{X}}](\psi_h^{\tau,p}) + \|(\widetilde{\mathbf{A}} - \widetilde{\mathbf{A}}_p - \nabla F_p)\psi_h^{\tau,p}\|^2 \\ &\quad + 2 \operatorname{Re} \int_{\Pi_{\mathbb{X}}} (-ih\nabla + \widetilde{\mathbf{A}}_p)\psi_h^{\tau,p}(\mathbf{x}) \cdot (\widetilde{\mathbf{A}} - \widetilde{\mathbf{A}}_p - \nabla F_p)(\mathbf{x}) \overline{\psi_h^{\tau,p}(\mathbf{x})} \, d\mathbf{x}. \end{aligned}$$

By analogy with (6.36), we define

$$\widehat{a}_h = \frac{\|(\widetilde{\mathbf{A}} - \widetilde{\mathbf{A}}_p - \nabla F_p)\psi_h^{\tau,p}\|}{\|\psi_h^{\tau,p}\|}.$$

Recall that we have denoted Λ_p for $E(\widetilde{\mathbf{B}}_p, \Pi_{\mathbb{X}})$. Using Lemma A.8 we get like for (6.28)

$$(6.53) \quad \frac{q_h[\widetilde{\mathbf{A}}_p, \Pi_{\mathbf{x}_0}](\psi_h^{\tau,p})}{\|\psi_h^{\tau,p}\|^2} = \frac{q_h[\widetilde{\mathbf{A}}_p, \Pi_{\mathbb{X}}](\chi_h \Psi_h^p)}{\|\chi_h \Psi_h^p\|^2} = h\Lambda_p + h^2\rho_h \quad \text{with} \quad \rho_h = \frac{\|\nabla \chi_h\| \|\Psi_h^p\|^2}{\|\chi_h \Psi_h^p\|^2}.$$

Here ρ_h satisfies the estimates as given in Lemma 6.10 because of the uniformly exponential decay (6.46).

Combining (6.52) and (6.53), we obtain

$$(6.54) \quad \frac{q_h[\widetilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\widehat{\psi}_h^\tau)}{\|\widehat{\psi}_h^\tau\|^2} \leq h\Lambda_p + h^2\rho_h + 2\sqrt{h}\sqrt{\Lambda_p + h\rho_h} \widehat{a}_h + \widehat{a}_h^2$$

Compared with (6.35), we have \widehat{a}_h instead of a_h . The following Lemma provides an improvement when compared to (6.38), due to estimates (6.49) which replace (6.37).

Lemma 6.14. *There exist constants $C(\Omega) > 0$ and $h_0 > 0$ such that for all $h \in (0, h_0)$*

$$(6.55) \quad \widehat{a}_h = \frac{\|(\widetilde{\mathbf{A}} - \widetilde{\mathbf{A}}_p - \nabla F_p)\psi_h^{\tau,p}\|}{\|\psi_h^{\tau,p}\|} \leq C(\Omega) \|\widetilde{\mathbf{A}}\|_{W^{3,\infty}(\mathcal{V}_{\mathbf{x}_0})} h^{\min(1, \frac{1}{2} + \delta, 3\delta)}.$$

Proof. Let us write the admissible generalized eigenvector Ψ^p (6.44) directly in coordinates $\mathbf{x}^\natural = (y, \mathbf{z})$:

$$\Psi^{\natural,p}(\mathbf{x}^\natural) = e^{i\varphi_p(y,\mathbf{z})} \Phi^p(\mathbf{z}).$$

Let $\psi_h^{\natural,p}(\mathbf{x}^\natural) := \psi_h^{\tau,p}(\mathbf{x})$ with $\mathbf{x}^\natural = \underline{\mathbf{u}}_p(\mathbf{x} - \mathbf{p})$. From (6.50) we deduce

$$(\widetilde{\mathbf{A}} - \widetilde{\mathbf{A}}_p - \nabla F_p)\psi_h^{\tau,p}(\mathbf{x}) = J(\mathbf{A}^\natural - \mathbf{A}_0^\natural - \nabla_\natural F^\natural)\psi_h^{\natural,p}(\mathbf{x}^\natural)$$

and combining with (6.49) we obtain

$$|(\widetilde{\mathbf{A}} - \widetilde{\mathbf{A}}_p - \nabla F_p)\psi_h^{\tau,p}(\mathbf{x})| \leq C(\mathcal{V}_{\mathbf{x}_0}) \|\widetilde{\mathbf{A}}\|_{W^{3,\infty}(\mathcal{V}_{\mathbf{x}_0})} (|y|^3 + |y||\mathbf{z}| + |\mathbf{z}|^2) |\psi_h^{\natural,p}(\mathbf{x}^\natural)|.$$

We note that

$$|\psi_h^{\mathfrak{h},\mathfrak{p}}(\mathbf{x}^{\mathfrak{h}})| = \underline{\chi}_R \left(\frac{|\mathbf{x}^{\mathfrak{h}}|}{h^\delta} \right) \left| \Psi^{\mathfrak{h},\mathfrak{p}} \left(\frac{\mathbf{x}^{\mathfrak{h}}}{h^{1/2}} \right) \right| = \underline{\chi}_R \left(\frac{|\mathbf{x}^{\mathfrak{h}}|}{h^\delta} \right) \left| \Phi^{\mathfrak{p}} \left(\frac{\mathbf{z}}{h^{1/2}} \right) \right|.$$

Using the uniform exponential decay (6.46) and the changes of variables $\mathbf{Z} = \mathbf{z}h^{-1/2}$ and $Y = yh^{-\delta}$, we find the bounds

$$\begin{aligned} \left\| |y|^3 \underline{\chi}_R \left(\frac{|\mathbf{x}^{\mathfrak{h}}|}{h^\delta} \right) \Phi^{\mathfrak{p}} \left(\frac{\mathbf{z}}{h^{1/2}} \right) \right\| &\leq h^{3\delta} \|\psi_h^{\mathfrak{h},\mathfrak{p}}\| \\ \left\| |y| |\mathbf{z}| \underline{\chi}_R \left(\frac{|\mathbf{x}^{\mathfrak{h}}|}{h^\delta} \right) \Phi^{\mathfrak{p}} \left(\frac{\mathbf{z}}{h^{1/2}} \right) \right\| &\leq h^{\delta+1/2} \|\psi_h^{\mathfrak{h},\mathfrak{p}}\| \\ \left\| |\mathbf{z}|^2 \underline{\chi}_R \left(\frac{|\mathbf{x}^{\mathfrak{h}}|}{h^\delta} \right) \Phi^{\mathfrak{p}} \left(\frac{\mathbf{z}}{h^{1/2}} \right) \right\| &\leq h \|\psi_h^{\mathfrak{h},\mathfrak{p}}\|. \end{aligned}$$

Summing up the latter three estimates leads to the lemma. \square

We now take the same arguments as in Section 6.3.2 but instead of (6.39) we use Lemma 6.14 to estimate \hat{a}_h and (6.54) becomes

$$\frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\hat{\psi}_h^\tau)}{\|\hat{\psi}_h^\tau\|^2} \leq h\Lambda_{\mathfrak{p}} + C(\Omega)(1 + \|\mathbf{A}\|_{W^{3,\infty}(\Omega)}^2)(h^{3/2} + h^{2-2\delta} + h^{1+\delta} + h^{3\delta+1/2} + h^{6\delta}).$$

Now we use Lemma 4.19 to obtain the estimate $|\Lambda_{\mathfrak{p}} - \Lambda_0| \leq C|\mathfrak{p}| \leq Ch^\delta$ (as previously Λ_0 denotes $E(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0})$). Hence

$$(6.56) \quad \frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\hat{\psi}_h^\tau)}{\|\hat{\psi}_h^\tau\|^2} \leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{3,\infty}(\Omega)}^2)(h^{3/2} + h^{2-2\delta} + h^{1+\delta} + h^{3\delta+1/2} + h^{6\delta}).$$

Let \hat{f}_h be the quasimode defined in the same way as in (6.41). The same arguments as in Section 6.3.3 and Lemma 4.19 combined with (6.56) yields

$$(6.57) \quad \frac{q_h[\mathbf{A}, \Omega](\hat{f}_h)}{\|\hat{f}_h\|^2} \leq h\Lambda_0 + C(\Omega)(1 + \|\mathbf{A}\|_{W^{3,\infty}(\Omega)}^2)(h^{2-2\delta} + h^{3\delta+1/2} + h^{6\delta} + h^{1+\delta}).$$

We optimize this upper bound by taking $\delta = \frac{1}{3}$. The min-max principle provides Theorem 6.3 in the case (G2).

- *Assume we are in situation (G1).* In situation (G1) the generalized eigenfunction has exponential decay in one variable z . The upper bound (6.35) obtained by a Cauchy-Schwarz inequality is too rough and we will deal with the previous identity (6.34). A Feynman-Hellmann formula will simplify the cross term in (6.34) and the exponential decay in one variable will provide the desired result.

In situation (G1) $\Pi_{\mathbb{X}}$ is a half-plane and $\mathbf{B}_{\mathbf{x}_0}$ is tangent to its boundary. Denote by $(\mathbf{y}, \mathbf{z}) = (y_1, y_2, z) \in \mathbb{R}^2 \times \mathbb{R}_+$ a system of coordinates of $\Pi_{\mathbb{X}}$ such that $\mathbf{B}_{\mathbf{x}_0}$ is tangent to the y_2 -axis. In these coordinates, the magnetic field $\mathbf{B}_{\mathbf{x}_0}$ writes $(0, b, 0)$.

In the rest of this proof, we will assume without restriction that $b = 1$. Indeed, once quasimodes are constructed for $b = 1$, Lemmas A.5 and A.6 allow to convert them into quasimodes for any b .

Let $\underline{U} : \Pi_{\mathbb{X}} \mapsto \mathbb{R}^2 \times \mathbb{R}_+$ be the rotation associated to the coordinates $(\mathbf{y}, \mathbf{z}) = \mathbf{x}^{\natural}$ and J be its associated matrix. The magnetic potential and its linear part becomes (cf. Remark 4.5)

$$(6.58) \quad \mathbf{A}^{\natural}(\mathbf{x}^{\natural}) = J^{\top}(\tilde{\mathbf{A}}(\mathbf{x})), \quad \mathbf{A}_0^{\natural}(\mathbf{x}^{\natural}) = J^{\top}(\tilde{\mathbf{A}}_0(\mathbf{x})).$$

Let us define the canonical reference potential

$$\underline{\mathbf{A}}(\mathbf{x}^{\natural}) = (z, 0, 0).$$

such that $\text{curl } \underline{\mathbf{A}} = (0, 1, 0)$. Since \mathbf{A}_0^{\natural} and $\underline{\mathbf{A}}$ are both affine with $\text{curl } \mathbf{A}_0^{\natural} = \text{curl } \underline{\mathbf{A}}$, there exists a polynomial function of degree two F^{\natural} such that

$$(6.59) \quad \mathbf{A}_0^{\natural} - \nabla F^{\natural} = \underline{\mathbf{A}}.$$

We know (see Section 4.8) that

$$\underline{\Psi}_h(\mathbf{y}, \mathbf{z}) := e^{-i\sqrt{\Theta_0}y_1/\sqrt{h}} \Phi\left(\frac{z}{\sqrt{h}}\right)$$

is a generalized eigenvector of $H_h(\underline{\mathbf{A}}, \mathbb{R}_+^3)$ where Φ is a normalized eigenvector associated with the first eigenvalue of the de Gennes operator $-\partial_z^2 + (z - \sqrt{\Theta_0})^2$ on \mathbb{R}_+ .

Therefore, setting

$$\Psi_h^{\natural}(\mathbf{y}, \mathbf{z}) := e^{-iF^{\natural}(\mathbf{x}^{\natural})/h} \underline{\Psi}_h(\mathbf{y}, \mathbf{z})$$

we obtain an admissible generalized eigenvector for the operator $H(\mathbf{A}_0^{\natural}, \mathbb{R}_+^3)$.

We define ψ_h^{\natural} in the same way as in (6.21) but for convenience we take a cut-off function in tensor product form (here for simplicity we denote $\underline{\chi}_R$ by χ)

$$(6.60) \quad \psi_h^{\natural}(\mathbf{x}^{\natural}) := \chi\left(\frac{|\mathbf{y}|}{h^\delta}\right) \chi\left(\frac{z}{h^\delta}\right) \Psi_h^{\natural}(\mathbf{x}^{\natural}).$$

Let $\boldsymbol{\tau}$ be the vector introduced in Definition 6.6 and $\boldsymbol{\tau}^{\natural} = \underline{U}\boldsymbol{\tau}$. We note that $\boldsymbol{\tau}^{\natural}$ has no component in the z direction, because $\boldsymbol{\tau}$ is tangent to the boundary of the half-space $\Pi_{\mathbb{X}}$, see Remark 6.7. We define $\mathbf{p}^{\natural} = \boldsymbol{\tau}^{\natural}h^\delta$, which we write $\mathbf{p}^{\natural} = (\mathbf{p}_y, 0) = (p_1, p_2, 0)$ in coordinates \mathbf{x}^{\natural} .

We introduce our quasimode for $H_h(\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0})$:

$$(6.61) \quad \psi_h^{\boldsymbol{\tau}}(\mathbf{x}) = \psi_h^{\natural, \boldsymbol{\tau}}(\mathbf{x}^{\natural}) \quad \text{with} \quad \psi_h^{\natural, \boldsymbol{\tau}}(\mathbf{x}^{\natural}) = e^{i\langle \mathbf{A}_0^{\natural}(0) - \mathbf{A}_0^{\natural}(\mathbf{p}^{\natural}), \mathbf{x}/h \rangle} \psi_h^{\natural}(\mathbf{x}^{\natural} - \mathbf{p}^{\natural})$$

We set

$$\underline{\psi}_h^{\boldsymbol{\tau}}(\mathbf{x}^{\natural}) = \underline{\psi}_h(\mathbf{x}^{\natural} - \mathbf{p}^{\natural}) \quad \text{with} \quad \underline{\psi}_h(\mathbf{x}^{\natural}) := \chi\left(\frac{|\mathbf{y}|}{h^\delta}\right) \chi\left(\frac{z}{h^\delta}\right) \underline{\Psi}_h(\mathbf{x}^{\natural}).$$

Since $\underline{\mathbf{A}}$ does not depend on \mathbf{y} and \mathbf{p}^{\natural} has no component in z , we note that the above formula coincides with

$$\underline{\psi}_h^\tau(\mathbf{x}^{\natural}) = e^{i(\underline{\mathbf{A}}(\mathbf{0}) - \underline{\mathbf{A}}(\mathbf{p}^{\natural}), \mathbf{x}/h)} \underline{\psi}_h(\mathbf{x}^{\natural} - \mathbf{p}^{\natural})$$

We have $\psi_h^{\natural, \tau}(\mathbf{x}^{\natural}) = e^{-iF^{\natural}(\mathbf{x}^{\natural} - \mathbf{p}^{\natural})/h} \underline{\psi}_h^\tau(\mathbf{x}^{\natural})$. Hence

$$(6.62) \quad \begin{aligned} q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\psi_h^\tau) &= q_h[\mathbf{A}^{\natural}, \mathbb{R}_+^3](\psi_h^{\natural, \tau}) \\ &= q_h[\mathbf{A}^{\natural} - \nabla F^{\natural}, \mathbb{R}_+^3](\underline{\psi}_h^\tau). \end{aligned}$$

Now we apply (A.7) with $\mathbf{A} = \mathbf{A}^{\natural} - \nabla F^{\natural}$ and $\mathbf{A}' = \underline{\mathbf{A}}$. Using (6.59) we find $\mathbf{A} - \mathbf{A}' = \mathbf{A}^{\natural} - \mathbf{A}_0^{\natural}$, and thus

$$(6.63) \quad q_h[\mathbf{A}^{\natural} - \nabla F^{\natural}, \mathbb{R}_+^3](\underline{\psi}_h^\tau) = q_h[\underline{\mathbf{A}}, \mathbb{R}_+^3](\underline{\psi}_h^\tau)$$

$$(6.64) \quad + 2 \operatorname{Re} \int_{\mathbb{R}_+^3} (-ih\nabla + \underline{\mathbf{A}})\underline{\psi}_h^\tau(\mathbf{x}^{\natural}) \cdot (\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural})(\mathbf{x}^{\natural}) \overline{\underline{\psi}_h^\tau(\mathbf{x}^{\natural})} \, d\mathbf{x}^{\natural}$$

$$(6.65) \quad + \|(\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural})\underline{\psi}_h^\tau\|^2.$$

We bound from above the term (6.65) like in (6.38):

$$(6.66) \quad \|(\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural})\underline{\psi}_h^\tau\|^2 \leq C(\Omega) \|\mathbf{A}^{\natural}\|_{W^{2,\infty}(\operatorname{supp}(\underline{\psi}_h^\tau))}^2 h^{4\delta} \|\underline{\psi}_h^\tau\|^2.$$

Let us now deal with the term (6.64). We have to calculate $(-ih\nabla + \underline{\mathbf{A}})\underline{\psi}_h^\tau(\mathbf{x}^{\natural})$. By definition

$$\underline{\psi}_h^\tau(\mathbf{x}^{\natural}) = \chi\left(\frac{|\mathbf{y} - \mathbf{p}_y|}{h^\delta}\right) \chi\left(\frac{z}{h^\delta}\right) e^{-i\sqrt{\Theta_0}(y_1 - p_1)/\sqrt{h}} \Phi\left(\frac{z}{\sqrt{h}}\right).$$

Whence

$$\begin{aligned} (-ih\nabla + \underline{\mathbf{A}})\underline{\psi}_h^\tau &= e^{-i\sqrt{\Theta_0}(y_1 - p_1)/\sqrt{h}} \times \\ &\left\{ \chi\left(\frac{|\mathbf{y} - \mathbf{p}_y|}{h^\delta}\right) \chi\left(\frac{z}{h^\delta}\right) \begin{bmatrix} (z - \sqrt{h\Theta_0}) \Phi\left(\frac{z}{\sqrt{h}}\right) \\ 0 \\ -i\sqrt{h} \Phi'\left(\frac{z}{\sqrt{h}}\right) \end{bmatrix} - ih^{1-\delta} \begin{bmatrix} \frac{y_1 - p_1}{|\mathbf{y} - \mathbf{p}_y|} \chi'\left(\frac{|\mathbf{y} - \mathbf{p}_y|}{h^\delta}\right) \chi\left(\frac{z}{h^\delta}\right) \\ \frac{y_2 - p_2}{|\mathbf{y} - \mathbf{p}_y|} \chi'\left(\frac{|\mathbf{y} - \mathbf{p}_y|}{h^\delta}\right) \chi\left(\frac{z}{h^\delta}\right) \\ \chi\left(\frac{|\mathbf{y}|}{h^\delta}\right) \chi'\left(\frac{z}{h^\delta}\right) \end{bmatrix} \Phi\left(\frac{z}{\sqrt{h}}\right) \right\}. \end{aligned}$$

Since Φ and χ are real valued functions, the term (6.64) reduces to a single term:

$$(6.67) \quad \begin{aligned} \operatorname{Re} \int_{\mathbb{R}_+^3} (-ih\nabla + \underline{\mathbf{A}})\underline{\psi}_h^\tau(\mathbf{x}^{\natural}) \cdot (\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural})(\mathbf{x}^{\natural}) \overline{\underline{\psi}_h^\tau(\mathbf{x}^{\natural})} \, d\mathbf{x}^{\natural} \\ = \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) |\underline{\psi}_h^\tau(\mathbf{x}^{\natural})|^2 A_2^{(\operatorname{rem},2)}(\mathbf{x}^{\natural}) \, d\mathbf{x}^{\natural} \\ = \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) \left| \Phi\left(\frac{z}{\sqrt{h}}\right) \right|^2 \left| \chi\left(\frac{|\mathbf{y} - \mathbf{p}_y|}{h^\delta}\right) \right|^2 \left| \chi\left(\frac{z}{h^\delta}\right) \right|^2 A_2^{(\operatorname{rem},2)}(\mathbf{x}^{\natural}) \, d\mathbf{x}^{\natural}, \end{aligned}$$

where $A_2^{(\operatorname{rem},2)}$ denotes the second component of $\mathbf{A}^{\natural} - \mathbf{A}_0^{\natural}$. We write

$$A_2^{(\operatorname{rem},2)}(\mathbf{x}^{\natural}) = P_2(\mathbf{y}) + R_2(\mathbf{x}^{\natural}) + A_2^{(\operatorname{rem},3)}(\mathbf{x}^{\natural}),$$

where $A_2^{(\text{rem},3)}$ is the Taylor remainder of degree 3 of the second component of \mathbf{A}^{\natural} at $\mathbf{0}$, whereas $P_2(\mathbf{y}) + R_2(\mathbf{x}^{\natural})$ is a representation of its quadratic part in the form

$$P_2(\mathbf{y}) = a_1 y_1^2 + a_2 y_2^2 + a_3 y_1 y_2 \quad \text{and} \quad R_2(\mathbf{x}^{\natural}) = b_1 z^2 + b_2 z y_1 + b_3 z y_2.$$

As in (A.2) there holds

$$\|A_2^{(\text{rem},3)}\|_{L^\infty(\text{supp}(\underline{\psi}_h))} \leq C \|\tilde{\mathbf{A}}\|_{W^{3,\infty}(\text{supp}(\underline{\psi}_h^\tau))} h^{3\delta},$$

leading to, with the help of the variable change $Z = z/\sqrt{h}$ and the exponential decay of Φ :

$$(6.68) \quad \left| \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) |\underline{\psi}_h^\tau(\mathbf{x}^{\natural})|^2 A_2^{(\text{rem},3)}(\mathbf{x}^{\natural}) d\mathbf{x}^{\natural} \right| \leq C h^{\frac{1}{2}+3\delta} \|\underline{\psi}_h^\tau\|^2.$$

Likewise, combining the exponential decay of Φ , the change of variable $Z = z/\sqrt{h}$ and the localization of the support in balls of size Ch^δ , we deduce

$$(6.69) \quad \left| \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) |\underline{\psi}_h^\tau(\mathbf{x}^{\natural})|^2 R_2(\mathbf{x}^{\natural}) d\mathbf{x}^{\natural} \right| \leq C h^{\min(\frac{3}{2}, 1+\delta)} \|\underline{\psi}_h^\tau\|^2.$$

Let us now deal with the term involving $\mathbf{y} \mapsto P_2(\mathbf{y})$. Due to a Feynman-Hellmann formula applied to the de Gennes operator $\mathcal{H}(\tau)$ at $\tau = -\sqrt{\Theta_0}$ (cf. [16, Lemma A.1]) we find by the scaling $z \mapsto z/\sqrt{h}$ the identity

$$\int_{\mathbb{R}_+} (z - \sqrt{h\Theta_0}) \left| \Phi\left(\frac{z}{\sqrt{h}}\right) \right|^2 dz = 0.$$

Thus we can write

$$\begin{aligned} & \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) |\underline{\psi}_h^\tau(\mathbf{x}^{\natural})|^2 P_2(\mathbf{y}) d\mathbf{x}^{\natural} \\ &= \int_{\mathbb{R}^2} P_2(\mathbf{y}) \left| \chi\left(\frac{|\mathbf{y}-\mathbf{p}_y|}{h^\delta}\right) \right|^2 d\mathbf{y} \int_{z \in \mathbb{R}_+} (z - \sqrt{h\Theta_0}) \left| \Phi\left(\frac{z}{\sqrt{h}}\right) \right|^2 \chi\left(\frac{z}{h^\delta}\right)^2 dz \\ &= \int_{\mathbb{R}^2} P_2(\mathbf{y}) \left| \chi\left(\frac{|\mathbf{y}-\mathbf{p}_y|}{h^\delta}\right) \right|^2 d\mathbf{y} \int_{z \in \mathbb{R}_+} (z - \sqrt{h\Theta_0}) \left| \Phi\left(\frac{z}{\sqrt{h}}\right) \right|^2 \left(\chi\left(\frac{z}{h^\delta}\right)^2 - 1 \right) dz \end{aligned}$$

The support of the integral in z is contained in $z \geq Rh^\delta$ with $\delta < \frac{1}{2}$. Therefore, using once more the changes of variables $\mathbf{Y} = (\mathbf{y} - \mathbf{p}_y)/h^\delta$ and $Z = z/\sqrt{h}$, we find:

$$\left| \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) |\underline{\psi}_h^\tau(\mathbf{x}^{\natural})|^2 P_2(\mathbf{y}) d\mathbf{x}^{\natural} \right| \leq C \|\mathbf{A}^{\natural}\|_{W^{2,\infty}(\text{supp}(\underline{\psi}_h^{\natural,\tau}))} h^{4\delta + \frac{1}{2}} e^{-ch^{\delta-1/2}}.$$

Since $\|\underline{\psi}_h^\tau\|^2 \geq Ch^{2\delta + \frac{1}{2}}$ (see (6.32)), this leads to:

$$(6.70) \quad \left| \int_{\mathbb{R}_+^3} (z - \sqrt{h\Theta_0}) |\underline{\psi}_h^\tau(\mathbf{x}^{\natural})|^2 P_2(\mathbf{y}) d\mathbf{x}^{\natural} \right| \leq C \|\mathbf{A}^{\natural}\|_{W^{2,\infty}(\text{supp}(\underline{\psi}_h^{\natural,\tau}))} e^{-ch^{\delta-1/2}} \|\underline{\psi}_h^\tau\|^2.$$

Collecting (6.68), (6.69), and (6.70) in (6.64), we find the upper bound

$$(6.71) \quad \left| \operatorname{Re} \int (-ih\nabla + \mathbf{A}_0^h) \underline{\psi}_h^\tau(\mathbf{x}^h) \cdot (\mathbf{A}^h - \mathbf{A}_0^h) \overline{\underline{\psi}_h^\tau(\mathbf{x}^h)} dx^h \right| \\ \leq C(1 + \|\mathbf{A}^h\|_{W^{3,\infty}(\operatorname{supp}(\underline{\psi}_h^\tau))}^2) h^{\min(\frac{1}{2}+3\delta, 1+\delta)} \|\underline{\psi}_h^\tau\|^2.$$

Returning to (6.62) via (6.63) and combining (6.71) with (6.66), we deduce

$$\frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\underline{\psi}_h^\tau)}{\|\underline{\psi}_h^\tau\|^2} \leq \frac{q_h[\mathbf{A}, \mathbb{R}_+^3](\underline{\psi}_h^\tau)}{\|\underline{\psi}_h^\tau\|^2} + C(1 + \|\mathbf{A}^h\|_{W^{3,\infty}(\operatorname{supp}(\underline{\psi}_h^\tau))}^2) (h^{\min(1+\delta, \frac{1}{2}+3\delta)} + h^{4\delta}).$$

Inserting the cut-off error for $q_h[\mathbf{A}, \mathbb{R}_+^3](\underline{\psi}_h^\tau)$ we obtain

$$\frac{q_h[\tilde{\mathbf{A}}, \Pi_{\mathbf{x}_0}](\underline{\psi}_h^\tau)}{\|\underline{\psi}_h^\tau\|^2} \leq hE(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) + C(1 + \|\tilde{\mathbf{A}}\|_{W^{3,\infty}(\operatorname{supp}(\underline{\psi}_h^\tau))}^2) (h^{2-2\delta} + h^{\min(1+\delta, \frac{1}{2}+3\delta)} + h^{4\delta}).$$

As already seen, we have $\|\tilde{\mathbf{A}}\|_{W^{3,\infty}(\operatorname{supp}(\underline{\psi}_h^\tau))} \leq (1 + C(\Omega)h^\delta)\|\mathbf{A}\|_{W^{3,\infty}(\Omega)}$. The quasimode \hat{f}_h on Ω is defined as in (6.41) and the estimation of Section 6.3.3 provides

$$(6.72) \quad \frac{q_h[\mathbf{A}, \Omega](\hat{f}_h)}{\|\hat{f}_h\|^2} \leq hE(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}_0}) + C(\Omega)(1 + \|\mathbf{A}\|_{W^{3,\infty}(\Omega)}^2)(h^{2-2\delta} + h^{3\delta+\frac{1}{2}} + h^{4\delta} + h^{1+\delta}).$$

Choosing $\delta = \frac{1}{3}$ and using the min-max principle we get Theorem 6.3 in situation (G1).

The proof of Theorem 6.3 is now complete since we have explored all possible geometric situations for $(\mathbf{B}_{\mathbf{x}_0}, \Pi_{\mathbf{x}})$.

6.5. Improvement for a straight polyhedron with constant magnetic field. In this paragraph we improve Theorem 6.1 for a straight polyhedral domain with constant magnetic field. Since there is no curvature, we expect smaller remainders in the asymptotics of $\lambda_h(\mathbf{B}, \Omega)$. Moreover, in that case, we will see that the function $\mathbf{x} \mapsto E(\mathbf{B}, \Pi_{\mathbf{x}})$ attains its minimum at a vertex of Ω .

Theorem 6.15. *Let Ω be a straight polyhedron and \mathbf{B} be a constant magnetic field with $|\mathbf{B}| = 1$. Then*

$$\mathcal{E}(\mathbf{B}, \Omega) = \min_{\mathbf{v} \in \mathfrak{V}} E(\mathbf{B}, \Pi_{\mathbf{v}})$$

where \mathfrak{V} denotes the set of the vertices of Ω . We have

$$\lambda_h(\mathbf{B}, \Omega) \leq h\mathcal{E}(\mathbf{B}, \Omega) + Ch^2.$$

If there exists $\mathbf{v} \in \mathfrak{V}$ such that $E(\mathbf{B}, \Pi_{\mathbf{v}}) = \mathcal{E}(\mathbf{B}, \Omega) < \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{v}})$, then there exist positive constants C, c such that

$$\lambda_h(\mathbf{B}, \Omega) \leq h\mathcal{E}(\mathbf{B}, \Omega) + Ce^{-ch^{-1/2}}.$$

Proof. Since the polyhedral domain is assumed to have straight faces and edges and the magnetic field is constant, the function $\mathbf{x} \mapsto E(\mathbf{B}, \Pi_{\mathbf{x}})$ is constant on each stratum of $\overline{\Omega}$. Let $\mathbf{v} \in \mathfrak{V}$. We apply Theorem 4.7 and relations (4.8) and (4.19) with $\Pi = \Pi_{\mathbf{v}}$:

$$E(\mathbf{B}, \Pi_{\mathbf{v}}) \leq \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{v}}) = \min_{\mathbf{e} \in \mathfrak{E}_{\mathbf{v}}} E(\mathbf{B}, \Pi_{\mathbf{e}}),$$

with $\mathfrak{E}_{\mathbf{v}}$ the subset of \mathfrak{E} such that for any $\mathbf{e} \in \mathfrak{E}_{\mathbf{v}}$, $\mathbf{v} \in \partial \mathbf{e}$ and $\Pi_{\mathbf{e}}$ the wedge associated with the edge \mathbf{e} . In the same way we prove for each edge $\mathbf{e} \in \mathfrak{E}$:

$$E(\mathbf{B}, \Pi_{\mathbf{e}}) \leq \min_{\mathbf{f} \in \mathfrak{F}_{\mathbf{e}}} E(\mathbf{B}, \Pi_{\mathbf{f}}) \leq 1$$

where $\mathfrak{F}_{\mathbf{e}}$ denotes the set of the faces adjacent to an edge \mathbf{e} . Therefore

$$\min_{\mathbf{v} \in \mathfrak{V}} E(\mathbf{B}, \Pi_{\mathbf{v}}) = \mathcal{E}(\mathbf{B}, \Omega).$$

Let \mathbf{v}_0 be a vertex minimizing $\mathbf{x} \mapsto E(\mathbf{B}, \Pi_{\mathbf{x}})$. Let $\Pi_{\mathbf{x}}$ be the tangent cone given by Theorem 4.7. If $E(\mathbf{B}, \Pi_{\mathbf{v}_0}) < \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{v}_0})$ then $\Pi_{\mathbf{x}} = \Pi_{\mathbf{v}_0}$ and we have an (admissible generalized) eigenfunction on $\Pi_{\mathbf{v}_0}$ associated with $E(\mathbf{B}, \Pi_{\mathbf{v}_0})$. If $E(\mathbf{B}, \Pi_{\mathbf{v}}) = \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{v}})$, then there exists a stratum \mathbf{t} of Ω associated with $\Pi_{\mathbf{x}}$ such that $\Pi_{\mathbf{x}}$ is the tangent cone to any point of \mathbf{t} . Moreover for any point $\mathbf{x} \in \mathbf{t}$ we have $E(\mathbf{B}, \Pi_{\mathbf{x}}) < \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{x}})$ therefore there exists a generalized eigenfunction on $\Pi_{\mathbf{x}}$ associated to $E(\mathbf{B}, \Pi_{\mathbf{x}})$. In both cases we have found a point $\mathbf{x} \in \overline{\Omega}$ such that there exists a generalized eigenfunction on $\Pi_{\mathbf{x}}$ associated to $\mathcal{E}(\mathbf{B}, \Omega)$. There exists $R_{\mathbf{x}} > 0$ such that

$$(6.73) \quad \Omega \cap \mathcal{B}(0, 2R_{\mathbf{x}}) = \Pi_{\mathbf{x}} \cap \mathcal{B}(0, 2R_{\mathbf{x}}).$$

We define the quasimode ψ_h as in (6.22) with $\delta = 0$, $\boldsymbol{\tau} = 0$ and $R = R_{\mathbf{x}}$. We have $\psi_h \in H^1(\Omega)$ and $q_h[\mathbf{A}, \Pi_{\mathbf{x}}](\psi_h) = q_h[\mathbf{A}, \Omega](\psi_h)$. Using (6.34) and the fact that \mathbf{A} equals its affine part, we have:

$$(6.74) \quad \frac{q_h[\mathbf{A}, \Omega](\psi_h)}{\|\psi_h\|^2} = hE(\mathbf{B}, \Pi_{\mathbf{x}}) + h^2 \rho_h.$$

Applying Lemma 6.9 with χ_h as defined in (6.20), $\delta = 0$ and $R = R_{\mathbf{x}}$, we have

$$(6.75) \quad \rho_h = \begin{cases} O(1) & \text{if } k < 3, \\ O(e^{-ch^{-1/2}}) & \text{if } k = 3. \end{cases}$$

Then, by the min-max principle and (6.74), we deduce when $k < 3$:

$$\lambda_h(\mathbf{B}, \Omega) \leq h \inf_{\mathbf{x} \in \overline{\Omega}} E(\mathbf{B}, \Pi_{\mathbf{x}}) + O(h^2) = h\mathcal{E}(\mathbf{B}, \Omega) + Ch^2.$$

If there exists $\mathbf{v} \in \mathfrak{V}$ such that $E(\mathbf{B}, \Pi_{\mathbf{v}}) = \mathcal{E}(\mathbf{B}, \Omega) < \mathcal{E}^*(\mathbf{B}, \Pi_{\mathbf{v}})$, we use Theorem 4.7, Proposition 4.15 and there exists an (admissible generalized) eigenfunction with $k = 3$ of $H(\mathbf{A}, \Pi_{\mathbf{v}})$ for $E(\mathbf{B}, \Pi_{\mathbf{v}})$. According to (6.74) and (6.75), we have:

$$(6.76) \quad \lambda_h(\mathbf{B}, \Omega) \leq \frac{q_h[\mathbf{A}, \Omega](\psi_h)}{\|\psi_h\|^2} \leq hE(\mathbf{B}, \Pi_{\mathbf{v}}) + Ce^{-ch^{-1/2}}.$$

□

7. Lower bound for first eigenvalues

In this section we give a general lower bound on the first eigenvalue, namely:

Theorem 7.1. *Let $\Omega \in \overline{\mathcal{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{A} \in W^{2,\infty}(\overline{\Omega})$ be a twice differentiable magnetic potential such that the associated magnetic field \mathbf{B} does not vanish on $\overline{\Omega}$. Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that*

$$(7.1) \quad \forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \geq h \mathcal{E}(\mathbf{B}, \Omega) - C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2)h^{5/4}.$$

We recall that the quantity $\mathcal{E}(\mathbf{B}, \Omega)$ is the lowest local energy defined in (1.5).

- *Idea of the proof.* We first make a partition of the unity of $\overline{\Omega}$ such that on each element we are able to use the change of variable given in (3.1). The local energy of the associated tangent model problem with frozen magnetic field is then bounded from below by $h \mathcal{E}(\mathbf{B}, \Omega)$. As above we then estimate the remainders due to the cut-off effects, the change of variables and the linearization of the potential.

- *IMS localization.* Let $\delta \in (0, \frac{1}{2})$. For h small enough, let us denote by $(\mathbf{x}_{j,h}, r_{j,h})$ a finite set of pairs (center, radius) provided by Lemma B.2 for $\rho = h^\delta$. Relying on Lemma B.7, we choose a finite associate partition of the unity $(\chi_{j,h})_j$ with $\chi_{j,h} \in \mathcal{C}_0^\infty(\mathcal{B}(\mathbf{x}_{j,h}, 2r_{j,h}))$ satisfying

$$\sum_j \chi_{j,h}^2 = 1 \quad \text{on } \overline{\Omega}$$

and the uniform estimate of gradients

$$(7.2) \quad \exists C > 0, \quad \forall h \in (0, h_0), \quad \forall j, \quad \|\nabla \chi_{j,h}\|_{L^\infty(\Omega)}^2 \leq Ch^{-2\delta}.$$

The IMS formula (see A.7) provides for all $f \in H^1(\Omega)$

$$q_h[\mathbf{A}, \Omega](f) = \sum_j q_h[\mathbf{A}, \Omega](\chi_{j,h}f) - h^2 \sum_j \|\nabla \chi_{j,h}f\|_{L^2(\Omega)}^2$$

and using (7.2) and the finite covering property of the covering we get $C(\Omega) > 0$ such that

$$(7.3) \quad q_h[\mathbf{A}, \Omega](f) \geq \sum_j q_h[\mathbf{A}, \Omega](f_{j,h}) - C(\Omega)h^{2-2\delta}\|f\|_{L^2(\Omega)}^2.$$

where $f_{j,h}$ denotes the localized function $\chi_{j,h}f$.

- *Local control of the energy.* We estimate each term $q_h[\mathbf{A}, \Omega](f_{j,h})$ appearing in (7.3). By construction $\text{supp}(f_{j,h}) \subset \mathcal{U}_{\mathbf{x}_{j,h}}$. Let $\tilde{\mathbf{A}}^{j,h}$ defined as in (A.4) with $\mathbf{x} = \mathbf{x}_{j,h}$. Lemma 6.5 applied with $r = Kr_{j,h} \leq Ch^\delta$ provides $C(\Omega) > 0$ such that

$$(7.4) \quad \frac{q_h[\mathbf{A}, \Omega](f_{j,h})}{\|f_{j,h}\|^2} \geq (1 - C(\Omega)h^\delta) \frac{q_h[\tilde{\mathbf{A}}^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h})}{\|\psi_{j,h}\|^2}$$

where we have denoted $\psi_{j,h} = f_{j,h} \circ U_{\mathbf{x}_{j,h}}^{-1}$. Let $\mathbf{A}_0^{j,h}$ be the affine part of $\tilde{\mathbf{A}}^{j,h}$ at the origin of $\Pi_{\mathbf{x}_{j,h}}$. We use (A.7) with $\psi = \psi_{j,h}$ and $\mathcal{O} = \Pi_{\mathbf{x}_{j,h}}$:

$$(7.5) \quad q_h[\tilde{\mathbf{A}}^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) = q_h[\mathbf{A}_0^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) \\ + 2 \operatorname{Re} \langle (-ih\nabla + \mathbf{A}_0^{j,h})\psi_{j,h}, (\tilde{\mathbf{A}}^{j,h} - \mathbf{A}_0^{j,h})\psi_{j,h} \rangle + \|(\tilde{\mathbf{A}}^{j,h} - \mathbf{A}_0^{j,h})\psi_{j,h}\|^2.$$

Therefore Using the Cauchy-Schwarz inequality:

$$q_h[\tilde{\mathbf{A}}^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) \geq q_h[\mathbf{A}_0^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) - 2 (q_h[\mathbf{A}_0^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}))^{1/2} \|(\tilde{\mathbf{A}}^{j,h} - \mathbf{A}_0^{j,h})\psi_{j,h}\|.$$

We cannot conclude like in (6.35) because we do not have any *a priori* upper bound on $q_h[\mathbf{A}_0^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h})$. That is why we use the parametric estimate

$$\forall \eta > 0, \quad q_h[\tilde{\mathbf{A}}^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) \geq (1 - \eta)q_h[\mathbf{A}_0^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) - \eta^{-1} \|(\tilde{\mathbf{A}}^{j,h} - \mathbf{A}_0^{j,h})\psi_{j,h}\|^2$$

based on the simple inequality $2ab \leq \eta a^2 + \eta^{-1} b^2$. Since $\operatorname{curl} \mathbf{A}_0^{j,h} = \mathbf{B}_{\mathbf{x}_{j,h}}$ we have

$$q_h[\mathbf{A}_0^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) \geq hE(\mathbf{B}_{\mathbf{x}_{j,h}}, \Pi_{\mathbf{x}_{j,h}}) \|\psi_{j,h}\|^2.$$

Moreover using (6.37) and the same arguments as in Section 6.3.2 we get

$$\|(\tilde{\mathbf{A}}^{j,h} - \mathbf{A}_0^{j,h})\psi_{j,h}\|^2 \leq C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) h^{4\delta} \|\psi_{j,h}\|^2.$$

We deduce for all $\eta > 0$:

$$q_h[\tilde{\mathbf{A}}^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) \geq (1 - \eta)hE(\mathbf{B}_{\mathbf{x}_{j,h}}, \Pi_{\mathbf{x}_{j,h}}) \|\psi_{j,h}\|^2 - \eta^{-1}C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) h^{4\delta} \|\psi_{j,h}\|^2.$$

Choosing $\eta = h^{2\delta-1/2}$ we get

$$(7.6) \quad q_h[\tilde{\mathbf{A}}^{j,h}, \Pi_{\mathbf{x}_{j,h}}](\psi_{j,h}) \geq \left(hE(\mathbf{B}_{\mathbf{x}_{j,h}}, \Pi_{\mathbf{x}_{j,h}}) - C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) h^{2\delta+1/2} \right) \|\psi_{j,h}\|^2 \\ \geq \left(h\mathcal{E}(\mathbf{B}, \Omega) - C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) h^{2\delta+1/2} \right) \|\psi_{j,h}\|^2.$$

• *Conclusion.* Combining the previous localized estimate (7.6) with (7.4) we deduce:

$$q_h[\mathbf{A}, \Omega](f_{j,h}) \geq \left(h\mathcal{E}(\mathbf{B}, \Omega) - C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) (h^{2\delta+1/2} + h^{1+\delta}) \right) \|f_{j,h}\|^2.$$

Summing up in j and using that $\sum_j f_{j,h}^2 = f^2$ we obtain

$$(7.7) \quad \sum_j q_h[\mathbf{A}, \Omega](f_{j,h}) \geq \left(h\mathcal{E}(\mathbf{B}, \Omega) - C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) (h^{2\delta+1/2} + h^{1+\delta}) \right) \|f\|^2,$$

and combining (7.7) with (7.3) we get $C(\Omega) > 0$ such that

$$(7.8) \quad \forall f \in H^1(\Omega), \\ \frac{q_h[\mathbf{A}, \Omega](f)}{\|f\|^2} \geq h\mathcal{E}(\mathbf{B}, \Omega) - C(\Omega)(1 + \|\mathbf{A}\|_{W^{2,\infty}(\Omega)}^2) (h^{2\delta+1/2} + h^{1+\delta} + h^{2-2\delta}).$$

We optimize this by taking $\delta = \frac{3}{8}$ and we deduce Theorem 6.1 from the min-max principle.

Like in the last section, we have a result using only a Hölder norm of the magnetic field:

Corollary 7.2. *Let $\Omega \in \overline{\mathfrak{D}}(\mathbb{R}^3)$ be a polyhedral domain, $\mathbf{B} \in W^{1+\alpha,\infty}(\overline{\Omega})$ be a non-vanishing Hölder continuous magnetic field of order $1 + \alpha$ with some $\alpha \in (0, 1)$. Then there exist $C(\Omega) > 0$ and $h_0 > 0$ such that*

$$(7.9) \quad \forall h \in (0, h_0), \quad \lambda_h(\mathbf{B}, \Omega) \geq h \mathcal{E}(\mathbf{B}, \Omega) - C(\Omega)(1 + \|\mathbf{B}\|_{W^{1+\alpha,\infty}(\Omega)}^2)h^{5/4}.$$

Appendix A. Technical lemmas

A.1. Gauge transform.

Lemma A.1. *Let $\mathcal{O} \subset \mathbb{R}^3$ be a domain and let φ be a regular function on $\overline{\mathcal{O}}$. Let \mathbf{A} be a regular potential. Then*

$$\forall \psi \in \text{Dom}(q_h[\mathbf{A}, \mathcal{O}]), \quad q_h[\mathbf{A} + \nabla\varphi, \mathcal{O}](e^{-i\varphi/h}\psi) = q_h[\mathbf{A}, \mathcal{O}](\psi).$$

Furthermore a function ψ is an eigenfunction for the operator $H_h(\mathbf{A}, \mathcal{O})$ if and only if $e^{-i\varphi/h}\psi$ is an eigenfunction for $H_h(\mathbf{A} + \nabla\varphi, \mathcal{O})$ associated with the same eigenvalue.

Proof. It is enough to use the formula

$$(-ih\nabla + \mathbf{A} + \nabla\varphi)(e^{-i\varphi/h}\psi) = e^{-i\varphi/h}(-ih\nabla + \mathbf{A})\psi.$$

□

For the sake of completeness we provide the following standard lemma describing the effect of a translation when the potential is affine:

Lemma A.2 (Translation). *Let $\mathcal{O} \subset \mathbb{R}^3$ be a domain and \mathbf{A} be an affine magnetic potential. Let $\mathbf{d} \in \mathbb{R}^3$ be a vector and $\mathcal{O}_d := \mathcal{O} + \mathbf{d}$ be the translated domain. For $\psi \in \text{Dom}(H_h(\mathbf{A}, \mathcal{O}))$, we define the translated function on \mathcal{O}_d by $\psi_d : \mathbf{x} \mapsto \psi(\mathbf{x} - \mathbf{d})$. Let $\varphi_d(\mathbf{x}) := \langle \mathbf{A}(\mathbf{d}) - \mathbf{A}(\mathbf{0}), \mathbf{x} \rangle$. Then*

$$q_h[\mathbf{A}, \mathcal{O}_d](e^{-i\varphi_d/h}\psi_d) = q_h[\mathbf{A}, \mathcal{O}](\psi).$$

and ψ is an eigenfunction of $H_h(\mathbf{A}, \mathcal{O})$ if and only if $e^{-i\varphi_d/h}\psi_d$ is an eigenfunction of $H_h(\mathbf{A}, \mathcal{O}_d)$.

Lemma A.3. *Let \mathcal{O} be a bounded domain such that $\mathbf{0} \in \overline{\mathcal{O}}$. Let $\mathbf{A} \in W^{3,\infty}(\mathcal{O})$ be a magnetic potential. Let \mathbf{A}_0 denote the affine part of \mathbf{A} at $\mathbf{0}$. Let ℓ be an index in $\{1, 2, 3\}$.*

(a) *There exists a change of gauge ∇F where F is a polynomial function of degree 3, so that*

- (1) *The affine part of $\mathbf{A} - \nabla F$ at $\mathbf{0}$ is still \mathbf{A}_0 ,*
- (2) *The second derivative of $\mathbf{A} - \nabla F$ with respect to u_ℓ cancels at $\mathbf{0}$:*

$$\partial_{u_\ell}^2(\mathbf{A} - \nabla F)(\mathbf{0}) = 0.$$

- (3) *The coefficients of F are bounded by $\|\mathbf{A}\|_{W^{2,\infty}(\mathcal{O})}$.*

(b) Let us choose $\ell = 1$ for instance. We have the estimate

$$(A.1) \quad |\mathbf{A}(\mathbf{u}) - \mathbf{A}_0(\mathbf{u}) - \nabla F(\mathbf{u})| \leq C(\mathcal{O}) \|\mathbf{A}\|_{W^{3,\infty}(\mathcal{O})} (|u_1|^3 + |u_1 u_2| + |u_1 u_3| + |u_2|^2 + |u_3|^2),$$

where the constant $C(\mathcal{O})$ depends only on the outer diameter of \mathcal{O} .

Proof. The Taylor expansion of \mathbf{A} at $\mathbf{0}$ takes the form

$$\mathbf{A} = \mathbf{A}_0 + \mathbf{A}^{(2)} + \mathbf{A}^{(\text{rem},3)},$$

where $\mathbf{A}^{(2)}$ is a homogeneous polynomial of degree 2 with 3 components and $\mathbf{A}^{(\text{rem},3)}$ is a remainder:

$$(A.2) \quad |\mathbf{A}^{(\text{rem},3)}(\mathbf{u})| \leq \|\mathbf{A}\|_{W^{3,\infty}(\mathcal{O})} |\mathbf{u}|^3 \quad \text{for } \mathbf{u} \in \mathcal{O}.$$

Let us write the m -th component $A_m^{(2)}$ of $\mathbf{A}^{(2)}$ as

$$A_m^{(2)}(\mathbf{u}) = \sum_{|\alpha|=2} a_{m,\alpha} u_1^{\alpha_1} u_2^{\alpha_2} u_3^{\alpha_3} \quad \text{for } \mathbf{u} = (u_1, u_2, u_3) \in \mathcal{O}.$$

(a) Now, the polynomial F can be explicitly determined. It suffices to take

$$F(\mathbf{u}) = u_\ell^2 (a_{1,\alpha^*} u_1 + a_{2,\alpha^*} u_2 + a_{3,\alpha^*} u_3 - \frac{2}{3} a_{\ell,\alpha^*} u_\ell),$$

where α^* is such that $\alpha_\ell^* = 2$ (and the other components are 0). Then

$$\nabla F(\mathbf{u}) = u_\ell^2 \begin{pmatrix} a_{1,\alpha^*} \\ a_{2,\alpha^*} \\ a_{3,\alpha^*} \end{pmatrix}$$

and point (a) of the lemma is proved.

(b) Choosing $\ell = 1$, we see that the m -th components of $\mathbf{A}^{(2)} - \nabla F$ is

$$A_m^{(2)}(\mathbf{u}) - (\nabla F)_m(\mathbf{u}) = a_{m,(1,1,0)} u_1 u_2 + a_{m,(1,0,1)} u_1 u_3 + a_{m,(0,1,1)} u_2 u_3 + a_{m,(0,2,0)} u_2^2 + a_{m,(0,0,2)} u_3^2.$$

Hence $\mathbf{A}^{(2)} - \nabla F$ satisfies the estimate

$$|(\mathbf{A}^{(2)}(\mathbf{u}) - \nabla F(\mathbf{u}))| \leq \|\mathbf{A}\|_{W^{2,\infty}(\mathcal{O})} (|u_1 u_2| + |u_1 u_3| + |u_2|^2 + |u_3|^2).$$

But

$$\mathbf{A} - \mathbf{A}_0 - \nabla F = \mathbf{A}^{(2)} - \nabla F + \mathbf{A}^{(\text{rem},3)}.$$

Therefore, with (A.2)

$$|\mathbf{A}(\mathbf{u}) - \mathbf{A}_0(\mathbf{u}) - \nabla F(\mathbf{u})| \leq \|\mathbf{A}\|_{W^{2,\infty}(\mathcal{O})} (|u_1 u_2| + |u_1 u_3| + |u_2|^2 + |u_3|^2) + \|\mathbf{A}\|_{W^{3,\infty}(\mathcal{O})} |\mathbf{u}|^3.$$

Using finally that $|\mathbf{u}|^3 \leq 12(|u_1|^3 + |u_2|^3 + |u_3|^3) \leq C(\mathcal{O})(|u_1|^3 + |u_2|^2 + |u_3|^2)$, we conclude the proof of estimate (A.1). \square

A.2. **Change of variables.** Let G be a metric of \mathbb{R}^3 , that is a 3×3 positive symmetric matrix with regular coefficients. For a smooth magnetic potential, the quadratic form of the associated magnetic Laplacian with the metric G is denoted by $q_h[\mathbf{A}, \mathcal{O}, G]$ and is defined in (1.15). The following lemma describes how this quadratic form is involved when using a change of variables:

Lemma A.4. *Let $U : \mathcal{O} \rightarrow \mathcal{O}'$, $\mathbf{u} \mapsto \mathbf{v}$ be a diffeomorphism with $\mathcal{O}, \mathcal{O}'$ domains. We denote by*

$$J := d(U^{-1})$$

the jacobian matrix of the inverse of U . Let \mathbf{A} be a smooth magnetic potential and $\mathbf{B} = \text{curl } \mathbf{A}$ the associated magnetic field. Let f be a function of $\text{Dom}(q_h[\mathbf{A}, \mathcal{O}])$ and $\psi := f \circ U^{-1}$ defined in \mathcal{O}' . For any $h > 0$ we have

$$(A.3) \quad q_h[\mathbf{A}, \mathcal{O}](f) = q_h[\tilde{\mathbf{A}}, \mathcal{O}', G](\psi) \quad \text{and} \quad \|f\|_{L^2(\mathcal{O})} = \|\psi\|_{L^2_{\mathcal{G}}(\mathcal{O}')}$$

where the new magnetic potential and the metric are respectively given by

$$(A.4) \quad \tilde{\mathbf{A}} := J^\top(\mathbf{A} \circ U^{-1}) \quad \text{and} \quad G := J^{-1}(J^{-1})^\top.$$

The magnetic field $\tilde{\mathbf{B}} = \text{curl } \tilde{\mathbf{A}}$ in the new variables is given by

$$(A.5) \quad \tilde{\mathbf{B}} := |\det J| J^{-1}(\mathbf{B} \circ U^{-1}).$$

Let $\rho > 0$, using the previous Lemma with the scaling $U_\rho := \mathbf{x} \mapsto \sqrt{\rho} \mathbf{x}$ we get

Lemma A.5. *Let $\mathcal{O} \subset \mathbb{R}^3$ be a domain and for $r > 0$, we denote by $r\mathcal{O}$ the domain $\{\mathbf{x} \in \mathbb{R}^3, \mathbf{x} = r\mathbf{x}' \text{ with } \mathbf{x}' \in \mathcal{O}\}$. Let \mathbf{B} be a constant magnetic field. Then*

$$\forall \rho > 0, \quad E(\mathbf{B}, \mathcal{O}) = \rho E\left(\frac{\mathbf{B}}{\rho}, \sqrt{\rho} \mathcal{O}\right).$$

A.3. **Miscellaneous.**

- *Orientation of the magnetic field.* Let \mathbf{B} be a magnetic field. It is known that changing \mathbf{B} into $-\mathbf{B}$ does not affect the spectrum of the associated magnetic Laplacian. More precisely we have:

Lemma A.6. *Let $\mathcal{O} \subset \mathbb{R}^3$ be a domain, \mathbf{B} be a magnetic field and \mathbf{A} an associated potential. Then $H_h(-\mathbf{A}, \mathcal{O})$ and $H_h(\mathbf{A}, \mathcal{O})$ are unitary equivalent. We have*

$$\forall \psi \in \text{Dom}(q_h[\mathbf{A}, \mathcal{O}]), \quad q_h[-\mathbf{A}, \mathcal{O}](\bar{\psi}) = q_h[\mathbf{A}, \mathcal{O}](\psi)$$

and ψ is an eigenfunction of $H_h(\mathbf{A}, \mathcal{O})$ if and only if $\bar{\psi}$ is an eigenfunction of $H_h(-\mathbf{A}, \mathcal{O})$.

- *Model linear potential.* Let us remark that if \mathbf{B} is a constant magnetic field, an associated magnetic potential is given by

$$(A.6) \quad \mathbf{A}^S(\mathbf{x}) := \frac{1}{3} \mathbf{B} \wedge \mathbf{x} .$$

Indeed we have

$$\operatorname{curl} \mathbf{A}^S = \frac{1}{3} \nabla \wedge (\mathbf{B} \wedge \mathbf{x}) = \frac{1}{3} ((\nabla \cdot \mathbf{x}) \mathbf{B} - (\nabla \cdot \mathbf{B}) \mathbf{x}) = \mathbf{B} .$$

- *Comparison between two potentials.* Let $\mathcal{O} \subset \mathbb{R}^3$ be a domain and let \mathbf{A} and \mathbf{A}' be two magnetic potentials. Then, for any function ψ of $\operatorname{Dom}(q_h[\mathbf{A}, \mathcal{O}]) \cap \operatorname{Dom}(q_h[\mathbf{A}', \mathcal{O}])$, we have:

$$(A.7) \quad q_h[\mathbf{A}, \mathcal{O}](\psi) = q_h[\mathbf{A}', \mathcal{O}](\psi) + 2 \operatorname{Re} \int_{\mathcal{O}} (-ih \nabla + \mathbf{A}') \psi(\mathbf{x}) \cdot (\mathbf{A} - \mathbf{A}')(\mathbf{x}) \overline{\psi(\mathbf{x})} \, d\mathbf{x} + \|(\mathbf{A} - \mathbf{A}') \psi\|^2 .$$

A.4. **Cut-off effect.** In this section we recall standard IMS⁶ formulas. This kind of formulas appear for Schrödinger operators in [9], but they can also be found in older works like [24]. In this section \mathbf{A} denotes a regular magnetic potential and \mathcal{O} a generic domain of \mathbb{R}^3 .

The first formula describes the effect of a partition of the unity on the energy of a function which is in the form domain, see for example [35, Lemma 3.1]:

Lemma A.7 (IMS formula). *Assume that $\chi_1, \dots, \chi_L \in \mathcal{C}^\infty(\overline{\mathcal{O}})$ are such that*

$$\sum_{\ell=1}^L \chi_\ell^2 \equiv 1 \quad \text{on } \mathcal{O} .$$

Then, for any $\psi \in \operatorname{Dom}(q_h[\mathbf{A}, \mathcal{O}])$

$$q_h[\mathbf{A}, \mathcal{O}](\psi) = \sum_{\ell=1}^L q_h[\mathbf{A}, \mathcal{O}](\chi_\ell \psi) - h^2 \sum_{\ell=1}^L \|\psi \nabla \chi_\ell\|_{L^2(\mathcal{O})}^2$$

Recall that $\operatorname{Dom}_{\text{loc}}(H_h(\mathbf{A}, \mathcal{O}))$ denotes the functions that are locally in the domain of the operator, see (1.17). In particular they satisfy the Neumann boundary condition. The second formula describes the energy of such a function when applying a cut-off function, see for example [16, (6.11)]:

Lemma A.8. *Let $\chi \in \mathcal{C}_0^\infty(\overline{\mathcal{O}})$ a real smooth function. Then for any $\psi \in \operatorname{Dom}_{\text{loc}}(H_h(\mathbf{A}, \mathcal{O}))$*

$$(A.8) \quad q_h[\mathbf{A}, \mathcal{O}](\chi \psi) = \operatorname{Re} \int_{\mathcal{O}} \chi(\mathbf{x})^2 H_h(\mathbf{A}, \mathcal{O}) \psi(\mathbf{x}) \overline{\psi(\mathbf{x})} \, d\mathbf{x} + h^2 \|\nabla \chi \psi\|_{L^2(\mathcal{O})}^2 .$$

⁶IMS stands for Ismagilov-Morgan-Sigal (or Simon)

Appendix B. Partition of unity suitable for IMS formulas

We need a preliminary definition.

Definition B.1. Let $\Omega \in \mathfrak{D}(M)$ with $M = \mathbb{R}^n$ or $M = \mathbb{S}^n$. Let $\mathbf{x} \in \overline{\Omega}$ and \mathcal{U} be an open neighborhood of \mathbf{x} in M . We say that \mathcal{U} is a *map-neighborhood* of \mathbf{x} for Ω if there exists a local smooth diffeomorphism $U_{\mathbf{x}}$ which maps the neighborhood \mathcal{U} onto a neighborhood \mathcal{V} of 0 in \mathbb{R}^n and such that

$$(B.1) \quad U_{\mathbf{x}}(\mathcal{U} \cap \Omega) = \mathcal{V} \cap \Pi_{\mathbf{x}} \quad \text{and} \quad U_{\mathbf{x}}(\mathcal{U} \cap \partial\Omega) = \mathcal{V} \cap \partial\Pi_{\mathbf{x}},$$

where $\Pi_{\mathbf{x}}$ is the tangent cone to Ω at \mathbf{x} (compare with (3.1)).

Lemma B.2. Let $n \geq 1$ be the space dimension. M denotes \mathbb{R}^n or \mathbb{S}^n . Let $\Omega \in \mathfrak{D}(M)$ and $K > 1$. There exist a positive integer L and two positive constants ρ_{\max} and $\kappa \leq 1$ (depending on Ω and K) such that for all $\rho \in (0, \rho_{\max}]$, there exists a (finite) set $\mathcal{Z} \subset \overline{\Omega} \times [\kappa\rho, \rho]$ satisfying the following three properties

- (1) We have the inclusion $\overline{\Omega} \subset \cup_{(\mathbf{x}, r) \in \mathcal{Z}} \overline{\mathcal{B}(\mathbf{x}, r)}$
- (2) For any $(\mathbf{x}, r) \in \mathcal{Z}$, the ball $\mathcal{B}(\mathbf{x}, Kr)$ is a map-neighborhood of \mathbf{x} for Ω
- (3) Each point \mathbf{x}_0 of $\overline{\Omega}$ belongs to at most L different balls $\mathcal{B}(\mathbf{x}, Kr)$.

Here are preparatory notations and lemmas.

Let $\Omega \in \mathfrak{D}(M)$ and $K > 1$. If the assertions of Lemma B.2 are true for this Ω and this K , we say that Property $\mathcal{P}(\Omega, K)$ holds. We may also specify that the assertion by the sentence

$$\text{Property } \mathcal{P}(\Omega, K) \text{ holds with parameters } (L, \rho_{\max}, \kappa).$$

Let $\mathcal{U}^* \subset \subset \mathcal{U}$ be two nested open sets. We say that the property $\mathcal{P}(\Omega, K; \mathcal{U}^*, \mathcal{U})$ holds⁷ if the assertions of Lemma B.2 are true for this Ω and this K , with discrete sets $\mathcal{Z} \subset (\mathcal{U}^* \cap \overline{\Omega}) \times [\kappa_{\Omega}\rho, \rho]$ and with (1)-(3) replaced by

- (1) We have the inclusion $\mathcal{U}^* \cap \overline{\Omega} \subset \cup_{(\mathbf{x}, r) \in \mathcal{Z}} \overline{\mathcal{B}(\mathbf{x}, r)}$
- (2) For any $(\mathbf{x}, r) \in \mathcal{Z}$, the ball $\mathcal{B}(\mathbf{x}, Kr)$ is included in \mathcal{U} and is a map-neighborhood of \mathbf{x} for Ω
- (3) Each point \mathbf{x}_0 of $\mathcal{U} \cap \overline{\Omega}$ belongs to at most L different balls $\mathcal{B}(\mathbf{x}, Kr)$.

Like above the specification is

$$\text{Property } \mathcal{P}(\Omega, K; \mathcal{U}^*, \mathcal{U}) \text{ holds with parameters } (L, \rho_{\max}, \kappa).$$

In the process of proof, we will construct coverings which are not exactly balls, but domains uniformly comparable to balls. Let us introduce the local version of this new assertion. For $0 < a \leq a'$ we say that

$$\text{Property } \mathcal{P}[a, a'](\Omega, K; \mathcal{U}^*, \mathcal{U}) \text{ holds with parameters } (L, \rho_{\max}, \kappa)$$

⁷This is the localized version of property $\mathcal{P}(\Omega, K)$.

if for all $\rho \in (0, \rho_{\max}]$, there exists a finite set $\mathcal{Z} \subset (\mathcal{U}^* \cap \overline{\Omega}) \times [\kappa_{\Omega}\rho, \rho]$ and open sets $\mathcal{D}(\mathbf{x}, r)$ satisfying the following four properties

- (1) We have the inclusion $\mathcal{U}^* \cap \overline{\Omega} \subset \bigcup_{(\mathbf{x}, r) \in \mathcal{Z}} \overline{\mathcal{D}(\mathbf{x}, r)}$
- (2) For any $(\mathbf{x}, r) \in \mathcal{Z}$, the set⁸ $\mathcal{D}(\mathbf{x}, Kr)$ is included in \mathcal{U} and is a map-neighborhood of \mathbf{x} for Ω
- (3) Each point \mathbf{x}_0 of $\mathcal{U} \cap \overline{\Omega}$ belongs to at most L different sets $\mathcal{D}(\mathbf{x}, Kr)$
- (4) For any $(\mathbf{x}, r) \in \mathcal{Z}$, we have the inclusions $\mathcal{B}(\mathbf{x}, ar) \subset \mathcal{D}(\mathbf{x}, r) \subset \mathcal{B}(\mathbf{x}, a'r)$.

Note that $\mathcal{P}[1, 1](\Omega, K; \mathcal{U}^*, \mathcal{U}) = \mathcal{P}(\Omega, K; \mathcal{U}^*, \mathcal{U})$.

Lemma B.3. *If Property $\mathcal{P}[a, a'](\Omega, K; \mathcal{U}^*, \mathcal{U})$ holds with parameters (L, ρ_{\max}, κ) , then Property $\mathcal{P}(\Omega, \frac{a}{a'}K; \mathcal{U}^*, \mathcal{U})$ holds with parameters $(L, a'\rho_{\max}, \kappa)$.*

Proof. Starting from the covering of $\mathcal{U}^* \cap \overline{\Omega}$ by the sets $\overline{\mathcal{D}(\mathbf{x}, r)}$ and using condition (4), we can consider the covering of $\mathcal{U}^* \cap \overline{\Omega}$ by the balls $\mathcal{B}(\mathbf{x}, a'r)$. Then $r' := a'r \in [\kappa a'\rho, a'\rho] = [\kappa\rho', \rho']$ with $\rho' < a'\rho_{\max}$.

Concerning conditions (2) and (3), it suffices to note the inclusions

$$\mathcal{B}(\mathbf{x}, \frac{a}{a'}Kr') \subset \mathcal{D}(\mathbf{x}, \frac{1}{a'}r'K) = \mathcal{D}(\mathbf{x}, rK).$$

The lemma is proved. □

Proof. of Lemma B.2. The principle of the proof is a recursion on the dimension n .

Step 1. Explicit construction when $n = 1$.

The domain Ω and the localizing open sets \mathcal{U}^* and \mathcal{U} are then open intervals. Let us assume for example that $\mathcal{U}^* = (-\ell, \ell)$, $\mathcal{U} = (-\ell - \delta, \ell + \delta)$ and $\Omega = (0, \ell + \delta')$ with $\ell, \delta > 0$ and $\delta' > \delta$. Let $K \geq 1$. We can take

$$\rho_{\max} = \min \left\{ \frac{\ell}{K}, \delta \right\}$$

and for any $\rho \leq \rho_{\max}$ the following set of couples (\mathbf{x}_j, r_j) , $j = 0, 1, \dots, J$

$$\mathbf{x}_0 = 0, r_0 = \rho \quad \text{and} \quad \mathbf{x}_j = \rho + \frac{2j-1}{K}\rho, r_j = \frac{\rho}{K} \quad \text{for } j = 1, \dots, J$$

with J such that $\mathbf{x}_J < \ell$ and $\rho + \frac{2J+1}{K}\rho \geq \ell$. If $\mathbf{x}_j < \ell - \frac{\rho}{K}$, we add the point $\mathbf{x}_{j+1} = \rho + \frac{2j}{K}\rho$. The covering condition (1) is obvious.

Concerning condition (2), we note that the bound $\rho_{\max} \leq \frac{\ell}{K}$ implies that $[0, Kr_0) = [0, K\rho)$ is a map-neighborhood for the boundary of Ω , and the bound $\rho_{\max} \leq \delta$ implies that when $j \geq 1$, the “balls” $(\mathbf{x}_j - Kr_j, \mathbf{x}_j + Kr_j) = (\mathbf{x}_j - \rho, \mathbf{x}_j + \rho)$ are map-neighborhoods for the interior of Ω .

Concerning condition (3), we can check that $L = K + 2$ is suitable.

⁸Here $\mathcal{D}(\mathbf{x}, Kr)$ is the set of \mathbf{y} such that $\mathbf{x} + (\mathbf{y} - \mathbf{x})/K \in \mathcal{D}(\mathbf{x}, r)$.

Step 2. Localization.

Let $\Omega \in \mathfrak{D}(\mathbb{R}^n)$ or $\Omega \in \mathfrak{D}(\mathbb{S}^n)$. For any $\mathbf{x} \in \overline{\Omega}$, there exists a ball $\mathcal{B}(\mathbf{x}, r_{\mathbf{x}})$ with positive radius $r_{\mathbf{x}}$ that is a map-neighborhood for Ω . We extract a finite covering of $\overline{\Omega}$ by open sets $\mathcal{B}(\mathbf{x}^{(\ell)}, \frac{1}{2}r^{(\ell)})$. We set

$$\mathcal{U}_{\ell}^* = \mathcal{B}(\mathbf{x}^{(\ell)}, \frac{1}{2}r^{(\ell)}) \quad \text{and} \quad \mathcal{U}_{\ell} = \mathcal{B}(\mathbf{x}^{(\ell)}, r^{(\ell)}).$$

The map $U_{\ell} := U_{\mathbf{x}^{(\ell)}}$ transforms \mathcal{U}_{ℓ}^* and \mathcal{U}_{ℓ} into neighborhoods \mathcal{V}_{ℓ}^* and \mathcal{V}_{ℓ} of 0 in the tangent cone $\Pi_{\ell} := \Pi_{\mathbf{x}^{(\ell)}}$. Thus we are reduced to prove the local property $\mathcal{P}(\Pi_{\ell}, K; \mathcal{V}_{\ell}^*, \mathcal{V}_{\ell})$ for any ℓ . Indeed

- The local diffeomorphism U_{ℓ} allows to deduce Property $\mathcal{P}(\Omega, K; \mathcal{U}_{\ell}^*, \mathcal{U}_{\ell})$ from Property $\mathcal{P}(\Pi_{\ell}, K'; \mathcal{V}_{\ell}^*, \mathcal{V}_{\ell})$ for a ratio K'/K that only depends on U_{ℓ} (this relies on Lemma B.3).
- Properties $\mathcal{P}(\Omega, K; \mathcal{U}_{\ell}^*, \mathcal{U}_{\ell})$ imply Property $\mathcal{P}(\Omega, K; \cup_{\ell} \mathcal{U}_{\ell}^*, \cup_{\ell} \mathcal{U}_{\ell}) = \mathcal{P}(\Omega, K)$ (it suffices to merge the (finite) union of the sets \mathcal{L} corresponding to each \mathcal{U}_{ℓ}).

Step 3. Core recursive argument: If Ω_0 is the section of the cone Π , Property $\mathcal{P}(\Omega_0, K)$ implies Property $\mathcal{P}(\Pi, K'; \mathcal{B}(0, 1), \mathcal{B}(0, 2))$ for a suitable ratio K'/K . We are going to prove this separately in several lemmas (B.4 to B.6). Then the proof Lemma B.2 will be complete. \square

Lemma B.4. *Let Γ be a cone in \mathfrak{P}^{n-1} . For $\ell = 1, 2$, let \mathcal{B}_{ℓ} and \mathcal{I}_{ℓ} be the ball $\mathcal{B}(0, \ell)$ of \mathbb{R}^{n-1} and the interval $(-\ell, \ell)$, respectively. We assume that Property $\mathcal{P}(\Gamma, K; \mathcal{B}_1, \mathcal{B}_2)$ holds (with parameters (L, ρ_{\max}, κ)). Then Property $\mathcal{P}[1, \sqrt{2}](\Gamma \times \mathbb{R}, K; \mathcal{B}_1 \times \mathcal{I}_1, \mathcal{B}_2 \times \mathcal{I}_2)$ holds.*

Proof. Let us denote by \mathbf{y} and z coordinates in Γ and \mathbb{R} , respectively. For $\rho \leq \rho_{\max}$, let \mathcal{Z}_{Γ} be an associate set of couples $(\mathbf{y}, r_{\mathbf{y}})$. For each \mathbf{y} we consider the unique set of equidistant points $\mathcal{Z}_{\mathbf{y}} = \{z_j \in [-1, 1], j = 1, \dots, J_{\mathbf{y}}\}$ such that

$$z_j - z_{j-1} = 2r_{\mathbf{y}} \quad \text{and} \quad z_1 + 1 = 1 - z_{J_{\mathbf{y}}} < r_{\mathbf{y}}.$$

Then we define

$$(B.2) \quad \mathcal{Z}^{(\rho)} = \{(\mathbf{x}, r_{\mathbf{x}}), \quad \text{for } \mathbf{x} = (\mathbf{y}, z) \text{ with } (\mathbf{y}, r_{\mathbf{y}}) \in \mathcal{Z}_{\Gamma}, z \in \mathcal{Z}_{\mathbf{y}} \text{ and } r_{\mathbf{x}} = r_{\mathbf{y}}\}.$$

The associate open set $\mathcal{D}(\mathbf{x}, r_{\mathbf{x}})$ is the product

$$\mathcal{D}(\mathbf{x}, r_{\mathbf{x}}) = \mathcal{B}(\mathbf{y}, r_{\mathbf{y}}) \times (z - r_{\mathbf{y}}, z + r_{\mathbf{y}}).$$

We have the inclusions $\mathcal{B}(\mathbf{x}, r_{\mathbf{x}}) \subset \mathcal{D}(\mathbf{x}, r_{\mathbf{x}}) \subset \mathcal{B}(\mathbf{x}, \sqrt{2}r_{\mathbf{x}})$ and it is easy to check that Property $\mathcal{P}[1, \sqrt{2}](\Gamma \times \mathbb{R}, K; \mathcal{B}(0, 1) \times \mathcal{I}_1, \mathcal{B}(0, 2) \times \mathcal{I}_2)$ holds with parameters $(L', \rho_{\max}, \kappa)$ with $L' = LK$. \square

Lemma B.5. *Let Ω be a section in $\mathfrak{D}(\mathbb{S}^{n-1})$, let Π be the corresponding cone, and let \mathcal{I}_{ℓ} be the interval $(2^{-\ell}, 2^{\ell})$ for $\ell = 1, 2$. We define the annuli*

$$\mathcal{A}_{\ell} = \left\{ \mathbf{x} \in \Pi, \quad |\mathbf{x}| \in \mathcal{I}_{\ell} \quad \text{and} \quad \frac{\mathbf{x}}{|\mathbf{x}|} \in \Omega \right\}.$$

We assume that Property $\mathcal{P}(\Omega, K)$ holds (with parameters (L, ρ_{\max}, κ)). Then Property $\mathcal{P}[a, a'](\Pi, K; \mathcal{A}_1, \mathcal{A}_2)$ holds for suitable constants a and a' (independent of Ω and K).

Proof. Let us consider the diffeomorphism

$$(B.3) \quad \begin{aligned} T : \Omega \times (-2, 2) &\longrightarrow \mathcal{A}_2 \\ \mathbf{x} = (\mathbf{y}, z) &\longmapsto \check{\mathbf{x}} = 2^z \mathbf{y} \end{aligned}$$

in view of proving Property $\mathcal{P}[a, a'](\Pi, K; \mathcal{A}_1, \mathcal{A}_2)$, for a given $\rho \leq \rho_{\max}$, we define a suitable set $\check{\mathcal{Z}}^{(\rho)}$ using the set $\mathcal{Z}^{(\rho)}$ introduced in (B.2)

$$(B.4) \quad \check{\mathcal{Z}}^{(\rho)} = \{(\check{\mathbf{x}}, r_{\check{\mathbf{x}}}), \text{ for } \check{\mathbf{x}} = T\mathbf{x} \text{ with } (\mathbf{x}, r_{\mathbf{x}}) \in \mathcal{Z}^{(\rho)}\},$$

and the associated open sets

$$\check{\mathcal{D}}(\check{\mathbf{x}}, r_{\check{\mathbf{x}}}) = T(\mathcal{D}(\mathbf{x}, r_{\mathbf{x}})).$$

We can check that

$$\mathcal{B}(\check{\mathbf{x}}, ar_{\check{\mathbf{x}}}) \subset \check{\mathcal{D}}(\check{\mathbf{x}}, r_{\check{\mathbf{x}}}) \subset \mathcal{B}(\check{\mathbf{x}}, a'r_{\check{\mathbf{x}}})$$

with $a = \frac{1}{8} \log 2$ and $a' = 8\sqrt{2} \log 2$ and that Property $\mathcal{P}[a, a'](\Pi, K; \mathcal{A}_1, \mathcal{A}_2)$ holds with parameters $(L', \rho_{\max}, \kappa)$ for $L' = NLK$ with an integer N independent of L and K . \square

Lemma B.6. *Let Ω be a section in $\mathfrak{D}(\mathbb{S}^{n-1})$, let Π be the corresponding cone, and let \mathcal{B}_ℓ be the balls $\mathcal{B}(0, \ell)$ of \mathbb{R}^n for $\ell = 1, 2$. We assume that Property $\mathcal{P}(\Omega, K)$ holds with parameters (L, ρ_{\max}, κ) for a $\rho_{\max} \leq 1$. Then Property $\mathcal{P}[a, a'](\Pi, K; \mathcal{B}_1, \mathcal{B}_2)$ holds for suitable constants a and a' (independent of Ω and K) and with parameters $(L', 1, \kappa\rho_{\max})$.*

Proof. Let $\rho \leq 1$ and let M be the natural number such that

$$2^{-M-1} < \rho \leq 2^{-M}.$$

On the model of (B.3)-(B.4), we set

$$\check{\mathcal{Z}}^m = \{(2^{-m}T\mathbf{x}, 2^{-m}r_{\mathbf{x}}), \text{ with } (\mathbf{x}, r_{\mathbf{x}}) \in \mathcal{Z}^{(2^m\rho_{\max}\rho)}\}, \quad m = 0, \dots, M,$$

and the associated open sets are

$$(B.5) \quad 2^{-m}T(\mathcal{D}(\mathbf{x}, r_{\mathbf{x}})) \text{ with } (\mathbf{x}, r_{\mathbf{x}}) \in \mathcal{Z}^{(2^m\rho_{\max}\rho)}.$$

The set $\check{\mathcal{Z}}$ associated with the cone Π in the ball \mathcal{B}_1 is

$$\{(0, \rho)\} \cup \bigcup_{m=0}^M \check{\mathcal{Z}}^m$$

and the associated open sets are the reunion of the sets (B.5) for $m = 0, \dots, M$ and of the ball $\mathcal{B}(0, \rho)$. As the radii $r_{\check{\mathbf{x}}}$ belong to $[\kappa 2^m \rho_{\max} \rho, 2^m \rho_{\max} \rho]$, we have $2^{-m}r_{\check{\mathbf{x}}} \in [\kappa \rho_{\max} \rho, \rho_{\max} \rho]$. Since ρ itself belongs to the full collection of radii r , we finally find $r \in [\kappa \rho_{\max} \rho, \rho]$. The finite covering holds with $L' = 3NLK + 1$ for the same integer N appearing at the end of the proof of Lemma B.5. \square

Lemma B.7. *Let $\Omega \in \mathfrak{D}(\mathbb{R}^n)$. Let (L, ρ_{\max}, κ) be the parameters provided by Lemma B.2, for the Property $\mathcal{P}(\Omega, 2)$ to hold. For any $\rho \in (0, \rho_{\max}]$ let $\mathcal{Z} \subset \bar{\Omega} \times [\kappa\rho, \rho]$ be an associate set of pairs (center, radius). Then there exists a collection of smooth functions $(\chi_{(\mathbf{x}, r)})_{(\mathbf{x}, r) \in \mathcal{Z}}$ with $\chi_{(\mathbf{x}, r)} \in \mathcal{C}_0^\infty(\mathcal{B}(\mathbf{x}, 2r))$ satisfying the identity (partition of unity)*

$$\sum_{(\mathbf{x}, r) \in \mathcal{Z}} \chi_{(\mathbf{x}, r)}^2 = 1 \quad \text{on } \bar{\Omega}$$

and the uniform estimate of gradients

$$\exists C > 0, \quad \forall (\mathbf{x}, r) \in \mathcal{Z}, \quad \|\nabla \chi_{(\mathbf{x}, r)}\|_{L^\infty(\Omega)} \leq C\rho^{-1},$$

where C only depends on Ω . By construction any ball $\mathcal{B}(\mathbf{x}, 2r)$ is a map-neighborhood of \mathbf{x} for Ω .

Proof. Let $\xi_{(\mathbf{x}, r)} \in \mathcal{C}_0^\infty(\mathcal{B}(\mathbf{x}, 2r))$, with the property that $\xi_{(\mathbf{x}, r)} \equiv 1$ in $\mathcal{B}(\mathbf{x}, r)$, and satisfying

$$\|\nabla \xi_{(\mathbf{x}, r)}\|_{L^\infty(\mathbb{R}^3)} \leq Cr^{-1}$$

where C is a universal constant. Then we set for each $(\mathbf{x}_0, r_0) \in \mathcal{Z}$

$$\chi_{(\mathbf{x}_0, r_0)} = \frac{\xi_{(\mathbf{x}_0, r_0)}}{(\sum_{(\mathbf{x}, r) \in \mathcal{Z}} \xi_{(\mathbf{x}, r)}^2)^{1/2}}.$$

Due to property (1) in Lemma B.2, $\sum_{(\mathbf{x}, r) \in \mathcal{Z}} \xi_{(\mathbf{x}, r)}^2 \geq 1$ and due to property (3),

$$\left\| \sum_{(\mathbf{x}, r) \in \mathcal{Z}} \nabla \xi_{(\mathbf{x}, r)}^2 \right\|_{L^\infty(\mathbb{R}^3)} \leq CL_\Omega.$$

We deduce the lemma. □

References

- [1] S. Agmon. *Bounds on exponential decay of eigenfunctions of Schrödinger operators*, volume 1159 of *Lecture Notes in Math*. Springer, Berlin 1985.
- [2] J. Aramaki. Asymptotics of the eigenvalues for the Neumann Laplacian with non-constant magnetic field associated with superconductivity. *Far East J. Math. Sci. (FJMS)* **25**(3) (2007) 529–584.
- [3] A. Bernoff, P. Sternberg. Onset of superconductivity in decreasing fields for general domains. *J. Math. Phys.* **39**(3) (1998) 1272–1284.
- [4] V. Bonnaillie. On the fundamental state energy for a Schrödinger operator with magnetic field in domains with corners. *Asymptot. Anal.* **41**(3-4) (2005) 215–258.
- [5] V. Bonnaillie-Noël, M. Dauge. Asymptotics for the low-lying eigenstates of the Schrödinger operator with magnetic field near corners. *Ann. Henri Poincaré* **7** (2006) 899–931.
- [6] V. Bonnaillie-Noël, M. Dauge, D. Martin, G. Vial. Computations of the first eigenpairs for the Schrödinger operator with magnetic field. *Comput. Methods Appl. Mech. Engrg.* **196**(37-40) (2007) 3841–3858.
- [7] V. Bonnaillie-Noël, M. Dauge, N. Popoff, N. Raymond. Discrete spectrum of a model Schrödinger operator on the half-plane with Neumann conditions. *ZAMP* **63**(2) (2012) 203–231.
- [8] M. Costabel, A. McIntosh. On Bogovskii and regularized Poincaré integral operators for de Rham complexes on Lipschitz domains. *Math. Z.* **265**(2) (2010) 297–320.
- [9] H. Cycon, R. Froese, W. Kirsch, B. Simon. *Schrödinger operators with application to quantum mechanics and global geometry*. Texts and Monographs in Physics. Springer-Verlag, Berlin, study edition 1987.
- [10] M. Dauge. *Elliptic boundary value problems on corner domains*, volume 1341 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin 1988. Smoothness and asymptotics of solutions.
- [11] M. Dauge, B. Helffer. Eigenvalues variation. I. Neumann problem for Sturm-Liouville operators. *J. Differential Equations* **104**(2) (1993) 243–262.
- [12] N. Dombrowski, N. Raymond. Semiclassical analysis with vanishing magnetic fields. *J. Spectr. Theory* **3**(3) (2013) 423–464.
- [13] S. Fournais, B. Helffer. Accurate eigenvalue estimates for the magnetic Neumann Laplacian. *Annales Inst. Fourier* **56**(1) (2006) 1–67.
- [14] S. Fournais, B. Helffer. *Spectral methods in surface superconductivity*. Progress in Nonlinear Differential Equations and their Applications, 77. Birkhäuser Boston Inc., Boston, MA 2010.
- [15] B. Helffer, A. Mohamed. Semiclassical analysis for the ground state energy of a Schrödinger operator with magnetic wells. *J. Funct. Anal.* **138**(1) (1996) 40–81.
- [16] B. Helffer, A. Morame. Magnetic bottles in connection with superconductivity. *J. Funct. Anal.* **185**(2) (2001) 604–680.
- [17] B. Helffer, A. Morame. Magnetic bottles for the Neumann problem: the case of dimension 3. *Proc. Indian Acad. Sci. Math. Sci.* **112**(1) (2002) 71–84. Spectral and inverse spectral theory (Goa, 2000).
- [18] B. Helffer, A. Morame. Magnetic bottles for the Neumann problem: curvature effects in the case of dimension 3 (general case). *Ann. Sci. École Norm. Sup. (4)* **37**(1) (2004) 105–170.
- [19] H. Jadallah. The onset of superconductivity in a domain with a corner. *J. Math. Phys.* **42**(9) (2001) 4101–4121.
- [20] L. D. Landau, E. M. Lifshitz. *Quantum mechanics: non-relativistic theory. Course of Theoretical Physics, Vol. 3*. Addison-Wesley Series in Advanced Physics. Pergamon Press Ltd., London-Paris 1958. Translated from the Russian by J. B. Sykes and J. S. Bell.
- [21] K. Lu, X.-B. Pan. Eigenvalue problems of Ginzburg-Landau operator in bounded domains. *J. Math. Phys.* **40**(6) (1999) 2647–2670.
- [22] K. Lu, X.-B. Pan. Surface nucleation of superconductivity in 3-dimensions. *J. Differential Equations* **168**(2) (2000) 386–452. Special issue in celebration of Jack K. Hale's 70th birthday, Part 2 (Atlanta, GA/Lisbon, 1998).
- [23] V. G. Maz'ya, B. A. Plamenevskii. Elliptic boundary value problems on manifolds with singularities. *Probl. Mat. Anal.* **6** (1977) 85–142.

- [24] A. Melin. Lower bounds for pseudo-differential operators. *Arkiv för Matematik* **9**(1) (1971) 117–140.
- [25] S. A. Nazarov, B. A. Plamenevskii. *Elliptic Problems in Domains with Piecewise Smooth Boundaries*. Expositions in Mathematics 13. Walter de Gruyter, Berlin 1994.
- [26] X.-B. Pan. Upper critical field for superconductors with edges and corners. *Calc. Var. Partial Differential Equations* **14**(4) (2002) 447–482.
- [27] A. Persson. Bounds for the discrete part of the spectrum of a semi-bounded Schrödinger operator. *Math. Scand.* **8** (1960) 143–153.
- [28] N. Popoff. *Sur l'opérateur de Schrödinger magnétique dans un domaine diédral*. PhD thesis, Université de Rennes 1 2012.
- [29] N. Popoff. The Schrödinger operator on a wedge with a tangent magnetic field. *J. Math. Phys.* **54**(4) (2013).
- [30] N. Popoff. The model Magnetic Laplacian on wedges. Research Report posted on HAL: <http://hal.archives-ouvertes.fr/hal-00845305>, 2013.
- [31] N. Popoff, N. Raymond. When the 3d-magnetic laplacian meets a curved edge in the semi-classical limit. *To app. in SIAM J. Math. Anal.* (2013).
- [32] N. Raymond. Sharp asymptotics for the Neumann Laplacian with variable magnetic field: case of dimension 2. *Ann. Henri Poincaré* **10**(1) (2009) 95–122.
- [33] N. Raymond. On the semi-classical 3D Neumann Laplacian with variable magnetic field. *Asymptotic Analysis* **68**(1-2) (2010) 1– 40.
- [34] D. Saint-James, P.-G. de Gennes. Onset of superconductivity in decreasing fields. *Physics Letters* **7** (Dec. 1963) 306–308.
- [35] B. Simon. Semiclassical analysis of low lying eigenvalues. I. Nondegenerate minima: asymptotic expansions. *Ann. Inst. H. Poincaré Sect. A (N.S.)* **38**(3) (1983) 295–308.