J. Phys. A: Math. Theor. 47 (2014) 155203 (19pp)

doi:10.1088/1751-8113/47/15/155203

# Spectral asymptotics of a broken $\delta$ -interaction

# V Duchêne<sup>1</sup> and N Raymond<sup>2</sup>

<sup>1</sup> IRMAR, CNRS, Université de Rennes 1, Campus de Beaulieu, F-35042 Rennes cedex, France

<sup>2</sup> IRMAR, Université de Rennes 1, Campus de Beaulieu, F-35042 Rennes cedex, France

E-mail: vincent.duchene@univ-rennes1.fr and nicolas.raymond@univ-rennes1.fr

Received 18 December 2013, revised 4 March 2014 Accepted for publication 7 March 2014 Published 28 March 2014

#### Abstract

This paper is concerned with the spectral analysis of a Hamiltonian with a  $\delta$ -interaction supported along a broken line with angle  $\theta$ . The bound states with energy slightly below the threshold of the essential spectrum are estimated in the semiclassical regime  $\theta \rightarrow 0$ .

Keywords: delta interaction, semiclassical analysis, bound states PACS numbers: 03.65.Sq, 02.30.Xx

# 1. Motivation and results

#### 1.1. Motivation

1.1.1. Why breaking the  $\delta$ -interaction? The  $\delta$ -interaction supported on various geometries has attracted a lot of interest recently as an alternative to standard quantum graphs (see for instance [4]). In particular the reader may consult the review by Exner [12] for an introduction to leaky quantum graphs and the lecture notes by Post [23] for convergence results between the two objects. Our aim is to investigate the spectrum of a broken  $\delta$ -interaction. Before defining the main operator analyzed in this paper we shall present our initial motivation. In the paper by Exner and Němcová [16, section 5] (see also their related paper [15]) the authors were concerned by the existence and estimates of the discrete spectrum of a Hamiltonian with a  $\delta$ -interaction supported on a star. In particular they analyzed the simple case of a star with two branches in section 5.2 for which their general result establishes the existence of discrete spectrum below the essential spectrum (see also [13] for the case when the  $\delta$ -interaction is supported on a curve). What's more is that they prove that the number of bound states tends to infinity when the angle between two branches of their stars is small: they even get an explicit lower bound (see [16, remark 5.10]). Moreover they also provide numerical simulations of the eigenvalues and eigenfunctions (see [16, figures 8 and 11] and also [15, figures 1 and 4]). The spectral behaviors which show up there should be compared with recent results about broken waveguides by Dauge, Lafranche and Raymond [9, figure 11] and [10] where similar phenomena are observed. In this work, we will precisely quantify the number of eigenvalues generated by the breaking of the support of a  $\delta$ -interaction and provide their asymptotic expansions when the breaking is strong (such spectral questions are quite natural as we can see in the related works [14] and [19]). We will complete the considerations of [16] (and also [7]) when the number of branches is two thanks to the light of semiclassical analysis. At the same time the present paper will provide some insight into Open problem 7.3 in [12].

1.1.2. Definition of the main operator. Let us now define our main operator. For  $\alpha > 0$ , we introduce the following quadratic form

$$\mathcal{Q}_{\theta,\alpha}(\psi) = \int_{\mathbb{R}^2} |\nabla\psi|^2 \,\mathrm{d} u \,\mathrm{d} v - \alpha \int_{\mathbb{R}} |\psi(|s|\cos\theta, s\sin\theta)|^2 \,\mathrm{d} s, \quad \forall \psi \in H^1(\mathbb{R}^2), \tag{1.1}$$

where  $\theta \in (0, \frac{\pi}{2})$  is the breaking angle. This is well-known that  $\mathcal{Q}_{\theta,\alpha}$  is semi-bounded (see [6]). In particular we may consider its Friedrichs extension  $\mathcal{H}_{\theta,\alpha}$ . We can formally write

$$\mathcal{H}_{\theta,\alpha} = -\Delta - \alpha \delta_{\Sigma_{\theta}},$$

where

$$\Sigma_{\theta} = \{ (|s| \cos \theta, s \sin \theta), \quad s \in \mathbb{R} \}.$$

The following characterization of the essential spectrum is well-known (see [13]).

Lemma 1.1. We have

$$\sigma_{\mathsf{ess}}(\mathcal{H}_{\theta,\alpha}) = \left[-\frac{\alpha^2}{4}, +\infty\right).$$

We would like to describe the spectrum below the essential spectrum in the strong breaking limit  $\theta \rightarrow 0$ . For that purpose we shall perform the following rescaling:

$$x = \alpha \frac{\sin \theta}{\cos^2 \theta} u, \quad y = \alpha \frac{1}{\cos \theta} v,$$
 (1.2)

which permits to rephrase the problem into a semiclassical problem. We introduce the unitary transform, defined for  $\psi \in L^2(\mathbb{R}^2)$  by

$$U_{\theta,\alpha}\psi(x,y) = \frac{\cos^{3/2}\theta}{\alpha\sin^{1/2}\theta}\psi\left(\frac{\cos^2\theta}{\alpha\sin\theta}x,\frac{\cos\theta}{\alpha}y\right).$$

We have  $\mathcal{H}_{\theta,\alpha} = \alpha^2 (1 + h^2) U_{\theta,\alpha}^{-1} \mathfrak{H}_h U_{\theta,\alpha}$  where  $\mathfrak{H}_h$  is the Friedrichs extension of the rescaled quadratic form:

$$\mathfrak{Q}_{h}(\psi) = \int_{\mathbb{R}^{2}} h^{2} |\partial_{x}\psi|^{2} + |\partial_{y}\psi|^{2} \,\mathrm{d}x \,\mathrm{d}y - \int_{\mathbb{R}} |\psi(|s|, s)|^{2} \,\mathrm{d}s, \quad \forall \psi \in H^{1}(\mathbb{R}^{2}), \tag{1.3}$$

and where  $h = \tan \theta$ . Formally we may write

$$\mathfrak{H}_h = -h^2 \partial_x^2 - \partial_y^2 - \delta_{\Sigma_{\frac{\pi}{4}}}.$$
(1.4)

In particular, we notice that:

$$\sigma_{\mathsf{ess}}(\mathfrak{H}_h) = \left[-\frac{1}{4(1+h^2)}, +\infty\right).$$

**Notation 1.2.** We denote by  $\lambda_n(h)$  the nth eigenvalue, if it exists, of  $\mathfrak{H}_h$ . More generally for a semi-bounded quadratic form  $\mathfrak{Q}_h^{\natural}$ , we denote by  $\mathfrak{H}_h^{\natural}$  the corresponding Friedrichs extension and by  $\lambda_n^{\natural}(h)$  the nth eigenvalue, if it exists. Let us also recall the min–max characterization of the nth eigenvalue. We have

$$\lambda_n^{\natural}(h) = \inf_{\substack{G \subset \mathsf{Dom}(\mathfrak{Q}_h^{\natural}) \\ \dim G = n}} \sup_{\psi \in G} \frac{\mathfrak{Q}_h^{\natural}(\psi)}{\|\psi\|^2}.$$

Here and below  $\|\cdot\|$  denotes the standard  $L^2$  norm on  $\mathbb{R}^2$  while  $\|\cdot\|_{L^2(\mathbb{R})}$  is the  $L^2$  norm on  $\mathbb{R}$ . We will also denote by  $\langle\cdot,\cdot\rangle_{L^2(\mathbb{R}_v)}$  the partial scalar product defined by:

$$\langle \psi_1, \psi_2 \rangle_{L^2(\mathbb{R}_y)} = \int_{\mathbb{R}_y} \psi_1(x, y) \psi_2(x, y) \,\mathrm{d}y$$

and by  $\|\cdot\|_{L^2(\mathbb{R}_v)}$  the corresponding norm.

By using this semiclassical reformulation we will easily get an explicit lower bound for  $Q_{\theta,\alpha}$ .

**Proposition 1.3.** For all  $\psi \in H^1(\mathbb{R}^2)$  and  $\theta \in (0, \frac{\pi}{2})$ :

$$\mathcal{Q}_{\theta,\alpha}(\psi) \geqslant -\frac{\alpha^2}{\cos^2 \theta} \|\psi\|^2.$$

**Remark 1.4.** In fact this lower bound permits to define directly the Friedrichs extension associated with  $\mathcal{H}_{\theta,\alpha}$  without using the general result of [6]. This lower bound degenerates when  $\theta$  goes to  $\frac{\pi}{2}$  but, as we will see, it is more and more accurate when  $\theta$  goes to 0. A fine lower bound (independently from  $\theta$ ) is obtained in [20]. In the regime  $\theta \to 0$ , an easy corollary of one of our main results will provide a description of the optimal lower bound.

#### 1.2. Main results and organization of the paper

Let us now state the main results of this paper. Our first result is an estimate of the number of eigenvalues of  $\mathfrak{H}_h$  below the threshold of the essential spectrum. For this purpose we shall introduce some notation.

**Notation 1.5.** We denote by  $W : [-e^{-1}, +\infty) \rightarrow [-1, +\infty)$  the Lambert function defined as the inverse of  $[-1, +\infty) \ni w \mapsto w e^w \in [-e^{-1}, +\infty)$ .

**Notation 1.6.** *Given*  $\mathfrak{H}$  *a semi-bounded self-adjoint operator and*  $a < \inf \sigma_{ess}(\mathfrak{H})$ *, we denote*  $\mathcal{N}(\mathfrak{H}, a) = \#\{\lambda \in \sigma(\mathfrak{H}) : \lambda \leq a\} < +\infty.$ 

The eigenvalues are counted with multiplicity.

**Theorem 1.7.** There exists  $M_0 > 0$  such that for all  $C(h) \ge M_0 h$  with  $C(h) \xrightarrow[h \to 0]{} C_0 \ge 0$ :

$$\mathcal{N}\left(\mathfrak{H}_{h}, -\frac{1}{4} - C(h)\right) \underset{h \to 0}{\sim} \frac{1}{\pi h} \int_{x=0}^{+\infty} \sqrt{\left(-\frac{1}{4} - C_{0} + \left(\frac{1}{2} + \frac{1}{2x}W(xe^{-x})\right)^{2}\right)_{+}} dx,$$

with the notation  $f_+(x) = \max\{0, f(x)\}.$ 

**Remark 1.8.** It is important to notice that in the above result, we estimate the counting function below a potentially moving (w.r.t. *h*) threshold. In particular, the distance between  $-\frac{1}{4} - C(h)$  and the bottom of the essential spectrum is allowed to vanish in the semiclassical limit. Therefore our statement is slightly unusual as customary results would typically concern  $\mathcal{N}(\mathfrak{H}, E)$  with *E* fixed and satisfying  $E < -\frac{1}{4}$ , so as to insure a fixed security distance to the bottom of the essential spectrum (see for instance the related work [22]).

**Remark 1.9.** In the small angle limit, this result is a refinement of [16, remark 5.10]. Indeed, Exner and Němcová show that the number of bound states grows as  $n \gtrsim \frac{C}{\pi h}$  with  $C = \frac{3^{3/2}}{8\sqrt{5}} \approx 0.290$ , whereas our result implies a better constant  $C \approx 1.379$ .

Our second result concerns the asymptotics of the low lying spectrum of  $\mathfrak{H}_h$ . Let us first recall the definition of the Airy operator.

**Notation 1.10.** The Airy operator is the Dirichlet realization on  $L^2((0, +\infty))$  of  $-\partial_x^2 + x$ . Its *n*th eigenvalue is nothing but the absolute value of the nth zero (counted in decreasing order), denoted by  $z_{Ai}(n)$ , of the standard Airy function.

**Theorem 1.11.** For all  $n \ge 1$ , we have:

$$\lambda_n(h) \underset{h \to 0}{=} -1 + 2^{2/3} z_{\mathsf{A}\mathsf{i}}(n) h^{2/3} + O(h).$$

**Remark 1.12.** This asymptotic expansion explains the behavior of the spectral curves of [16, figure 8] when the angle approaches zero: the behavior of the first eigenvalues is governed by the Airy operator. Our result is a refinement (in the small angle limit) of [7] since we have an accurate description of the first eigenvalues and not only an upper bound of the first one (see also the upper bounds of the first eigenvalue obtained in [8] for star graphs).

From theorem 1.11 this is possible to deduce a quasi-tensorial structure of the first eigenfunctions.

**Theorem 1.13.** For all  $C_0 > 0$ , there exist  $h_0 > 0$ , C > 0 such that for all  $h \in (0, h_0)$  and all eigenpairs  $(\lambda, \psi)$  such that  $\lambda \leq -1 + C_0 h^{2/3}$ , we have

$$\int_{\mathbb{R}^2} |\psi - \Pi_0 \psi|^2 \, \mathrm{d}x \, \mathrm{d}y \leqslant C h^{2/3} \|\psi\|^2,$$

where  $\Pi_0 \psi = \langle \psi, e^{-|y|} \rangle_{L^2(\mathbb{R}_y)} e^{-|y|}$ .

*Remarks on*  $\delta$ *-interactions on crossing lines.* Let us consider, as in [20] and after the rescaling (1.2), the following quadratic form, defined for  $\psi \in H^1(\mathbb{R}^2)$  by

$$\mathfrak{Q}_h^{\times}(\psi) = \int_{\mathbb{R}^2} h^2 |\partial_x \psi|^2 + |\partial_y \psi|^2 \, \mathrm{d}x \, \mathrm{d}y - \int_{\mathbb{R}} \left( |\psi(-s,s)|^2 + |\psi(s,s)|^2 \right) \, \mathrm{d}s$$

The strategy of our proofs can apply modulo straightforward modifications and we get the following asymptotics

$$\mathcal{N}\left(\mathfrak{H}_{h}^{\times},-\frac{1}{4}-C(h)\right) \underset{h\to 0}{\sim} \frac{2}{\pi h} \int_{x=0}^{+\infty} \sqrt{\left(-\frac{1}{4}-C_{0}+\left(\frac{1}{2}+\frac{1}{2x}W(xe^{-x})\right)^{2}\right)_{+}} dx.$$

In the same way, we have, for  $n \ge 1$ ,

$$\lambda_{2n}^{\times}(h) = -1 + 2^{2/3} z_{\mathsf{Ai}}(n) h^{2/3} + O(h),$$

and

$$\lambda_{2n-1}^{\times}(h) = -1 + 2^{2/3} z_{\mathsf{A}\mathsf{i}'}(n) h^{2/3} + O(h)$$

where  $z_{Ai'}(n)$  is the absolute value of the *n*th zero (counted in decreasing order) of the derivative of the Airy function.

*Philosophy of the proofs.* Let us now discuss the general philosophy of the proofs. As suggested by the expression (1.4), the main ingredient in this paper is a dimensional reduction in the spirit of the famous Born–Oppenheimer approximation (see [5, 18, 21]). Such dimensional reductions where used by Balazard-Konlein in [3] in a pseudo-differential context (and thus in a very regular framework) to estimate numbers of eigenvalues. Let us also mention the paper by Morame and Truc [22] where this kind of questions appears (with a regular electric potential). It turns out that our framework is strongly excluded by the assumptions of [3] since the  $\delta$ -interaction is not even an electric potential. Nevertheless we will see that a pure variational analysis can overturn this difficulty.

Organization of the paper. This paper is organized as follows. In section 2 we introduce the double  $\delta$ -well in dimension one and we recall their basic spectral properties. In particular we will prove proposition 1.3. Section 3 is devoted to the dimensional reduction of  $\mathfrak{H}_h$  to model operators in dimension one (see proposition 3.5). Finally section 4 is concerned with the analysis of one-dimensional operators and with the proof of theorems 1.7, 1.11 and 1.13.

# 2. Double δ-well

For  $x \ge 0$ , we introduce the quadratic form  $\mathfrak{q}_x$  defined for  $\psi \in H^1(\mathbb{R})$  by

$$q_{x}(\psi) = \int_{\mathbb{R}} |\psi'(y)|^{2} \, \mathrm{d}y - |\psi(-x)|^{2} - |\psi(x)|^{2}.$$
(2.1)

This is standard (see [2, chapter II.2] and also [6]) that  $q_x$  is a semi-bounded and closed quadratic form on  $H^1(\mathbb{R})$ . Therefore we may introduce the associated self-adjoint operator denoted by  $\mathfrak{D}_x$  whose domain is

 $\mathsf{Dom}(\mathfrak{D}_x) = \{ \psi \in H^1(\mathbb{R}) \cap H^2(\mathbb{R} \setminus \{\pm x\}) : \lim_{\varepsilon \searrow 0} \left( \psi'(\pm x + \varepsilon) - \psi'(\pm x - \varepsilon) \right) = -\psi(\pm x) \}$ 

and defined as  $\mathfrak{D}_x \psi(y) = -\psi''(y)$ . We can write formally

 $\mathfrak{D}_x = -\partial_y^2 - \delta_{-x} - \delta_x.$ 

Let us describe the spectrum of  $\mathfrak{D}_x$ . The following lemma is obvious.

**Lemma 2.1.** For all  $x \ge 0$ , the essential spectrum of  $\mathfrak{D}_x$  is given by

$$\sigma_{\mathsf{ess}}(\mathfrak{D}_x) = [0, +\infty).$$

**Notation 2.2.** For  $x \ge 0$ , we denote by  $\mu_1(x)$  the lowest eigenvalue of  $\mathfrak{D}_x$  and by  $u_x$  the corresponding positive and  $L^2$ -normalized eigenfunction.

In fact we can give an explicit expression of the pair  $(\mu_1(x), u_x)$ . The following proposition is essentially well-known, except maybe its last two points.

**Proposition 2.3.** *For*  $x \ge 0$ *, we have* 

$$\mu_1(x) = -\left(\frac{1}{2} + \frac{1}{2x}W(xe^{-x})\right)^2.$$

*The second eigenvalue*  $\mu_2(x)$  *only exists for* x > 1 *and is given by* 

$$\mu_2(x) = -\left(\frac{1}{2} + \frac{1}{2x}W(-xe^{-x})\right)^2$$

By convention we set  $\mu_2(x) = 0$  when  $x \leq 1$ . In particular we have the following properties (see illustration in figure 1):



**Figure 1.** The eigenvalues of  $\mathfrak{D}_x$  as functions of *x*:  $\mu_1(x)$  (solid) and  $\mu_2(x)$  (dashed).

(1) 
$$\mu_1(x) = -1 + 2x + O(x^2),$$
  
(2)  $\mu_1(x) = -\frac{1}{4} - \frac{e^{-x}}{2} + O(xe^{-2x}), \ \mu_2(x) = -\frac{1}{4} + \frac{e^{-x}}{2} + O(xe^{-2x}),$   
(3) For all  $x \ge 0, -1 \le \mu_1(x) < -\frac{1}{4}$  and for all  $x > 1, \ \mu_2(x) > -\frac{1}{4},$   
(4)  $\mu_1$  admits a unique minimum at 0,  
(5) For all  $x \ge 0$  and all  $f \in H^1(\mathbb{R}),$  we have  $q_x(f) \ge -\|f\|_{L^2(\mathbb{R})}^2,$   
(6)  $R(x) := \|\partial_x u_x\|_{L^2(\mathbb{R}_y)}^2$  defines a bounded function for  $x > 0,$   
(7)  $\|\partial_y u_x\|_{L^2(\mathbb{R}_y)}^2$  defines a bounded function for  $x \ge 0.$ 

Proof. Let us solve the eigenvalue equation

$$\mathfrak{D}_x\psi_x=-\lambda_x\psi_x.$$

Up to multiplicative constants and using the continuity of the elements of  $\mathsf{Dom}(\mathfrak{D}_x)$  we have the alternative

$$\psi_x = \psi_{x,1}$$
 or  $\psi = \psi_{x,2}$ .

where

$$\psi_{x,1}(y) = \begin{cases} e^{\sqrt{\lambda_x}(x+y)} & \text{if } y \leqslant -x\\ \frac{1}{\cosh(\sqrt{\lambda_x}x)} \cosh(\sqrt{\lambda_x}y) & \text{if } -x < y < x\\ e^{\sqrt{\lambda_x}(x-y)} & \text{if } y \geqslant x \end{cases}$$

and

$$\psi_{x,2}(y) = \begin{cases} e^{\sqrt{\lambda_x}(x+y)} & \text{if } y \leq -x \\ \frac{-1}{\sinh(\sqrt{\lambda_x}x)} \sinh(\sqrt{\lambda_x}y) & \text{if } -x < y < x \\ -e^{\sqrt{\lambda_x}(x-y)} & \text{if } y \geq x. \end{cases}$$

In the case  $\psi = \psi_{x,1}$ , the condition at  $\pm x$  becomes

$$(2\sqrt{\lambda}_{x,1}-1)\,\mathrm{e}^{2\sqrt{\lambda}_{x,1}x}=1,$$

and we see that  $\sqrt{\lambda_{x,1}} \ge \frac{1}{2}$ . In terms of the Lambert function, we have

$$\sqrt{\lambda_{x,1}} = \frac{1}{2} + \frac{1}{2x}W(xe^{-x}) =: \sqrt{-\mu_1(x)}.$$

In the case  $\psi = \psi_{x,2}$  we find in the same way, for x > 1,

$$\sqrt{\lambda_{x,2}} = \frac{1}{2} + \frac{1}{2x}W(-xe^{-x}) =: \sqrt{-\mu_2(x)}.$$

Recall that for  $x \in (0, 1]$ , we set  $\sqrt{\lambda_{x,2}} = 0$ . In addition, we find  $\sqrt{\lambda_{x,2}} \leq \frac{1}{2}$  for x > 1.

This is very standard to establish the points (1), (2), (3). For the point (4), we notice that  $\mu_1(x) = -1$  for x > 0 is equivalent to  $W(xe^{-x}) = x$  which admits no solution for x > 0. The point (5) is then obvious.

Let us now prove the point (6). We notice that  $\psi_{x,1}(y)$  can be rewritten in the form

$$\psi_{x,1}(y) = H(-x-y)e^{\sqrt{-\mu_1(x)}(x+y)} + H(-x+y)e^{\sqrt{-\mu_1(x)}(x-y)} + H(x+y)H(x-y)\frac{\cosh(\sqrt{-\mu_1(x)}y)}{\cosh(\sqrt{-\mu_1(x)}x)},$$

where  $H(\cdot)$  is the Heaviside function (with  $H(0) = \frac{1}{2}$ ). Now, one easily checks that

$$0 \leqslant \frac{\cosh(\sqrt{-\mu_{1}(x)}y)}{\cosh(\sqrt{-\mu_{1}(x)}x)} \leqslant e^{-\sqrt{-\mu_{1}(x)}(x+y)} + e^{\sqrt{-\mu_{1}(x)}(y-x)},$$

so that

 $H(-x-y) e^{\sqrt{-\mu_1(x)}(x+y)} + H(-x+y) e^{\sqrt{-\mu_1(x)}(x-y)}$  $\leqslant \psi_{x,1}(y) \leqslant e^{-\sqrt{-\mu_1(x)}|x+y|} + e^{-\sqrt{-\mu_1(x)}|y-x|},$ 

and therefore there exist positive constants c, C, independent of x, such that

$$0 < c \leq \left\| \psi_{x,1} \right\|_{L^2(\mathbb{R}_y)} \leq C < \infty.$$

In the same way, one can check the following estimates:

$$\begin{aligned} |\partial_x (e^{\sqrt{-\mu_1(x)}(x+y)})| &\leq (|(\sqrt{-\mu_1})'(x)||x+y| + \sqrt{-\mu_1(x)}) e^{-\sqrt{-\mu_1(x)}|x+y|}, \forall y \leq -x \\ |\partial_x (e^{\sqrt{-\mu_1(x)}(x-y)})| &\leq (|(\sqrt{-\mu_1})'(x)||x-y| + \sqrt{-\mu_1(x)}) e^{-\sqrt{-\mu_1(x)}|x-y|}, \forall y \geq x \end{aligned}$$

and, for  $y \in [-x, x]$ ,

$$\left|\partial_x \left(\frac{\cosh(\sqrt{-\mu_1(x)}y)}{\cosh(\sqrt{-\mu_1(x)x})}\right)\right| \le (e^{-\sqrt{-\mu_1(x)}(x+y)} + e^{\sqrt{-\mu_1(x)}(y-x)})(2x|(\sqrt{-\mu_1})'(x)| + \sqrt{-\mu_1(x)}).$$

Therefore, we deduce that there exists a positive constant, C', independent of x, such that

$$\left\|\partial_x\psi_{x,1}\right\|_{L^2(\mathbb{R}_y)}^2\leqslant C'<\infty.$$

By definition,  $u_x(y) = \frac{\psi_{x,1}(y)}{\|\psi_{x,1}\|_{L^2(\mathbb{R}_y)}}$ . It follows by elementary computations that

$$R(x) \leq 4 \frac{\|\partial_x \psi_{x,1}\|_{L^2(\mathbb{R}_y)}^2}{\|\psi_{x,1}\|_{L^2(\mathbb{R}_y)}^2},$$

and the point 6 is proved.

Finally, one obtains the point 7 by remarking

$$\|\partial_{y}u_{x}\|_{L^{2}(\mathbb{R}_{y})}^{2} = \mu_{1}(x) + |u_{x}(x)|^{2} + |u_{x}(-x)|^{2} = \mu_{1}(x) + \frac{2}{\|\psi_{x,1}\|_{L^{2}(\mathbb{R}_{y})}^{2}}.$$

As a direct application of proposition 2.3, we have

**Proposition 2.4.** For all  $\psi \in H^1(\mathbb{R}^2)$  and for all h > 0, we have:

$$\mathfrak{Q}_{h}(\psi) \geq \int_{\mathbb{R}_{x}} \int_{\mathbb{R}_{y}} (h^{2} |\partial_{x}\psi|^{2} + \tilde{\mu}_{1}(x)|\psi|^{2}) \,\mathrm{d}y \,\mathrm{d}x,$$

where  $\tilde{\mu}_1(x) = \mu_1(x)$ , for  $x \ge 0$  and 0 elsewhere. In particular, we have:

$$\mathfrak{Q}_h(\psi) \ge - \|\psi\|^2$$

or equivalently, for all  $\theta \in (0, \frac{\pi}{2})$ :

$$\mathcal{Q}_{\theta,\alpha}(\psi) \geqslant -\frac{\alpha^2}{\cos^2 \theta} \|\psi\|^2.$$

**Proof.** For  $\psi \in H^1(\mathbb{R}^2)$ , we have:

$$\mathfrak{Q}_h(\psi) = \int_{\mathbb{R}^2} h^2 |\partial_x \psi|^2 + |\partial_y \psi|^2 \, \mathrm{d}x \, \mathrm{d}y - \int_{\mathbb{R}} |\psi(|s|, s)|^2 \, \mathrm{d}s$$

so that:

$$\begin{split} \mathfrak{Q}_h(\psi) &= \int_{x \in \mathbb{R}^+} \left( \int_{\mathbb{R}_y} h^2 |\partial_x \psi|^2 \, \mathrm{d}y + \int_{\mathbb{R}_y} |\partial_y \psi|^2 \, \mathrm{d}y - |\psi(-x,x)|^2 - |\psi(x,x)|^2 \right) \, \mathrm{d}x \\ &+ \int_{x \in \mathbb{R}^-} \int_{\mathbb{R}_y} h^2 |\partial_x \psi|^2 + |\partial_y \psi|^2 \, \mathrm{d}y \, \mathrm{d}x. \end{split}$$

We infer that:

$$\mathfrak{Q}_{h}(\psi) \geq \int_{x \in \mathbb{R}^{+}} \int_{\mathbb{R}_{y}} (h^{2} |\partial_{x}\psi|^{2} + \mu_{1}(x)|\psi|^{2}) \,\mathrm{d}y \,\mathrm{d}x + \int_{x \in \mathbb{R}^{-}} \int_{\mathbb{R}_{y}} h^{2} |\partial_{x}\psi|^{2} \,\mathrm{d}y \,\mathrm{d}x,$$
  
sions follow.

and the conclusions follow.

# 3. Spectral reductions

Now we would like to use the spectral theory of  $\mathfrak{D}_x$  in order to compare the operator  $\mathfrak{H}_h$  with simpler operators.

### 3.1. Dimensional reduction

In order to deal with the singularity at x = 0, we introduce the following extension of  $u_x$ .

Notation 3.1. Let us define

$$\tilde{u}_x(y) = \begin{cases} u_x(y) & \text{if } x \ge 0\\ u_0(y) & \text{if } x < 0. \end{cases}$$

We also introduce the projections defined for  $\psi \in L^2(\mathbb{R}^2)$  by

$$\Pi \psi(x, y) = \langle \psi, \tilde{u}_x \rangle_{L^2(\mathbb{R}_v)} \tilde{u}_x(y), \quad \Pi^\perp \psi(x, y) = \psi(x, y) - \Pi \psi(x, y).$$

If  $\varphi = \varphi(x, y)$ , we denote  $\varphi_x(y) = \varphi(x, y)$ .

**Lemma 3.2.** For all  $\psi \in \text{Dom}(\mathfrak{Q}_h)$ , the function  $\Pi \psi$  belongs to  $\text{Dom}(\mathfrak{Q}_h)$  and we have

 $\mathcal{Q}_{h}(\Pi \psi) = \int_{\mathbb{R}_{x}} h^{2} |f'(x)|^{2} + (\hat{\mu}_{1}(x) + h^{2} \tilde{R}(x)) |f(x)|^{2} dx, \quad \text{with } f(x) = \langle \psi, \tilde{u}_{x} \rangle_{L^{2}(\mathbb{R}_{y})},$ where  $\hat{\mu}_{1}(x) = \mu_{1}(x)$  for  $x \ge 0$  and  $\hat{\mu}_{1}(x) = 1$  for x < 0 and  $\tilde{R}(x) = R(x)$  for x > 0 and  $\tilde{R}(x) = 0$  for  $x \le 0$ .

**Proof.** Recall that  $\mathsf{Dom}(\mathfrak{Q}_h) = H^1(\mathbb{R}^2)$ . By proposition 2.3, one has

$$\operatorname{ess\,sup}_{x\in\mathbb{R}} \|\partial_x \tilde{u}_x\|_{L^2(\mathbb{R}_y)}^2 = \sup_{x>0} R(x) < \infty$$

and

$$\operatorname{ess\,sup}_{x\in\mathbb{R}} \|\partial_y \tilde{u}_x\|_{L^2(\mathbb{R}_y)}^2 = \sup_{x\geqslant 0} \|\partial_y \tilde{u}_x\|_{L^2(\mathbb{R}_y)}^2 < \infty.$$

It follows immediately that, for any  $\psi \in H^1(\mathbb{R}^2)$ ,

$$\partial_x \big( \Pi \psi \big) = f(x) \partial_x \tilde{u}_x(y) + f'(x) \tilde{u}_x(y) \in L^2(\mathbb{R}^2),$$

since  $\operatorname{ess\,sup}_{x\in\mathbb{R}} f'(x) \leq \operatorname{ess\,sup}_{x\in\mathbb{R}} \langle \psi, \partial_x \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)} + \operatorname{ess\,sup}_{x\in\mathbb{R}} \langle \partial_x \psi, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)} < \infty$ , and  $\partial (\Pi \psi) - f(x) \partial \tilde{\mu}(y) \in L^2(\mathbb{R}^2)$ 

$$\partial_y (\Pi \psi) = f(x) \partial_y \tilde{u}_x(y) \in L^2(\mathbb{R}^2)$$

Thus one has  $\Pi \psi \in H^1(\mathbb{R}^2) = \mathsf{Dom}(\mathfrak{Q}_h)$ , and the calculations thereafter are valid. By definition, one has

$$\begin{split} \mathfrak{Q}_{h}(\Pi\psi) &= \int_{\mathbb{R}^{2}} h^{2} |f(x)\partial_{x}\tilde{u}_{x}(y) + f'(x)\tilde{u}_{x}(y)|^{2} + |f(x)|^{2} |\partial_{y}\tilde{u}_{x}(y)|^{2} \, dx \, dy - \int_{\mathbb{R}} |f(|s|)\tilde{u}_{|s|}(s)|^{2} \, ds \\ &= \int_{\mathbb{R}_{x}} h^{2} |f'(x)|^{2} + h^{2} |f(x)|^{2} \|\partial_{x}\tilde{u}_{x}(y)\|_{L^{2}(\mathbb{R}_{y})}^{2} \, dx + \int_{\mathbb{R}_{x}^{-}} |f(x)|^{2} \int_{\mathbb{R}_{y}} |\partial_{y}\tilde{u}_{x}(y)|^{2} \, dy \, dx \\ &+ \int_{\mathbb{R}_{x}^{+}} |f(x)|^{2} \int_{\mathbb{R}_{y}} |\partial_{y}\tilde{u}_{x}(y)|^{2} - \left(|\tilde{u}_{x}(-x)|^{2} + |\tilde{u}_{x}(x)|^{2}\right) \, dx \\ &= \int_{\mathbb{R}_{x}} h^{2} |f'(x)|^{2} \, dx + \int_{\mathbb{R}_{x}^{+}} h^{2} |f(x)|^{2} R(x) \, dx + \int_{\mathbb{R}_{x}^{-}} |f(x)|^{2} \, dx + \int_{\mathbb{R}_{x}^{+}} |f(x)|^{2} \mu_{1}(x) \, dx, \end{split}$$

where we used Fubini's theorem, and the following properties on  $\tilde{u}_x(y)$ :

•  $\forall x \in \mathbb{R}, \tilde{u}_x$  is normalized in  $L^2(\mathbb{R}_y)$ , and in particular, for any  $x \neq 0$ ,

$$2\langle \tilde{u}_x, \, \partial_x \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)} = \frac{\mathrm{d}}{\mathrm{d}x} \langle \tilde{u}_x, \, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)} = 0.$$

- $\forall x > 0$ , one has  $\mathfrak{q}_x(\tilde{u}_x) = \mu_1(x)$ .  $\forall x \leq 0$ , one has  $\int_{\mathbb{R}_y} |\partial_y \tilde{u}_x(y)|^2 dy = \int_{\mathbb{R}_y} |\partial_y u_0(y)|^2 dy = 1$ .

The result is now straightforward.

We get the same result for the corresponding bilinear form  $\mathfrak{B}_h$ .

**Lemma 3.3.** For all  $\psi_1, \psi_2 \in \mathsf{Dom}(\mathfrak{Q}_h)$ , we have

$$\mathfrak{B}_{h}(\Pi\psi_{1},\Pi\psi_{2}) = \int_{\mathbb{R}_{x}} h^{2} f_{1}'(x) f_{2}'(x) + (\hat{\mu}_{1}(x) + h^{2}\tilde{R}(x)) f_{1}(x) f_{2}(x) \,\mathrm{d}x,$$

with  $f_i(x) = \langle \psi_j, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)}$ .

Let us now use the orthogonal decomposition to bound  $\mathfrak{Q}_h$  from below.

**Proposition 3.4.** For all  $\psi \in \text{Dom}(\mathfrak{Q}_h)$  and all  $\varepsilon \in (0, 1)$ , we have

$$\begin{aligned} \mathfrak{Q}_{h}(\psi) &\geq \int_{\mathbb{R}_{x}} (1-\varepsilon)h^{2} |f'(x)|^{2} + (\tilde{\mu}_{1}(x) - 4\varepsilon^{-1}h^{2}\tilde{R}(x))|f(x)|^{2} \,\mathrm{d}x \\ &+ \int_{\mathbb{R}_{x}} (1-\varepsilon)h^{2} \|\partial_{x}\Pi^{\perp}\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} + (\tilde{\mu}_{2}(x) - 4\varepsilon^{-1}h^{2}\tilde{R}(x))\|\Pi^{\perp}\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} \,\mathrm{d}x, \end{aligned}$$

where  $\tilde{\mu}_i(x) = \mu_i(x)$  for  $x \ge 0$  and  $\tilde{\mu}_i(x) = 0$  for x < 0 ( $i \in \{1, 2\}$ );  $\tilde{R}(x) = R(x)$  for x > 0and  $\tilde{R}(x) = 0$  for  $x \leq 0$ .

**Proof.** By definition, one has for any  $\psi \in \mathsf{Dom}(\mathfrak{Q}_h) = H^1(\mathbb{R}^2)$ ,

$$\mathfrak{Q}_h(\psi) = \int_{\mathbb{R}^2} h^2 |\partial_x \psi|^2 \, \mathrm{d}x \, \mathrm{d}y + \int_{\mathbb{R}^-_x \times \mathbb{R}^+_y} |\partial_y \psi|^2 \, \mathrm{d}x \, \mathrm{d}y + \int_{\mathbb{R}^+_x} \mathfrak{q}_x(\psi_x) \, \mathrm{d}x$$

Since  $\psi \in \mathsf{Dom}(\mathfrak{Q}_h) = H^1(\mathbb{R}^2)$ , one has  $\Pi \psi \in H^1(\mathbb{R}^2)$  and  $\Pi^{\perp} \psi = \psi - \Pi \psi \in H^1(\mathbb{R}^2)$ . Moreover, for any fixed  $x \ge 0$ , recall that  $u_x$  is an eigenfunction corresponding to an eigenvalue of  $\mathfrak{D}_x$ , thus one has

$$\forall x \ge 0, \quad \mathfrak{q}_x(\psi_x) = \mathfrak{q}_x((\Pi\psi)_x) + \mathfrak{q}_x((\Pi^{\perp}\psi)_x) \ge \mu_1(x) \|\Pi\psi\|_{L^2(\mathbb{R}_v)}^2 + \mu_2(x) \|\Pi^{\perp}\psi\|_{L^2(\mathbb{R}_v)}^2$$

where we have applied the min-max principle to the quadratic form  $q_x$  and to the functions  $(\Pi \psi)_x$  and  $(\Pi^{\perp} \psi)_x$  which are orthogonal in  $L^2(\mathbb{R}_y)$ .

Now, one has  $\langle \Pi \varphi, \Pi^{\perp} \varphi \rangle_{L^2(\mathbb{R}_{\gamma})} = 0$ , for any  $\varphi \in L^2(\mathbb{R}^2)$ , therefore

$$\begin{split} \|\partial_{x}\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} &= \|\Pi\partial_{x}\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} + \|\Pi^{\perp}\partial_{x}\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} \\ &= \|\partial_{x}(\Pi\psi) - \mathcal{R}(x,y)\|_{L^{2}(\mathbb{R}_{y})}^{2} + \|\partial_{x}(\Pi^{\perp}\psi) + \mathcal{R}(x,y)\|_{L^{2}(\mathbb{R}_{y})}^{2}, \end{split}$$

with

$$\mathcal{R}(x, y) := [\partial_x, \Pi] \psi = \langle \psi, \partial_x \tilde{u}_x \rangle_{L^2(\mathbb{R}_v)} \tilde{u}_x(y) + \langle \psi, \tilde{u}_x \rangle_{L^2(\mathbb{R}_v)} \partial_x \tilde{u}_x(y).$$

It follows, for all  $\varepsilon \in (0, 1)$ ,

$$\begin{aligned} \|\partial_x \psi\|_{L^2(\mathbb{R}_y)}^2 &\ge (1-\varepsilon) \|\partial_x (\Pi \psi)\|_{L^2(\mathbb{R}_y)}^2 + (1-\varepsilon) \|\partial_x (\Pi^\perp \psi)\|_{L^2(\mathbb{R}_y)}^2 \\ &- 2(\varepsilon^{-1}-1) \|\mathcal{R}(x,y)\|_{L^2(\mathbb{R}_y)}^2. \end{aligned}$$

Now, notice, for any x > 0,

 $\|\mathcal{R}(x,y)\|_{L^{2}(\mathbb{R}_{y})}^{2} = \langle \psi, \partial_{x}\tilde{u}_{x} \rangle_{L^{2}(\mathbb{R}_{y})}^{2} + \langle \psi, \tilde{u}_{x} \rangle_{L^{2}(\mathbb{R}_{y})}^{2} \|\partial_{x}\tilde{u}_{x}\|_{L^{2}(\mathbb{R}_{y})}^{2} \leqslant 2 R(x) \|\psi_{x}\|_{L^{2}(\mathbb{R}_{y})}^{2},$ where we used proposition 2.3; and for any x < 0,  $\|\mathcal{R}(x, y)\|_{L^2(\mathbb{R}_+)}^2 \equiv 0$ .

Altogether, we proved

$$\begin{aligned} \mathfrak{Q}_{h}(\psi) &\geq \int_{\mathbb{R}_{x}} (1-\varepsilon)h^{2} \left( \|\partial_{x}(\Pi\psi)\|_{L^{2}(\mathbb{R}_{y})}^{2} + \|\partial_{x}(\Pi^{\perp}\psi)\|_{L^{2}(\mathbb{R}_{y})}^{2} \right) \mathrm{d}x \\ &+ \int_{x \geq 0} -4\varepsilon^{-1}h^{2}R(x)\|\psi_{x}\|_{L^{2}(\mathbb{R}_{y})}^{2} + \mu_{1}(x)\|\Pi\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} + \mu_{2}(x)\|\Pi^{\perp}\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} \mathrm{d}x, \end{aligned}$$

and the proof of proposition 3.4 is complete since  $\langle \tilde{u}_x, \partial_x \tilde{u}_x \rangle_{L^2(\mathbb{R}_v)} = 0$  yields

$$\|\partial_{x}(\Pi\psi)\|_{L^{2}(\mathbb{R}_{y})}^{2} = |f'(x)|^{2} + |f(x)|^{2} \|\partial_{x}\tilde{u}_{x}(y)\|_{L^{2}(\mathbb{R}_{y})}^{2} \ge |f'(x)|^{2}.$$

#### 3.2. Reduction to model operators

The aim of this section is to prove the following proposition.

**Proposition 3.5.** For all  $f \in H^1(\mathbb{R})$ , we let

$$\mathfrak{Q}_{h}^{\mathsf{modl}}(f) = \int_{\mathbb{R}} h^{2} |f'(x)|^{2} + \hat{\mu}_{1}(x) |f(x)|^{2} \, \mathrm{d}x,$$
$$\mathfrak{Q}_{h}^{\mathsf{mod2}}(f) = \int_{\mathbb{R}} h^{2} |f'(x)|^{2} + \tilde{\mu}_{1}(x) |f(x)|^{2} \, \mathrm{d}x,$$

and we denote by  $\mathfrak{H}_h^{\mathsf{mod}_j}$  the corresponding Friedrichs extensions. Set M' > M, where we denote

$$M = \sup_{x>0} R(x) = \sup_{x>0} \|\partial_x u_x\|_{L^2(\mathbb{R}_y)}^2$$

bounded by proposition 2.3. Then there exists  $M_0$ ,  $h_0 > 0$  such that for all  $h \in (0, h_0)$  and all  $C_h \ge M_0 h$ :

$$\mathcal{N}\left(\mathfrak{H}_{h}^{\mathsf{mod}1}, -\frac{1}{4} - C_{h} - h^{2}M\right) \leqslant \mathcal{N}\left(\mathfrak{H}_{h}, -\frac{1}{4} - C_{h}\right)$$
$$\leqslant \mathcal{N}\left(\mathfrak{H}_{h}^{\mathsf{mod}2}, \frac{-\frac{1}{4} - C_{h}}{1 - h} + (4M' + 1)h\right)$$

and

$$(1-h)\left\{\lambda_n^{\mathsf{mod2}}(h) - (4M'+1)h\right\} \leqslant \lambda_n(h) \leqslant \lambda_n^{\mathsf{mod1}}(h) + h^2 M.$$

**Remark 3.6.**  $M_0$  must be such that  $M_0 > 4M$  and  $\frac{-\frac{1}{4}-C_h}{1-h} + (4M'+1)h < \frac{-1}{4}$ , therefore one can chose  $M_0 = 4M' + \frac{3}{4}$ .

Let us now deal with the proof of proposition 3.5. Lemma 3.2 suggests we introduce the following reduced operator.

**Notation 3.7.** For all  $f \in H^1(\mathbb{R})$ , we let

$$\mathfrak{Q}_{h}^{\mathsf{red1}}(f) = \int_{\mathbb{R}_{x}} h^{2} |f'(x)|^{2} + (\hat{\mu}_{1}(x) + h^{2} \tilde{R}(x)) |f(x)|^{2} dx$$

and we denote by  $\mathfrak{H}_h^{\mathsf{red1}}$  the corresponding Friedrichs extension. We define  $(\lambda_n^{\mathsf{red1}}(h), f_n^{\mathsf{red1}})$  the nth  $L^2$ -normalized eigenpair which exists at least for  $n \in \{1, \ldots, \mathcal{N}(\mathfrak{H}_h^{\mathsf{red1}}, -\frac{1}{4})\}$ .

**Proposition 3.8.** For all  $n \in \{1, ..., \mathcal{N}(\mathfrak{H}_h^{\mathsf{red1}}, E)\}$ , with  $E < -\frac{1}{4}$ , and all h > 0 the nth eigenvalue of  $\mathfrak{H}_h$  exists and satisfies:

$$\lambda_n(h) \leqslant \lambda_n^{\mathsf{red1}}(h).$$

In particular, we have

$$\mathcal{N}(\mathfrak{H}_h, E) \geq \mathcal{N}(\mathfrak{H}_h^{\mathsf{red1}}, E).$$

**Proof.** The proof relies on the introduction of suitable test functions. For any  $n \in \{1, \ldots, \mathcal{N}(\mathfrak{H}_h^{\mathsf{red}}, E)\}$ , let us introduce the *n*-dimensional span

$$F_n = \operatorname{span}_{j \in \{1, \dots, n\}} f_j^{\mathsf{red1}}(x) \tilde{u}_x(y).$$

For all  $\psi \in F_n$  we have, with lemma 3.3 and noticing that the  $f_j^{\text{red1}}$  are orthogonal for the bilinear form associated with  $\mathfrak{Q}_h^{\text{red1}}$ ,

$$\mathfrak{Q}_h(\psi) \leqslant \lambda_n^{\mathsf{red1}}(h) \|\psi\|^2.$$

The conclusion follows from the min–max principle and the fact that  $-\frac{1}{4(1+h^2)} > -\frac{1}{4}$ .

We shall now analyze the reverse inequality. This is the aim of the following proposition.

Proposition 3.9. Let us consider the following quadratic form, defined on the product  $H^1(\mathbb{R}) \times H^1(\mathbb{R}^2)$ , by

$$\begin{aligned} \mathfrak{Q}_{h}^{\mathsf{tens}}(f,\varphi) &= \int_{\mathbb{R}_{x}} (1-h)h^{2} |f'(x)|^{2} + (\tilde{\mu}_{1}(x) - 4Mh)|f(x)|^{2} \, \mathrm{d}x \\ &+ \int_{\mathbb{R}^{2}} (1-h)h^{2} |\partial_{x}\varphi|^{2} + (\tilde{\mu}_{2}(x) - 4Mh)|\varphi|^{2} \, \mathrm{d}x \, \mathrm{d}y, \quad \forall (f,\varphi) \in H^{1}(\mathbb{R}) \times H^{1}(\mathbb{R}^{2}). \end{aligned}$$

If  $\mathfrak{H}_h^{\mathsf{tens}}$  denotes the associated operator, then we have, for all  $n \ge 1$ 

$$\lambda_n(h) \geqslant \lambda_n^{\mathsf{tens}}(h).$$

**Proof.** We use proposition 3.4 with  $\varepsilon = h$  and we get, for all  $\psi \in \mathsf{Dom}(\mathfrak{Q}_h)$ ,

$$\begin{aligned} \mathfrak{Q}_{h}(\psi) &\ge \int_{\mathbb{R}_{x}} (1-h)h^{2} |f'|^{2} + \left(\tilde{\mu}_{1}(x) - 4Mh\right) |f|^{2} \, \mathrm{d}x \\ &+ \int_{\mathbb{R}^{2}} (1-h)h^{2} |\partial_{x}\Pi^{\perp}\psi|^{2} + \left(\tilde{\mu}_{2}(x) - 4Mh\right) |\Pi^{\perp}\psi|^{2} \, \mathrm{d}x \, \mathrm{d}y. \end{aligned}$$
Thus we have

Thus we have

$$\mathfrak{Q}_{h}(\psi) \geqslant \mathfrak{Q}_{h}^{\mathsf{tens}}(\langle \psi, \tilde{u}_{x} \rangle_{L^{2}(\mathbb{R}_{y})}, \Pi^{\perp}\psi), \quad \|\psi\|^{2} = \|f\|_{L^{2}(\mathbb{R})}^{2} + \|\Pi^{\perp}\psi\|^{2}.$$
(3.1)

With notation 1.2 and (3.1) we infer

$$\lambda_n(h) \ge \inf_{\substack{G \subset H^1(\mathbb{R}^2) \\ \dim G = n}} \sup_{\psi \in G} \frac{\mathfrak{Q}_h^{\mathsf{tens}}(\langle \psi, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)}, \Pi^{\perp} \psi)}{\|\Pi \psi\|^2 + \|\Pi^{\perp} \psi\|^2}$$

Now, we define the linear injection

$$\mathcal{J}:\begin{cases} H^1(\mathbb{R}^2) & \to & H^1(\mathbb{R}) \times H^1(\mathbb{R}^2) \\ \psi & \mapsto & (\langle \psi, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)}, \Pi^{\perp} \psi). \end{cases}$$

So that we have

$$\inf_{\substack{G \subset H^1(\mathbb{R}^2) \ \psi \in G \\ \dim G = n}} \sup_{\substack{\mathcal{Q}_h^{\mathsf{tens}}(\langle \psi, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)}, \Pi^{\perp} \psi \|^2 \\ \|\Pi \psi\|^2 + \|\Pi^{\perp} \psi\|^2}} = \inf_{\substack{\tilde{G} \subset \mathcal{J}(H^1(\mathbb{R}^2)) \\ \dim \tilde{G} = n}} \sup_{\substack{\mathcal{Q}_h^{\mathsf{tens}}(f, \varphi) \\ \|f\|_{L^2(\mathbb{R})}^2 + \|\varphi\|^2}}$$

and

$$\inf_{\substack{\tilde{G} \subset \mathcal{J}(H^1(\mathbb{R}^2))\\\dim \tilde{G}=n}} \sup_{\substack{(f,\varphi) \in \tilde{G}}} \frac{\mathfrak{Q}_h^{\mathsf{tens}}(f,\varphi)}{\|f\|_{L^2(\mathbb{R})}^2 + \|\varphi\|^2} \ge \inf_{\substack{\tilde{G} \subset H^1(\mathbb{R}) \times H^1(\mathbb{R}^2)\\\dim \tilde{G}=n}} \sup_{\substack{(f,\varphi) \in \tilde{G}}} \frac{\mathfrak{Q}_h^{\mathsf{tens}}(f,\varphi)}{\|f\|_{L^2(\mathbb{R})}^2 + \|\varphi\|^2}$$

We recognize the *n*th Rayleigh quotient of  $\mathfrak{H}_{h}^{\mathsf{tens}}$  and the conclusion follows.

**Notation 3.10.** For all  $f \in H^1(\mathbb{R})$ , we let

$$\mathfrak{Q}_{h}^{\mathsf{red2}}(f) = \int_{\mathbb{R}} (1-h)h^{2} |f'(x)|^{2} + \left(\tilde{\mu}_{1}(x) - 4Mh\right) |f(x)|^{2} \,\mathrm{d}x$$

and we denote by  $\mathfrak{H}_h^{red2}$  the corresponding Friedrichs extension.

**Proposition 3.11.** For any h > 0 and  $C_h > 4Mh$ , one has

$$\lambda_n(h) \geqslant \lambda_n^{\mathsf{red2}}(h), \quad \forall n \in \left\{1, \dots, \mathcal{N}\left(\mathfrak{H}_h, -\frac{1}{4} - C_h\right)\right\}$$

and

$$\mathcal{N}(\mathfrak{H}_h, -\frac{1}{4} - C_h) \leqslant \mathcal{N}(\mathfrak{H}_h^{\mathsf{red2}}, -\frac{1}{4} - C_h).$$

V Duchêne and N Raymond

**Proof.** Notice that for any  $\varphi \in H^1(\mathbb{R}^2)$ , one has

$$\int_{\mathbb{R}^2} (1-h)h^2 |\partial_x \varphi|^2 + \left(\tilde{\mu}_2(x) - 4Mh\right)|\varphi|^2 \,\mathrm{d}x \,\mathrm{d}y > \left(-\frac{1}{4} - 4Mh\right) \|\varphi\|^2.$$

It follows that for any eigenstate of  $\mathfrak{H}_h^{\text{tens}}$  below the threshold  $(-\frac{1}{4} - 4Mh)$  is of the form (f, 0), with f an eigenstate of  $\mathfrak{H}_h^{\mathsf{red2}}$ . In other words, one has for any  $C \ge 4M$ ,

$$\{\lambda \in \sigma(\mathfrak{H}_h^{\mathsf{tens}}) : \lambda \leqslant -\frac{1}{4} - Ch\} = \{\lambda \in \sigma_{\mathsf{dis}}(\mathfrak{H}_h^{\mathsf{red2}}) : \lambda \leqslant -\frac{1}{4} - Ch\},$$
  
now follows from proposition 3.9.

and the result now follows from proposition 3.9.

Proposition 3.5 is a direct consequence of propositions 3.8 and 3.11, and straightforward computations. In particular, we use

$$\begin{aligned} \mathfrak{Q}_{h}^{\mathsf{red2}}(f) &= (1-h) \int_{\mathbb{R}} h^{2} |f'(x)|^{2} + \frac{\tilde{\mu}_{1}(x) - 4Mh}{1-h} |f(x)|^{2} \, \mathrm{d}x \\ &\geq (1-h) \int_{\mathbb{R}} h^{2} |f'(x)|^{2} + (\tilde{\mu}_{1}(x)(1+h) - 4Mh - Ch^{2}) |f(x)|^{2} \, \mathrm{d}x \\ &\geq (1-h) \int_{\mathbb{R}} h^{2} |f'(x)|^{2} + (\tilde{\mu}_{1}(x) - (4M'+1)h) |f(x)|^{2} \, \mathrm{d}x, \end{aligned}$$

which is valid for  $h \in (0, h_0)$  with  $h_0$  sufficiently small, C sufficiently large, and any M' > M(the last inequality comes from proposition 2.3, item 3). It follows

$$\mathcal{N}\left(\mathfrak{H}_{h}^{\mathsf{red2}}, -\frac{1}{4} - C_{h}\right) \leqslant \mathcal{N}\left(\mathfrak{H}_{h}^{\mathsf{mod2}}, \frac{-\frac{1}{4} - C_{h}}{1 - h} + (4M' + 1)h\right)$$

and for any  $n \leq \mathcal{N}\left(\mathfrak{H}_{h}^{\mathsf{mod2}}, -\frac{1}{4} - C_{h}\right)$ ,

$$\lambda_n^{\mathsf{red2}}(h) \geqslant (1-h) \big\{ \lambda_n^{\mathsf{mod2}}(h) - (4M'+1)h \big\}.$$

The condition  $C_h \ge M_0 h > (4M' + \frac{3}{4})h$  ensures  $\frac{-\frac{1}{4} - C_h}{1 - h} + (4M' + 1)h < -\frac{1}{4}$ , thus the above quantities are well-defined.

## 4. Models in dimension 1

Thanks to section 3 we have reduced the spectral analysis of  $\mathfrak{H}_h$  to the investigation of one dimensional models. This section is devoted to the proofs of theorems 1.7 and 1.11.

#### 4.1. Number of bound states

In order to prove theorem 1.7 we need the following extended Weyl's asymptotics which is not completely standard (see remark 4.2).

**Proposition 4.1.** Let us consider  $V : \mathbb{R} \to \mathbb{R}$  a piecewise Lipschitzian function with a finite number of discontinuities satisfying:

(1) V tends to  $\ell_{\pm\infty}$  when  $x \to \pm\infty$  with  $\ell_{+\infty} \leq \ell_{-\infty}$ , (2)  $\sqrt{(\ell_{+\infty} - V)_+}$  belongs to  $L^1(\mathbb{R})$ .

*Consider the operator*  $\mathfrak{h}_h = -h^2 \partial_x^2 + V(x)$  and a function  $(0, 1) \ni h \mapsto E(h) \in (-\infty, \ell_{+\infty})$ such that one has:

(1) for any 
$$h \in (0, 1)$$
,  $\{x \in \mathbb{R} : V(x) \leq E(h)\} = [x_{\min}(E(h)), x_{\max}(E(h))],$   
(2)  $h^{1/3}(x_{\max}(E(h)) - x_{\min}(E(h))) \xrightarrow[h \to 0]{} 0,$   
(3)  $E(h) \xrightarrow[h \to 0]{} E_0 \leq \ell_{+\infty}.$ 

J. Phys. A: Math. Theor. 47 (2014) 155203

V Duchêne and N Raymond

Then we have:

$$\mathcal{N}(\mathfrak{h}_h, E(h)) \underset{h \to 0}{\sim} \frac{1}{\pi h} \int_{\mathbb{R}} \sqrt{(E_0 - V)_+} \, \mathrm{d}x.$$

Proof. The strategy of the proof is well-known but we recall it since the usual result does not deal with a moving threshold E(h). We consider a subdivision of the real axis  $(s_i(h^{\alpha}))_{i \in \mathbb{Z}}$ , which contains the discontinuities of V, and such that there exists c > 0, C > 0 for which, for all  $j \in \mathbb{Z}$  and h > 0,  $ch^{\alpha} \leq s_{j+1}(h^{\alpha}) - s_j(h^{\alpha}) \leq Ch^{\alpha}$ , where  $\alpha > 0$  is to be determined. We introduce

$$J_{\min}(h^{\alpha}) = \min\{j \in \mathbb{Z} : s_j(h^{\alpha}) \ge x_{\min}(E(h))\},\$$
  
$$J_{\max}(h^{\alpha}) = \max\{j \in \mathbb{Z} : s_j(h^{\alpha}) \le x_{\max}(E(h))\}.$$

For  $j \in \mathbb{Z}$  we may introduce the Dirichlet (resp. Neumann) realization on  $(s_j(h^{\alpha}), s_{j+1}(h^{\alpha}))$ of  $-h^2 \partial_x^2 + V(x)$  denoted by  $\mathfrak{h}_{h,j}^{\mathsf{Dir}}$  (resp.  $\mathfrak{h}_{h,j}^{\mathsf{Neu}}$ ). The so-called Dirichlet–Neumann bracketing (see [24, chapter XIII, section 15]) implies:

$$\sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} \mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Dir}}, E(h)) \leqslant \mathcal{N}(\mathfrak{h}_{h}, E(h)) \leqslant \sum_{j=J_{\min}(h^{\alpha})-1}^{J_{\max}(h^{\alpha})+1} \mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Neu}}, E(h))$$

Let us estimate  $\mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Dir}}, E(h))$ . If  $\mathfrak{q}_{h,j}^{\mathsf{Dir}}$  denotes the quadratic form of  $\mathfrak{h}_{h,j}^{\mathsf{Dir}}$ , we have:  $\mathfrak{c}^{s_{j+1}(h^{\alpha})}$ 

$$q_{h,j}^{\mathsf{Dir}}(\psi) \leq \int_{s_{j}(h^{\alpha})}^{s_{j+1}(n)} h^{2} |\psi'(x)|^{2} + V_{j,\sup,h} |\psi(x)|^{2} \, \mathrm{d}x, \quad \forall \psi \in \mathcal{C}_{0}^{\infty}((s_{j}(h^{\alpha}), s_{j+1}(h^{\alpha}))),$$
  
where

$$V_{j,\sup,h} = \sup_{x \in (s_j(h^{\alpha}), s_{j+1}(h^{\alpha}))} V(x).$$

We infer that

$$\mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Dir}}, E(h)) \ge \# \left\{ n \ge 1 : n \leqslant \frac{1}{\pi h} (s_{j+1}(h^{\alpha}) - s_j(h^{\alpha})) \sqrt{\left(E(h) - V_{j,\sup,h}\right)_+} \right\}$$
  
so that:

$$\mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Dir}}, E(h)) \ge \frac{1}{\pi h} (s_{j+1}(h^{\alpha}) - s_j(h^{\alpha})) \sqrt{(E(h) - V_{j,\sup,h})_+} - 1$$

and thus:

$$\sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} \mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Dir}}, E(h)) \ge \frac{1}{\pi h} \sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} (s_{j+1}(h^{\alpha}) - s_{j}(h^{\alpha})) \sqrt{(E(h) - V_{j,\sup,h})_{+}} - (J_{\max}(h^{\alpha}) - J_{\min}(h^{\alpha}) + 1).$$

Let us consider the function

$$f_h(x) = \sqrt{(E(h) - V(x))_+}$$

and analyze

$$\begin{aligned} \left| \sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} (s_{j+1}(h^{\alpha}) - s_{j}(h^{\alpha})) \sqrt{\left(E(h) - V_{j,\sup,h}\right)_{+}} - \int_{\mathbb{R}} f_{h}(x) \, \mathrm{d}x \right| \\ & \leqslant \left| \sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} \int_{s_{j}(h^{\alpha})}^{s_{j+1}(h^{\alpha})} \sqrt{\left(E(h) - V_{j,\sup,h}\right)_{+}} - f_{h}(x) \, \mathrm{d}x \right| \\ & + \int_{s_{J_{\max}}(h^{\alpha})}^{x_{\max}(E(h))} f_{h}(x) \, \mathrm{d}x + \int_{x_{\min}(E(h))}^{s_{J_{\min}(h^{\alpha})}} f_{h}(x) \, \mathrm{d}x \\ & \leqslant \left| \sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} \int_{s_{j}(h^{\alpha})}^{s_{j+1}(h^{\alpha})} \sqrt{\left(E(h) - V_{j,\sup,h}\right)_{+}} - f_{h}(x) \, \mathrm{d}x \right| + \tilde{C}h^{\alpha}. \end{aligned}$$

Using the trivial inequality  $|\sqrt{a_+} - \sqrt{b_+}| \leq \sqrt{|a-b|}$ , we notice that

$$|f_h(x) - \sqrt{\left(E(h) - V_{j,\sup,h}\right)_+}| \leq \sqrt{|V(x) - V_{j,\sup,h}|}.$$

Since *V* is Lipschitzian on  $(s_j(h^{\alpha}), s_{j+1}(h^{\alpha}))$ , we get:

$$\left|\sum_{j=J_{\min}(h^{\alpha})}^{J_{\max}(h^{\alpha})} \int_{s_{j}(h^{\alpha})}^{s_{j+1}(h^{\alpha})} \sqrt{(E(h)-V_{j,\sup,h})_{+}} - f_{h}(x) \, \mathrm{d}x\right| \leq (J_{\max}(h^{\alpha}) - J_{\min}(h^{\alpha}) + 1)\tilde{C}h^{\alpha}h^{\alpha/2}.$$

This leads to the optimal choice  $\alpha = \frac{2}{3}$  and we get the lower bound:

$$\sum_{j=J_{\min}(h^{2/3})}^{J_{\max}(h^{2/3})} \mathcal{N}(\mathfrak{h}_{h,j}^{\mathsf{Dir}}, E(h)) \ge \frac{1}{\pi h} \left( \int_{\mathbb{R}} f_h(x) \, \mathrm{d}x - \tilde{C}h(J_{\max}(h^{2/3}) - J_{\min}(h^{2/3}) + 1) - \tilde{C}h^{2/3} \right).$$

Therefore we infer

. ....

$$\mathcal{N}(\mathfrak{h}_h, E(h)) \geq \frac{1}{\pi h} \left( \int_{\mathbb{R}} f_h(x) \, \mathrm{d}x - \tilde{C} h^{1/3}(x_{\max}(E(h)) - x_{\min}(E(h)) - \tilde{C} h^{2/3} \right).$$

We notice that:  $f_h(x) \leq \sqrt{(\ell_{+\infty} - V(x))_+}$  so that we can apply the dominate convergence theorem. We can deal with the Neumann realizations in the same way.

**Remark 4.2.** Classical results (see [11, 24–26]) impose a fixed security distance below the edge of the essential spectrum  $(E(h) = E_0 < l_{+\infty})$  or deal with non-negative potentials, *V*, with compact support. Both these cases are recovered by proposition 4.1. In our result, the maximal threshold for which one can ensure that the semiclassical behavior of the counting function holds is dictated by the convergence rate of the potential towards its limit at infinity, through the assumption

$$h^{1/3}(x_{\max}(E(h)) - x_{\min}(E(h))) \xrightarrow[h \to 0]{} 0$$

More precisely, assume that  $l_{-\infty} > l_{+\infty}$  so that  $x_{\min}(E(h)) \ge x_{\min}(l_{+\infty})$  is uniformly bounded for E(h) in a neighborhood of  $l_{+\infty}$ . Then

- If  $l_{+\infty} V(x) \leq Cx^{-\gamma}$  for any  $x \geq x_0$  and given  $x_0, C > 0$  and  $\gamma > 2$ , then one can choose  $E(h) = l_{+\infty} Ch^{\rho}$  and  $x_{\max}(E(h)) \leq h^{-\rho/\gamma}$ , provided  $\rho < \gamma/3$ .
- If  $l_{+\infty} V(x) \leq C_1 \exp(-C_2 x)$  for any  $x \geq x_0$  and given  $x_0, C_1, C_2 > 0$ , then one can choose  $E(h) = l_{+\infty} C_1 \exp(C_2 h^{-1/3} \times o(h))$  and the assumption is satisfied.

**Proof of theorem 1.7.** In order to prove theorem 1.7 we apply proposition 3.5 with  $C_h = C(h)$ . Then we apply proposition 4.1 to the operators  $\mathfrak{H}_h^{\text{mod}j}$ . Increasing  $M_0$  if necessary, we have  $E(h) \leq -\frac{1}{4} - Ch$  with any C > 0 and therefore the assumptions of proposition 4.1 are satisfied. Indeed from proposition 2.3,  $\tilde{\mu}_1$  and  $\hat{\mu}_1$  converge exponentially to  $-\frac{1}{4}$  as  $x \to \infty$ , and  $\tilde{\mu}_1, \hat{\mu}_1 > -\frac{1}{4}$  for x < 0.

#### 4.2. Low lying spectrum

Let us now deal with the proofs of theorem 1.11 and 1.13.

4.2.1. Proof of theorem 1.11. The following proposition provides the asymptotics of the lowest eigenvalues of the models  $\mathfrak{H}_h^{\text{mod}j}$  and is a direct consequence of the analysis of [10, section 3].

**Proposition 4.3.** For j = 1, 2 and for all  $n \ge 1$  we have:  $\lambda_n^{\text{mod}j}(h) = -1 + 2^{2/3} z_{\text{Ai}}(n) h^{2/3} + O(h).$ 

With proposition 3.5 this implies theorem 1.11. In fact, it is possible to establish some localization properties of the first eigenfunctions of  $\mathfrak{H}_h$ .

**Proposition 4.4.** Let  $\lambda \in (-1, 0)$  and  $\delta \in (0, 1)$ . For all h > 0 and all eigenpairs  $(\lambda, \psi)$  of  $\mathfrak{H}_h$ , we have

$$\int_{\mathbb{R}_{y}} \int_{-\infty}^{0} e^{2(1-\delta)\sqrt{-\lambda}h^{-1}|x|} |\psi|^{2} \, \mathrm{d}x \, \mathrm{d}y \leqslant \frac{1}{(-\lambda)\delta^{2}} \|\psi\|^{2}.$$
(4.1)

Moreover, for all  $C_0 > 0$ , there exist  $h_0 > 0$ , C > 0 and  $\varepsilon_0 > 0$  such that for all  $h \in (0, h_0)$ and all eigenpairs  $(\lambda, \psi)$  such that  $\lambda \leq -1 + C_0 h^{2/3}$ , we have

$$\int_{\mathbb{R}_{y}} \int_{0}^{+\infty} e^{2\varepsilon_{0}h^{-2/3}|x|} |\psi|^{2} dx dy \leq C \|\psi\|^{2}.$$
(4.2)

**Proof.** This is a consequence of proposition 2.4 and of Agmon type estimates inherited from the one dimensional operator  $-h^2 - \partial_x^2 + \tilde{\mu}_1(x)$  (see [10] and also the original references [1, 17]).

4.2.2. Proof of theorem 1.13. Let us consider an eigenpair  $(\lambda, \psi)$  such that  $\lambda \leq -1 + C_0 h^{2/3}$ . We can write

$$\mathfrak{Q}_h(\psi) = \lambda \|\psi\|^2$$

and

$$\mathfrak{Q}_{h}(\psi) = \mathfrak{Q}_{h,+}(\psi) + \mathfrak{Q}_{h,-}(\psi),$$

where

$$\mathfrak{Q}_{h,-}(\psi) = \int_{\mathbb{R}^-_x \times \mathbb{R}_y} h^2 |\partial_x \psi|^2 + |\partial_y \psi|^2 \,\mathrm{d}x \,\mathrm{d}y$$

$$\mathfrak{Q}_{h,+}(\psi) = \int_{\mathbb{R}^+_x \times \mathbb{R}_y} h^2 |\partial_x \psi|^2 + |\partial_y \psi|^2 \, \mathrm{d}x \, \mathrm{d}y - \int_{\mathbb{R}^+} |\psi(x,x)|^2 \, \mathrm{d}x - \int_{\mathbb{R}^+} |\psi(x,-x)|^2 \, \mathrm{d}x.$$
  
We infer that

We infer that

$$\mathfrak{Q}_{h,+}(\psi) + \mathfrak{Q}_{h,-}(\psi) + \int_{\mathbb{R}^+_x \times \mathbb{R}_y} |\psi|^2 \, \mathrm{d}x \, \mathrm{d}y + \int_{\mathbb{R}^-_x \times \mathbb{R}_y} |\psi|^2 \, \mathrm{d}x \, \mathrm{d}y \leqslant C_0 h^{2/3} \|\psi\|^2 \, \mathrm{d}x \, \mathrm{d}y \leq C_0 h^{2/3} \|\psi\|^2 \, \mathrm{d}y \, \mathrm{d}$$

By point (5) of proposition 2.3, we deduce that

$$0 \leq \mathfrak{Q}_{h,-}(\psi) + \int_{\mathbb{R}^{-}_{x} \times \mathbb{R}_{y}} |\psi|^{2} \, \mathrm{d}x \, \mathrm{d}y \leq C_{0} h^{2/3} \|\psi\|^{2}$$
(4.3)

and

$$0 \leq \mathfrak{Q}_{h,+}(\psi) + \int_{\mathbb{R}^+_x \times \mathbb{R}_y} |\psi|^2 \, \mathrm{d}x \, \mathrm{d}y \leq C_0 h^{2/3} \|\psi\|^2.$$
(4.4)

We recall the points (1) and (4) of proposition 2.3 to deduce from (4.4) that

$$\int_{\mathbb{R}^+_x \times \mathbb{R}_y} h^2 |\partial_x \psi|^2 \, \mathrm{d}x \, \mathrm{d}y + \int_{\mathbb{R}^+} \mathfrak{q}_x(\psi_x) \, \mathrm{d}x - \int_{\mathbb{R}^+_x \times \mathbb{R}_y} \mu_1(x) |\psi|^2 \, \mathrm{d}x \, \mathrm{d}y \leqslant C_0 h^{2/3} \|\psi\|^2$$

where we recall that

$$\mathfrak{q}_x(\psi_x) = \int_{\mathbb{R}_y} |\partial_y \psi_x|^2 \,\mathrm{d}y - |\psi(x, x)|^2 - |\psi(x, -x)|^2$$

We have

$$\mathfrak{q}_{x}(\psi_{x}) - \mu_{1}(x) \|\psi\|_{L^{2}(\mathbb{R}_{y})}^{2} = \mathfrak{q}_{x}(\psi - \Pi\psi) - \mu_{1}(x) \|\psi - \Pi\psi\|_{L^{2}(\mathbb{R}_{y})}^{2}$$

and then, due to the min-max principle,

$$\mathfrak{q}_x(\psi_x - (\Pi\psi)_x) \ge \mu_2(x) \|\psi - \Pi\psi\|_{L^2(\mathbb{R}_y)}^2.$$

We get

$$\int_{\mathbb{R}^+_x \times \mathbb{R}_y} (\mu_2(x) - \mu_1(x)) |\psi - \Pi \psi|^2 \, \mathrm{d}x \, \mathrm{d}y \leqslant C_0 h^{2/3} \|\psi\|^2.$$

Due to the simplicity of  $\mu_1$ , we can find  $\varepsilon_0 > 0$  such that for  $x \in [0, 1]$  we have

$$\mu_2(x) - \mu_1(x) \geqslant \varepsilon_0.$$

Then for  $x \ge 1$  we use the estimates of Agmon (4.2) and the boundedness of the  $\mu_j$  to get

$$\int_{\mathbb{R}_{y}} \int_{x>1} (\mu_{2}(x) - \mu_{1}(x)) |\psi - \Pi \psi|^{2} \, \mathrm{d}x \, \mathrm{d}y \leq C \int_{\mathbb{R}_{y}} \int_{x>1} |\psi|^{2} \, \mathrm{d}x \, \mathrm{d}y \leq C \, \mathrm{e}^{-2\varepsilon_{0}h^{-2/3}} \|\psi\|^{2},$$

where we have used

$$\|\Pi^{\perp}\psi\|_{L^2(\mathbb{R}_y)}^2 \leqslant \|\psi\|_{L^2(\mathbb{R}_y)}^2.$$

We deduce that

$$\int_{\mathbb{R}^+_x \times \mathbb{R}_y} |\psi - \Pi \psi|^2 \, \mathrm{d} x \, \mathrm{d} y \leqslant C h^{2/3} \|\psi\|^2.$$

We have proved (it follows from the point (vi) of proposition 2.3) that the application  $[0, +\infty) \ni x \mapsto \Pi_x = \langle \cdot, \tilde{u}_x \rangle_{L^2(\mathbb{R}_y)} \tilde{u}_x \in \mathcal{L}_c(L^2(\mathbb{R}_y), L^2(\mathbb{R}_y))$  is Lipschitzian (with Lipschitz constant K > 0) so that

$$\|(\Pi_x - \Pi_0)\psi_x\|_{L^2(\mathbb{R}_y)} \leq K|x|\|\psi\|_{L^2(\mathbb{R}_y)}.$$

Let us now consider for instance  $\eta \in (0, \frac{1}{100})$ . We infer that

$$\int_{\mathbb{R}_{y}} \int_{0 < x < h^{2/3-\eta}} |\Pi_{x} \psi_{x} - \Pi_{0} \psi|^{2} \, \mathrm{d}x \, \mathrm{d}y \leqslant K^{2} h^{4/3-2\eta} \|\psi\|^{2}.$$

Thanks to the estimates of Agmon, we have

$$\int_{\mathbb{R}_{y}} \int_{x > h^{2/3-\eta}} |\Pi_{x} \psi_{x} - \Pi_{0} \psi|^{2} \, \mathrm{d}x \, \mathrm{d}y \leqslant C \, \mathrm{e}^{-2\varepsilon_{0} h^{-2/3}} \|\psi\|^{2}$$

We deduce that

$$\int_{\mathbb{R}^+_x \times \mathbb{R}_y} |\psi - \Pi_0 \psi|^2 \, \mathrm{d} x \, \mathrm{d} y \leqslant C h^{2/3} \|\psi\|^2$$

and, since

$$\int_{\mathbb{R}^-_x \times \mathbb{R}_y} |\psi - \Pi_0 \psi|^2 \, \mathrm{d}x \, \mathrm{d}y \leqslant \int_{\mathbb{R}^-_x \times \mathbb{R}_y} |\psi|^2 \, \mathrm{d}x \, \mathrm{d}y.$$

the conclusion follows from (4.3).

## Acknowledgments

The second author would like to thank K Pankrashkin who gave him the initial impulse to study the broken  $\delta$ -interactions.

#### References

- Agmon S 1982 Lectures on Exponential Decay of Solutions of Second-Order Elliptic Equations: Bounds on Eigenfunctions of N-Body Schrödinger Operators (Mathematical Notes vol 29) (Princeton, NJ: Princeton University Press)
- [2] Albeverio S, Gesztesy F, Høegh-Krohn R and Holden H 1988 Solvable Models in Quantum Mechanics (Texts and Monographs in Physics) (New York: Springer)
- [3] Balazard-Konlein A 1985 Asymptotique semi-classique du spectre pour des opérateurs à symbole opératoriel C. R. Acad. Sci., Paris Sér. I Math. 301 903-6
- [4] Berkolaiko G and Kuchment P 2013 Introduction to Quantum Graphs (Mathematical Surveys and Monographs vol 186) (Providence, RI: American Mathematical Society)
- [5] Born M and Oppenheimer R 1927 Zur Quantentheorie der Molekeln Ann. Phys. 84 457–84
- [6] Brasche J F, Exner P, Kuperin Y A and Šeba P 1994 Schrödinger operators with singular interactions J. Math. Anal. Appl. 184 112–39
- Brown B M, Eastham M S P and Wood I G 2008 An example on the discrete spectrum of a star graph Analysis on Graphs and its Applications (Proc. Symp. Pure Math. vol 77) (Providence, RI: American Mathematical Society) pp 331–5
- [8] Brown B M, Eastham M S P and Wood I G 2009 Estimates for the lowest eigenvalue of a star graph J. Math. Anal. Appl. 354 24–30
- [9] Dauge M, Lafranche Y and Raymond N 2012 Quantum waveguides with corners Actes du Congrès SMAI 2011: ESAIM Proc. (Les Ulis: EDP Sciences) vol 35, pp 14–45
- [10] Dauge M and Raymond N 2012 Plane waveguides with corners in the small angle limit J. Math. Phys. 53 123529
- [11] Dimassi M and Sjöstrand J 1999 Spectral Asymptotics in the Semi-Classical Limit (London Mathematical Society Lecture Note Series vol 268) (Cambridge: Cambridge University Press)
- [12] Exner P 2008 Leaky quantum graphs: a review Analysis on Graphs and its Applications (Proc. Symp. Pure Math. vol 77) (Providence, RI: American Mathematical Society) pp 523–64
- [13] Exner P and Ichinose T 2001 Geometrically induced spectrum in curved leaky wires J. Phys. A: Math. Gen. 34 1439–50
- [14] Exner P and Jex M 2013 Spectral asymptotics of a strong δ' interaction on a planar loop J. Phys.
   A: Math. Theor. 46 345201
- [15] Exner P and Němcová K 2001 Bound states in point-interaction star graphs J. Phys. A: Math. Gen. 34 7783–94
- [16] Exner P and Němcová K 2003 Leaky quantum graphs: approximations by point-interaction Hamiltonians J. Phys. A: Math. Gen. 36 10173–93
- [17] Helffer B 1988 Semi-Classical Analysis for the Schrödinger Operator and Applications (Lecture Notes in Mathematics vol 1336) (Berlin: Springer)
- [18] Klein M, Martinez A, Seiler R and Wang X P 1992 On the Born–Oppenheimer expansion for polyatomic molecules *Commun. Math. Phys.* 143 607–39
- [19] Kondej S 2013 Schrödinger operator with a strong varying interaction on a curve in ℝ<sup>2</sup> J. Math. Phys. 54 093511
- [20] Lotoreichik V 2013 Note on 2D Schrödinger operators with δ-interactions on angles and crossing lines Nanosystems: Phys. Chem. Math. 4 166–72
- [21] Martinez A 1989 Développements asymptotiques et effet tunnel dans l'approximation de Born-Oppenheimer Ann. Inst. Henri Poincaré Phys. Théor. 50 239–57

- [22] Morame A and Truc F 2005 Remarks on the spectrum of the Neumann problem with magnetic field in the half-space J. Math. Phys. 46 012105
- [23] Post O 2012 Spectral Analysis on Graph-Like Spaces (Lecture Notes in Mathematics vol 2039) (Heidelberg: Springer)
- [24] Reed M and Simon B 1978 Methods of Modern Mathematical Physics: IV. Analysis of Operators (New York: Academic)
- [25] Robert D 1987 Autour de l'approximation Semi-Classique (Progress in Mathematics vol 68) (Boston, MA: Birkhäuser)
- [26] Zworski M 2012 Semiclassical Analysis (Graduate Studies in Mathematics vol 138) (Providence, RI: American Mathematical Society)